

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Li, Meijian; Lappi, Tuomas; Zhao, Xingbo; Salgado, Carlos

Title: On the momentum broadening of in-medium jet evolution using a light-front Hamiltonian approach

Year: 2024

Version: Published version

Copyright: © 2024 the Authors

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Li, M., Lappi, T., Zhao, X., & Salgado, C. (2024). On the momentum broadening of in-medium jet evolution using a light-front Hamiltonian approach. In *HardProbes2023: 11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions (Article 151)*. Sissa Medialab. POS Proceedings of Science, 438. <https://doi.org/10.22323/1.438.0151>

On the momentum broadening of in-medium jet evolution using a light-front Hamiltonian approach

Meijian Li,^{a,*} Tuomas Lappi,^{b,c} Xingbo Zhao^{d,e} and Carlos A. Salgado^a

^a*Instituto Galego de Fisica de Altas Enerxias (IGFAE), Universidade de Santiago de Compostela, E-15782 Galicia, Spain*

^b*Department of Physics, P.O. Box 35, FI-40014 University of Jyväskylä, Finland*

^c*Helsinki Institute of Physics, P.O. Box 64, FI-00014 University of Helsinki, Finland*

^d*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

^e*University of Chinese Academy of Sciences, Beijing 100049, China*

E-mail: meijian.li@usc.es, tuomas.v.v.lappi@jyu.fi, xbzhao@impcas.ac.cn, carlos.salgado@usc.es

We have developed a non-perturbative light-front Hamiltonian formalism to simulate the real-time evolution of a quark state in a $SU(3)$ colored medium, with a series of works. In this proceeding article, we focus on the transverse momentum broadening of an in-medium quark jet. We perform the numerical simulation of the quark jet evolution in the $|q\rangle + |qg\rangle$ Fock space at various medium densities. By analyzing the resulting jet light-front wavefunction, we extract the gluon emission rate and the non-eikonal quenching parameter. Additionally, we provide the analytical derivation of the eikonal expectation value of the quark-gluon state's transverse momentum for any color configuration and arbitrary spatial distribution. This study can help understand jet momentum broadening beyond the eikonal limit.

HardProbes2023

26-31 March 2023

Aschaffenburg, Germany

*Speaker

1. Introduction

In heavy-ion collisions, energetic quarks and gluons are produced at early stages, propagating through the dense and hot medium. Similar processes happen in deeply inelastic scattering where quark and gluon jets traverse cold nuclear matter. We developed a non-perturbative computational method, the time-dependent Basis Light-Front Quantization (tBLFQ) [1], for simulating the evolution of a quark jet inside a classical color background field, first for a $|q\rangle$ state [2], then also including $|qg\rangle$ components [3, 4]. Unlike pQCD-based approaches, tBLFQ calculates the evolution process at the amplitude level, and enables relaxation of the eikonal and collinear radiation approximations.

2. Methodology

We consider a high-energy quark jet moving in the positive z direction, traversing a medium moving in the negative z direction. We treat the quark as a quantum state and the medium as an external background field, with the interaction occurring over a finite distance $0 \leq x^+ \leq L_\eta$.

2.1 The light-front Hamiltonian in the $|q\rangle + |qg\rangle$ space

The Lagrangian for the process being considered is the QCD Lagrangian with an external field,

$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}{}_a F_{\mu\nu}{}^a + \bar{\Psi}(i\gamma^\mu D_\mu - m_q)\Psi, \quad (1)$$

where $F_a^{\mu\nu} \equiv \partial^\mu C_a^\nu - \partial^\nu C_a^\mu - g f^{abc} C_b^\mu C_c^\nu$, $D^\mu \equiv \partial^\mu + igC^\mu$, and $C^\mu = A^\mu + \mathcal{A}^\mu$ is the sum of the quantum gauge field A^μ and the background gluon field \mathcal{A}^μ . The light-front Hamiltonian is obtained through Legendre transformation in the light-cone gauge $A^+ = \mathcal{A}^+ = 0$. In the truncated Fock space $|q\rangle + |qg\rangle$, it consists of three parts, $P^-(x^+) = P_{KE}^- + V_{qg} + V_{\mathcal{A}}(x^+)$, the kinetic energy, the interaction between the quark and the dynamical gluon, and the medium interaction,

$$P_{KE}^- = \int dx^- d^2x_\perp \left\{ -\frac{1}{2}A_a^j (i\nabla_\perp)^2 A_j^a + \frac{1}{2}\bar{\Psi}\gamma^+ \frac{m_q^2 - \nabla_\perp^2}{2i\partial_-} \Psi \right\}, \quad (2a)$$

$$V_{qg} = \int dx^- d^2x_\perp g \bar{\Psi}\gamma^\mu T^a \Psi A_\mu^a, \quad (2b)$$

$$V_{\mathcal{A}}(x^+) = \int dx^- d^2x_\perp g \bar{\Psi}\gamma^+ T^a \Psi \mathcal{A}_+^a(x^+) + g f^{abc} \partial^+ A_b^i A_i^c \mathcal{A}_+^a(x^+). \quad (2c)$$

The background field \mathcal{A}^μ accounts for the target, and we describe it using the McLerran-Venugopalan (MV) model [5]. It is a classical field satisfying the reduced Yang-Mills equation,

$$(m_g^2 - \nabla_\perp^2)\mathcal{A}_a^-(\vec{x}_\perp, x^+) = \rho_a(\vec{x}_\perp, x^+), \quad \langle \rho_a(x)\rho_b(y) \rangle = g^2 \tilde{\mu}^2 \delta_{ab} \delta^2(\vec{x}_\perp - \vec{y}_\perp) \delta(x^+ - y^+). \quad (3)$$

The saturation scale is related to $\tilde{\mu}$ as $Q_s^2 = C_F (g^2 \tilde{\mu})^2 L_\eta / (2\pi)$, with $C_F = (N_c^2 - 1)/(2N_c) = 4/3$.

2.2 Evolution of the state in a basis representation

The evolution of quantum states is governed by the time-evolution equation on the light front. In the interaction picture (denoted by the subscript I), the equation reads

$$i \frac{\partial}{\partial x^+} |\psi; x^+\rangle_I = \frac{1}{2} V_I(x^+) |\psi; x^+\rangle_I, \quad (4)$$

with the interaction Hamiltonian $V_I(x^+) = e^{i\frac{1}{2}P_{KE}^- x^+} V(x^+) e^{-i\frac{1}{2}P_{KE}^- x^+}$. The interaction picture state is related to the Schrödinger picture state by $|\psi; x^+\rangle_I = e^{i\frac{1}{2}P_{KE}^- x^+} |\psi; x^+\rangle$.

We implement a non-perturbative treatment by decomposing the time-evolution operator into many small steps of the light-front time x^+ , then solving each timestep in the sequence numerically,

$$|\psi; x^+\rangle_I = \mathcal{T}_+ e^{-\frac{i}{2} \int_0^{x^+} dz^+ V_I(z^+)} |\psi; 0\rangle_I = \lim_{n \rightarrow \infty} \prod_{k=1}^n \mathcal{T}_+ e^{-\frac{i}{2} \int_{x_{k-1}^+}^{x_k^+} dz^+ V_I(z^+)} |\psi; 0\rangle_I, \quad (5)$$

with \mathcal{T}_+ denoting light-front time ordering. The step size is $\delta x^+ \equiv x^+/n$, and the intermediate time is $x_k^+ = k\delta x^+ (k = 0, 1, 2, \dots, n)$ with $x_0^+ = 0$ and $x_n^+ = x^+$.

We use a lattice with periodic boundary conditions in the transverse dimensions \vec{x}_\perp , ranging in $[-L_\perp, L_\perp]$ with $2N_\perp$ sites such that $a_\perp = L_\perp/N_\perp$ is the lattice spacing, and a loop with (anti-)periodic boundary condition in the x^- direction, of length $2L$, for the gluon(quark). The lattice introduces infrared (IR) and ultraviolet (UV) cutoffs in the transverse momentum space \vec{p}_\perp , $\lambda_{IR} = d_p = \pi/L_\perp$ and $\lambda_{UV} = \pi/a_\perp$. The longitudinal momentum p^+ is quantized in units of $2\pi/L$, and the gluon(quark) is allowed to take a positive (half-)integer number in this unit. The total momentum is denoted as $p^+ = K2\pi/L$ with K a half-integer. Then the longitudinal momentum fraction of the gluon, $z \equiv p_g^+/p^+$, has a resolution of $1/K$.

In the chosen discrete basis representation, the state is a column vector of basis coefficients, and the Hamiltonian is in the matrix form. The numerical method for this specific problem is optimized in Ref. [3]. In short, within each small time step δx^+ , we treat P_{KE}^- and $V_{\mathcal{A}}$ as time-constant and carry out matrix exponentiation in the momentum and coordinate space, respectively; the operation with V_{qg} uses the fourth-order Runge-Kutta method in the momentum space.

3. Result

In studying the phenomenon of jet momentum broadening inside a medium, we examine the expectation value of the transverse momentum square $\langle p_\perp^2(x^+) \rangle$, and the related quenching parameter defined as $\hat{q} = \Delta \langle p_\perp^2(x^+) \rangle / \Delta x^+$.

3.1 Eikonal analytical result

In the eikonal limit of $p^+ = \infty$, only the $V_{\mathcal{A}}$ term survives in the Hamiltonian, and the evolution operator reduces to the Wilson line, then $\langle p_\perp^2(x^+) \rangle$ and \hat{q} can be derived analytically using the Wilson line correlators. Here, we present the derivation result for the single quark/gluon state, and a novel derivation for the quark-gluon state following Ref. [4].

3.1.1 Single-particle state

The Wilson line of a quark is $U_F(0, x^+; \vec{x}_\perp) \equiv \mathcal{T}_+ \exp\left(-ig \int_0^{x^+} dz^+ \mathcal{A}_a^-(\vec{x}_\perp, z^+) T^a\right)$, in which T^a is the SU(3) generator in the fundamental representation. Replacing T^a by the generators in the adjoint representation, t^a , one gets the adjoint Wilson line for the gluon, $U_A(0, x^+; \vec{x}_\perp)$. The momentum transfer can be evaluated from the Wilson line correlator

$$S_F(0, x^+; r) = \frac{1}{N_c} \text{Tr} \langle U_F^\dagger(0, x^+; \vec{x}_\perp) U_F(0, x^+; \vec{y}_\perp) \rangle_m = e^{-C_F g^4 \hat{\mu}^2 x^+ [L(0) - L(r)]}, \quad (6)$$

with $r = |\vec{x}_\perp - \vec{y}_\perp|$ and $L(r) = \int_{\mathbf{p}} e^{-i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{y}_\perp)} / (m_g^2 + \vec{p}_\perp^2)^2 = m_g r K_1(m_g r) / (4\pi m_g^2)$,¹ as

$$\langle p_\perp^2(x^+) \rangle_{Eik} = \langle p_\perp^2(0) \rangle - \nabla_r^2 S_F(0, x^+; r)|_{r=0}. \quad (7)$$

The quenching parameter \hat{q} follows as,

$$\hat{q}_{Eik} = -C_F g^4 \tilde{\mu}^2 \nabla_r^2 L(r)|_{r=0} = \frac{C_F g^4 \tilde{\mu}^2}{4\pi} \left\{ \log \left[1 + \frac{1}{(m_g a_\perp / \pi)^2} \right] - \frac{1}{1 + (m_g a_\perp / \pi)^2} \right\}. \quad (8)$$

In analogy, one gets the gluon \hat{q} replacing C_F by $C_A = N_c$ in Eq. (8).

3.1.2 Quark-gluon state

The quark-gluon Wilson line is built as the tensor product of a quark and a gluon Wilson line, $U_{qg}(0, x^+; \vec{x}_\perp, \vec{y}_\perp) \equiv U_F(0, x^+; \vec{x}_\perp) \otimes U_A(0, x^+; \vec{y}_\perp)$. The probability distribution of the quark-gluon state is then given by Wilson line correlator $U_{qg}^\dagger U_{qg}$, and the state's transverse momentum square can be calculated accordingly. It contains three terms,

$$\langle p_\perp^2(x^+) \rangle_{qg,c;Eik} = \langle \vec{p}_{q,\perp}^2(x^+) \rangle_{Eik} + \langle \vec{p}_{g,\perp}^2(x^+) \rangle_{Eik} + 2 \langle \vec{p}_{q,\perp}(x^+) \cdot \vec{p}_{g,\perp}(x^+) \rangle_{c;Eik}. \quad (9)$$

The first two terms are the same as Eq. (7) with the corresponding Casimir, and the third term depends on the initial color configuration and the quark-gluon separation [4],

$$\langle \vec{p}_{q,\perp}(x^+) \cdot \vec{p}_{g,\perp}(x^+) \rangle_{c;Eik} = \begin{cases} 0, & c = 3 \otimes 8 \\ -\frac{N_c \sqrt{2}}{2} f_{12}, & c = 3 \\ -\frac{\sqrt{2}}{2} f_{12}, & c = \bar{6} \\ \frac{\sqrt{2}}{2} f_{12}, & c = 15 \end{cases}, \quad (10)$$

in which c represents the initial color configuration of the state, and

$$f_{12} = \int_{\mathbf{v}} f_{Rel}(\vec{v}_\perp) \nabla_{\mathbf{v}}^2 L(\mathbf{v}) \frac{2\sqrt{2}}{4N_c [L(0) - L(\mathbf{v})]} \left\{ e^{-g^4 \tilde{\mu}^2 N_c [L(0) - L(\mathbf{v})] x^+} - 1 \right\}. \quad (11)$$

The quantity $f_{Rel}(\vec{v}_\perp)$ is the distribution function of the quark-gluon relative coordinate $\vec{v}_\perp = \vec{x}_{q,\perp} - \vec{x}_{g,\perp}$, and it can be obtained by integrating the wavefunction square over the center-of-mass coordinate $\vec{R}_\perp = z\vec{x}_{q,\perp} + (1-z)\vec{y}_{g,\perp}$, $f_{Rel}(\vec{v}_\perp) \equiv \int_{\mathbf{R}} |\tilde{\phi}(\vec{x}_{q,\perp}, \vec{x}_{g,\perp})|^2$.

3.2 Non-eikonal numerical result

We perform the simulations in the $|q\rangle + |qg\rangle$ space, with the initial state as a single quark of $\vec{p}_\perp = \vec{0}_\perp$ at a finite p^+ .

To quantify the medium-induced gluon emission, we define $\delta P_{|qg\rangle}$ as the difference of the probability of the quark jet in the $|qg\rangle$ sector in the medium and that in the vacuum,

$$\delta P_{|qg\rangle}(Q_s, x^+) \equiv P_{|qg\rangle}(Q_s, x^+) - P_{|qg\rangle}(Q_s = 0, x^+). \quad (12)$$

The result is shown in Fig. 1. In the left panel, we see that the $\delta P_{|qg\rangle}$ curve forms a slight dip at an early time, then after around the point $x^+ = 12 \text{ GeV}^{-1}$, grows linearly in time. Additionally, a




Figure 1: The medium-induced gluon emission. Simulation parameters: $p^+ = 17$ GeV, $N_\perp = 16$, $L_\perp = 50$ GeV $^{-1}$, $L_\eta = 50$ GeV $^{-1}$ and $K = 8.5$.




Figure 2: The non-eikonal correction to $\langle p_\perp^2 \rangle$ and \hat{q} . Simulation parameters are the same as in Fig. 1.

larger $|qg\rangle$ component develops as Q_s increases. The right panel shows that the $\delta P_{|qg\rangle}$ of the final state ($x^+ = L_\eta$) is approximately proportional to Q_s^2 (c.f. [6] has a similar observation).

We then analyze the non-eikonal and radiative correction to the momentum broadening. We define $\delta \langle p_\perp^2 \rangle$ and $\delta \hat{q}$ as the difference of the quantity that is calculated from the total momentum of the quark jet in the $|q\rangle + |qg\rangle$ space, and the eikonal result of a bare quark [as in Eq. (8)],

$$\delta \langle p_\perp^2 \rangle \equiv \langle p_\perp^2 \rangle - \langle p_\perp^2 \rangle_{Eik}, \quad \delta \hat{q} \equiv \hat{q} - \hat{q}_{Eik}. \quad (13)$$

The results are shown in Fig. 2: $\delta \langle p_\perp^2 \rangle$ increases over the evolution time at various Q_s , and $\delta \hat{q}$ extracted from the final state $\delta \langle p_\perp^2 \rangle$ increases non-trivially when Q_s increases.

4. Summary

We present a study on the momentum broadening of in-medium jet evolution using the tBLFQ approach [3, 4], a non-perturbative light-front Hamiltonian formalism. We first provide a novel analytical derivation of the eikonal expectation value of the quark-gluon state's transverse momentum for any color and spatial distribution. We then perform the numerical simulation of the real-time jet evolution in the Fock space of $|q\rangle + |qg\rangle$ at various medium densities. With the obtained jet light-front wavefunction, we extract the gluon emission rate and the quenching parameter. We find their non-eikonal contributions sizable, time-dependent, and associated with saturation scale.

¹Here and throughout the paper we use the shorthand notation $\int_{\mathbf{p}} \equiv \int d^2 p_\perp / (2\pi)^2$ and $\int_r \equiv \int d^2 r_\perp$.

Acknowledgments

We are very grateful to Guillaume Beuf, Fabio Dominguez, Miguel A. Escobedo, Xabier Feal, Sigtryggur Hauksson, Cyrille Marquet, Wenyang Qian, Andrecia Ramnath, Andrey Sadofyev, Konrad Tywoniuk, James P. Vary, Xin-Nian Wang, and Bin Wu for helpful and valuable discussions.

XB is supported by new faculty startup funding by the Institute of Modern Physics, Chinese Academy of Sciences, by Key Research Program of Frontier Sciences, Chinese Academy of Sciences, Grant No. ZDBS-LY-7020, by the Natural Science Foundation of Gansu Province, China, Grant No. 20JR10RA067, by the Foundation for Key Talents of Gansu Province, by the Central Funds Guiding the Local Science and Technology Development of Gansu Province, Grant No. 22ZY1QA006 and by the Strategic Priority Research Program of the Chinese Academy of Sciences, Grant No. XDB34000000. ML and CS are supported by Xunta de Galicia (Centro singular de Investigación de Galicia accreditation 2019-2022), European Union ERDF, the “Maria de Maeztu” Units of Excellence program under project CEX2020-001035-M, the Spanish Research State Agency under project PID2020-119632GB-I00, and European Research Council under project ERC-2018-ADG-835105 YoctoLHC. TL is supported by the Academy of Finland, the Centre of Excellence in Quark Matter (project 346324) and project 321840. This work was also supported under the European Union’s Horizon 2020 research and innovation by the STRONG-2020 project (grant agreement No. 824093). The content of this article does not reflect the official opinion of the European Union and responsibility for the information and views expressed therein lies entirely with the authors.

References

- [1] J.P. Vary, H. Honkanen, J. Li, P. Maris, S.J. Brodsky, A. Harindranath et al., *Hamiltonian light-front field theory in a basis function approach*, *Phys. Rev. C* **81** (2010) 035205 [0905.1411].
- [2] M. Li, X. Zhao, P. Maris, G. Chen, Y. Li, K. Tuchin et al., *Ultrarelativistic quark-nucleus scattering in a light-front Hamiltonian approach*, *Phys. Rev. D* **101** (2020) 076016 [2002.09757].
- [3] M. Li, T. Lappi and X. Zhao, *Scattering and gluon emission in a color field: A light-front Hamiltonian approach*, *Phys. Rev. D* **104** (2021) 056014 [2107.02225].
- [4] M. Li, T. Lappi, X. Zhao and C.A. Salgado, *On the momentum broadening of in-medium jet evolution using a light-front Hamiltonian approach*, 2305.12490.
- [5] L.D. McLerran and R. Venugopalan, *Computing quark and gluon distribution functions for very large nuclei*, *Phys. Rev. D* **49** (1994) 2233 [hep-ph/9309289]; L.D. McLerran and R. Venugopalan, *Gluon distribution functions for very large nuclei at small transverse momentum*, *Phys. Rev. D* **49** (1994) 3352 [hep-ph/9311205]; L.D. McLerran and R. Venugopalan, *Fock space distributions, structure functions, higher twists and small x*, *Phys. Rev. D* **59** (1999) 094002 [hep-ph/9809427].
- [6] Y.-Y. Zhang and X.-N. Wang, *Parton rescattering and gluon saturation in dijet production at EIC*, *Phys. Rev. D* **105** (2022) 034015 [2104.04520].