

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Vainio, Veera; El Geneidy, Sami; Halme, Panu; Peura, Maiju; Kotiaho, Janne S.

Title: Biodiversity impact of the consumption of peat and wood-fired district heating

Year: 2024

Version: Published version


Copyright: © 2024 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Vainio, V., El Geneidy, S., Halme, P., Peura, M., & Kotiaho, J. S. (2024). Biodiversity impact of the consumption of peat and wood-fired district heating. *Environmental Impact Assessment Review*, 108, Article 107587. <https://doi.org/10.1016/j.eiar.2024.107587>



Biodiversity impact of the consumption of peat and wood-fired district heating

Veera Vainio^{a,b,*}, Sami El Geneidy^{b,c}, Panu Halme^{a,b}, Maiju Peura^{a,b}, Janne S. Kotiaho^{a,b}

^a Department of Biological and Environmental Science, University of Jyväskylä, P.O. Box 35, FIN-40014, Finland

^b School of Resource Wisdom, University of Jyväskylä, P.O. Box 35, FIN-40014, Finland

^c School of Business and Economics, University of Jyväskylä, P.O. Box 35, FIN-40014, Finland

ARTICLE INFO

Keywords:

Biodiversity impact
District heating
Biofuel
Energy-wood
Peat
Habitat hectare

ABSTRACT

A shift from fossil fuels to biofuels is becoming an increasingly important part of national and corporate climate strategies, but the biodiversity impacts of biofuels are often ignored and remain poorly understood. Here we developed a novel eight-step general framework for the assessment of consumption-based biodiversity impacts of biofuels and applied the framework to assess the biodiversity impacts of peat and wood-fired district heating in Finland. We utilize the habitat hectare approach for which information about the ecosystem condition before and after the impact is needed. The habitat hectare approach has not yet been used on assessments of consumption-based biodiversity impacts. Of the different fuel types used in peat and wood-fired district heating in Finland, peat had the highest biodiversity impact per unit area, followed by chips from roundwood and logging residue chips. By-products from sawmill industry are argued to have no additional biodiversity impacts. Interestingly, however, when we consider the biodiversity impacts per unit energy, chips from roundwood had the highest and peat the lowest biodiversity impact. Our results suggest there is a truly substantial trade-off between the climate and biodiversity impacts of wood and peat firing. We conclude that the habitat hectare approach combined with the presented general assessment framework is a promising approach to be used in the assessment of consumption-based biodiversity impacts in different land use contexts, such as biodiversity offsetting, around the globe. This protocol should be further developed and refined in different systems and with different bio-based materials and land use contexts.

1. Introduction

Energy used in buildings is close to 40% of the global energy consumption, and heating consumes roughly 60% of this (Pérez-Lombard et al., 2008). District heating is commonly used in Northern and Eastern Europe as well as in Russia and China (Werner, 2017; Paiho and Saastamoinen, 2018; Mazhar et al., 2018). For example, about half of the Finnish as well as Swedish population and buildings are dependent on district heating (Eriksson et al., 2007; Paiho and Saastamoinen, 2018). Moving away from fossil fuels is likely to increase the demand for wood and other renewable biomass-based district heating (Persson and Werner, 2011; Mazhar et al., 2018), and there indeed is some potential in bioenergy (Creutzig et al., 2015; Börjesson et al., 2017). However, expanding the use of bioenergy is already constrained for example by its availability (Szarka et al., 2017; Anttila et al., 2018; Verkerk et al., 2019) and competition of resources between energy and non-energy uses

(Mandley et al., 2020). Also, the pressure to increase forest carbon sinks is preventing significant increases in forest biomass harvests (Masson-Delmotte et al., 2018). Biodiversity impacts of the extraction of the biomass is likely to cause yet another constraint (Ranius et al., 2018; Weldu and Assefa, 2016), and while the extraction of forest biomass is already known to cause adverse biodiversity impacts (e.g., Toivanen et al., 2012; Ranius et al., 2018), these impacts are still to be quantified in relation to district heating.

Human activity has generated significant changes in the biosphere over the past several decades (Foley et al., 2005; Barnosky et al., 2012; Ellis et al., 2013). The abundance and diversity of nature are declining, species extinction rates are accelerating, and ecosystems are being degraded all over the world (Barnosky et al., 2012; Newbold et al., 2015; IPBES, 2018). The most important direct drivers of biodiversity loss and ecosystem degradation are extensive land-use and resource use including direct exploitation of species, but also climate change,

* Corresponding author at: Department of Biological and Environmental Science, University of Jyväskylä, P.O. Box 35, FIN-40014, Finland.
E-mail address: veera.s.vainio@jyu.fi (V. Vainio).

spreading of harmful alien species, and pollution of habitats (IPBES, 2019; Jaureguiberry et al., 2022). It has been estimated that the negative impacts of human land-use already affect up to 75% of the earth's surface (IPBES, 2018). Extracting biomass for district heating can be considered to fall under the category of direct exploitation and it clearly has potential to contribute to biodiversity loss.

Organizations, for example private businesses and public services such as district heating plants, hospitals, and education institutions, are the way we have organized the functions in our societies. To understand the role of organizations in enhancing or reducing planetary well-being (Kortetmäki et al., 2021), we need to be able to identify and quantify the environmental impacts, such as carbon emissions and biodiversity loss their operations are causing. Carbon footprint is a commonly used tool for assessing consumption-based climate impacts of human activities (Wiedmann and Minx, 2008), but tools and methodologies to assess consumption-based biodiversity impacts are still less developed but emerging (Lenzen, 2014; Marques et al., 2017; Winter et al., 2017; Crenna et al., 2020; Lammerant et al., 2021; El Geneidy et al., 2023). So far assessments of biodiversity impacts of any consumption at the level of an organization are rare (El Geneidy et al., 2021; Bull et al., 2022; Taylor et al., 2023).

In this paper we develop a framework for assessing the negative biodiversity impacts of the consumption of peat and wood-fired district heating and apply it to assess the biodiversity impacts of one district heating dependent organization, the University of Jyväskylä.

2. Assessing biodiversity impacts

2.1. Habitat hectares as a common currency for different biodiversity impacts

In general, human activities can have negative or positive impacts on species and ecosystems. Ecosystem degradation refers specifically to negative or harmful impacts (IPBES, 2018). For example, the impact that a construction project causes to any ecosystem, or the harm that logging causes to forest ecosystems can be considered as ecosystem degradation.

Detailed measurement of biodiversity is difficult since even a small area almost anywhere may contain hundreds or thousands of species and even more individuals. In practice it is simply impossible to count them all, and thus biodiversity is not readily amenable to very detailed measurement. In general, biotopes, ecosystems, habitat types or ecological features important for large numbers of species can be categorized and their condition mapped much more readily than occurrences of individual species or their population sizes. Thus, we approach the problem by looking at the condition of the ecosystems rather than species or their population. This approach has also been called the habitat hectares approach (Parkes et al., 2003).

The condition of any ecosystem can be considered to vary from 0 to 1. An area that has been completely degraded obtains value 0, and an area in a completely natural, undisturbed condition obtains value 1. In other words, habitat hectare is a condition weighted area where one habitat hectare equals one hectare of undisturbed ecosystem. For example, a value 0.4 is assigned to an area that has been degraded so that it has lost 60% of its pristine condition (Parkes et al., 2003; Kangas et al., 2023). Habitat hectare is a common currency that can be used when e.g., losses and gains are being balanced in biodiversity offsetting projects (see e.g., Moilanen and Kotiaho, 2018, 2020, 2021).

Life cycle assessment (LCA) and environmentally extended input-output (EEIO) analysis are commonly used to assess land use and associated biodiversity impacts based on consumption (Marques et al., 2017; Crenna et al., 2020; Damiani et al., 2023). While both LCA and EEIO can be used to assess the biodiversity impacts of consumption across the value chain, EEIO analysis is more appropriate for analyzing sectoral impacts, while LCA is more appropriate for analyzing product or process specific impacts (Marques et al., 2017). Here we utilize the habitat hectare approach, which is closer to LCA than EEIO, because we are

analyzing the impact of wood and peat for the district heating consumption with process-based impact factors.

Other metrics for assessing biodiversity impacts include e.g. mean species abundance (MSA, Alkemade et al., 2009) and potentially disappeared fraction of species (PDF, Verones et al., 2020) (for a review of different approaches, see e.g. Crenna et al., 2020; Damiani et al., 2023). Both of these are species-based metrics, and thus not directly amenable for the purpose of the current study. It has been argued that in the past there was a focus on indicators focused on species richness, whereas less attention was given to functional and structural attributes of biodiversity (Curran et al., 2011), which seems to be prevalent today as well (Crenna et al., 2020; Damiani et al., 2023). Lindqvist et al. (2016) explored the use of expert judgement and ecosystem condition valuation in assessing biodiversity impacts of forest management. Similarly, the habitat hectare approach presented in this paper supports the development of biodiversity impact indicators based on functional and structural attributes.

2.2. Biodiversity impacts of extracting wood for energy

Forest-based bioenergy can be sourced by extracting additional biomass from production forests or by using industrial by-product residues such as bark and sawdust. Extraction of logging residues results in biomass and thus nutrient removal, as well as disturbance of soils and several other environmental features important for biodiversity (Ranius et al., 2018). How much biomass is extracted at site-level varies among countries, with higher values reported from Sweden and Finland than for example from France and North America (Thiffault et al., 2016). Nevertheless, the evidence is clear that logging residue extraction can have significant negative impacts on biodiversity (Ranius et al., 2018).

Dead wood is an important resource for forest biodiversity. Woody biomass removal reduces the food and habitat available for dead wood dependent organisms (Bouget et al., 2012). Energy-wood extraction decreases the number of wood-inhabiting fungi due to decreased number and volume of habitat patches (Toivanen et al., 2012) and affects many other species groups by reducing the availability of shelter and nesting sites (Ecke et al., 2002). For insects the energy-wood extraction means actual removal of a large number of individuals from forest sites, because insects colonize the dead wood piles before the wood is transported to power plants (Hedin et al., 2008). Physical and geochemical changes due to logging residue extraction also affect field layer vegetation. Overall levels of soil organic matter and soil nutrients are reduced, with a potential loss of soil fertility and site productivity. Changes in soil physical properties due to increased heavy machinery traffic in the forest, thinner protective mats of logging residue, and changes in soil moisture and organic matter are examples of the changes caused by the extraction (Verkerk et al., 2019).

2.3. Biodiversity impacts of extracting peat for energy

In Finland many peatlands are threatened ecosystems and host a high number of threatened species. Currently the main threat to peatlands is habitat destruction due to ditching, peat mining, forestry, and agriculture (Alanen and Aapala, 2015; Hyvärinen et al., 2019; Kontula and Raunio, 2019). Peat mining has a detrimental effect on peatlands. Whole ecosystems are destroyed when peatlands are drained, and all the surface vegetation is removed (Finnish Ministry of the Environment, 2015). The direct impact of removing the surface layer including all the flora and fauna is evident, but indirect biodiversity impacts outside the direct extraction area via changes in hydrology can also occur. Hydrology is one of the most important features of peatland ecosystems and it largely determines peatland ecosystem functions and species diversity (Konar et al., 2013; Haapalehto et al., 2014; Kareksela et al., 2015).

Drainage of the extraction site unavoidably also affects the hydrology of the surrounding areas by lowering their groundwater levels (Tahvanainen, 2011; Alanen and Aapala, 2015; Paal et al., 2016).

Drainage can also increase nutrient leaching and the flow of suspended solids to downstream water systems (Finnish Ministry of the Environment, 2015). These changes degrade the peatland ecosystems (e.g., Mazerolle, 2003).

3. Framework for assessing the biodiversity impact of district heating consumption of an organization

There are several steps that need to be taken to assess an organization's biodiversity impact. In Fig. 1 we depict a framework for the assessment with eight steps. In this section, we describe each of the steps in detail and apply the framework to assess the biodiversity impact of the district heating consumed by the University of Jyväskylä.

3.1. Step 1. Set the limits of the assessment

To start the assessment, one needs to delimit the system in focus. In other words, one needs to determine which parts and operations of the organization will be included in the assessment of the biodiversity impacts. Once the limit has been decided, the total heat consumption of the delimited part needs to be calculated.

In the current study we focused on the consumption of district heating of the University of Jyväskylä and thus decided to include buildings used by the University that are located within the City of Jyväskylä district heating network. The areas of the buildings rented out for other organizations were deduced from the total consumption of each of the buildings. Data was available for the years 2019–2021, and 2021 is used as an example in the text. The other two years are included in Table 1.

Note that this study focuses on the direct ecological impacts of extracting fuels for district heat production. Therefore, indirect impacts from e.g., the machinery needed for extraction, or the infrastructure needed for heat production are excluded.

The total consumption of district heating within the scope mentioned above at the University of Jyväskylä was 26,321 MWh in 2021.

3.2. Step 2. Identify the fuel types used

The fuel types and the share of each with which the heat is produced must be identified, and an understanding about how and from where the fuels are being extracted needs to be developed.

The University of Jyväskylä purchases the district heating from the local energy company Alva that is owned by the City of Jyväskylä. Most of the power plants of Alva are combined heat and power plants (CHP), so they produce heat and electricity simultaneously. This means that the amount of fuel used for electricity production simultaneously produces heat on the side or vice versa. Since this study focuses only on heating, we need to determine what share of the biodiversity impacts is to be allocated to heat and to electricity respectively. For this we used the energy-based method that is used for emission allocation for CHP (Rosen, 2008) and statistics of Alva's annual energy production (Oy, 2022). The share of thermal energy or produced heat is calculated as follows:

$$f_Q = Q/(E + Q) \quad (1)$$

where f_Q is the fraction of impacts to be allocated to heating and E and Q are the net outputs of electricity and heat from the cogeneration system respectively. In 2021 Alva produced 1.3 TWh of heat and 0.7 TWh of electricity (Oy, 2022). 55,000 MWh of the heat was produced in a heat-only plant situated around 100 km out of Jyväskylä and thus was assumed to be excluded from the heat production considered in this study. Therefore, the share of impacts allocated for heating in this case equals $(1.3 - 0.055) \text{ TWh} / [0.7 \text{ TWh} + (1.3 - 0.055) \text{ TWh}] = 0.640$ or 64%. The allocation will be taken into account in Step 6 of the impact calculations.

The energy is produced by burning wood fuels (66.5%) and peat (32.5%). Coal, oil, and biogas are burnt in small quantities (1.0%) as backup fuels (Oy, 2023, Fig. 2). We decided to assess the biodiversity impacts of consuming district heat produced using the main fuels, wood and peat, and excluded the imported fossil fuels from the analysis. Wood fuels were comprised of forest fuels and timber industry residues. Forest fuels are further divided to logging residue chips and to chips from roundwood. Logging residue is mostly extracted from clear-cut sites, whereas roundwood is mostly extracted from thinnings of young forest stands.

Residues from timber industry include bark chips, wood residue chips and sawdust. Logging is necessary for obtaining timber industry residues, but the residues are not the primary reason for the logging. It is worth noting that if an increase in the demand for timber industry residues would encourage more logging, then the use of residues in district heating would cause an additional biodiversity impact. However,




Fig. 1. Framework and steps for assessing the biodiversity impact of the consumption of peat and wood-fired district heating of an organization. Blue color indicates common steps for all fuel types and orange color indicates the calculation steps that need to be completed separately for each fuel type. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. The fuel types and the share (percentage of the total energy produced) of each that are used for district heat production in the City of Jyväskylä (Alva, 2020, Official Statistics of Finland (OSF), 2021). The backup fuels indicated with gray color are not included in the biodiversity impact assessment.

currently the demand and price of timber industry residues is low compared to those of timber wood (Metsälehti, 2023). Therefore, we considered the timber industry residues to have zero biodiversity impact in district heating use. The biodiversity impact of logging for timber should be addressed to the primary timber industry such as sawmills.

Peat was assumed to be milled peat, which is the main form of peat used in district heating systems in Finland (Alakangas et al., 2016).

Of the total heat consumption of the University of Jyväskylä in 2021, 11,272 MWh was produced from timber industry residues, 2030 MWh from logging residue chips, 4183 MWh from roundwood chips, and 8554 MWh from peat (Table 1).

3.3. Step 3. Estimate the energy yield per volumetric unit of each fuel type

Steps 3–6 need to be conducted separately for each of the fuel types.

The energy yield per volumetric unit of each fuel type (MWh/m³) can be derived from literature. We utilized national statistics from the Technical Research Centre of Finland (Alakangas and Impola, 2014; Alakangas et al., 2016). The energy yields per volumetric unit for the different fuel types included in the assessment are listed in Table 1.

3.4. Step 4. Estimate the volumetric yield of each fuel type per unit of extraction area

The volumetric yield per unit of extraction area can be estimated based on literature. Detailed information on the different extraction practices of the fuel types is also necessary. This process is next described separately for each fuel type. For timber industry residues this step is not necessary, because, as was explained above, we assumed that the negative biodiversity impact due to logging for timber should be addressed for the timber sawmill industry and not for the district heating industry using their residues. It should be noted that the volumetric yield varies by region and country and each assessment should be based on the local context.

3.4.1. Logging residue chips

Logging residues contain branches and treetops from commercial logging, and they are customarily collected only after final felling and not from thinnings (Äijälä et al., 2014; Koistinen et al., 2019). Therefore, we assumed that all logging residues originated from final felling sites. The main method for final felling in Central Finland is clear-cutting (Official Statistics of Finland (OSF), 2018). The extraction of logging residues after clear-cutting causes an additional biodiversity impact because the amount of decaying wood material is significantly reduced (Toivanen et al., 2012; Mikkonen et al., 2023).

The average yield of logging residue is 50–60 m³/ha in Finland (Viitasaari, 2013).

3.4.2. Chips from roundwood

Roundwood from first thinning of young stands, where the average chest-height diameter is 8–16 cm, is used for energy production (Äijälä et al., 2014). Thus, the assessed biodiversity impact is the thinning of young stands.

The yield of timber from the first thinning is commonly 30–60 m³/ha (Asikainen et al., 2012; Lauhanen et al., 2014; Karttunen et al., 2016). We used the median value of 45 m³ as the volume of roundwood that is extracted from an average hectare of thinning of young stands.

3.4.3. Peat

Energy yields per hectare were readily available for peat in the literature. To follow the same steps as with other fuel types, the volumetric yield per hectare was calculated by dividing the energy yield per hectare by the energy yield per volumetric unit of peat (0.91 MWh/m³, see Table 1).

The average annual energy yields of peat in all of Finland vary from 400 to 500 MWh/ha (Väyrynen et al., 2008). Detailed information on the origin of the peat utilized by the local energy company is not available. Therefore, although it is known that some proportion of the peat is produced outside Central Finland (Flyktman, 2012), in the calculations we used the average annual energy yield of peat in Central Finland, which is 400 MWh/ha (Flyktman, 2012). Thus, the volumetric yield of peat is 439.6 m³/ha (400 MWh/ha divided by 0.91 MWh/m³).

3.5. Step 5. Quantify the biodiversity impact per unit area for each fuel type

To assess the biodiversity impact of the extraction of each of the fuel types in habitat hectares, information on two aspects of each of the ecosystems where extraction is performed is needed: i) the ecosystem condition of the extraction area prior to the extraction (habitat hectares per hectare (hha/ha)), and ii) the ecosystem condition of the extraction area after the extraction (hha/ha). The biodiversity impact is then calculated as the difference in per hectare condition before and after the extraction. Sometimes several calculations per fuel type may be needed if the ecosystem condition before extraction varies from site to site.

3.5.1. Logging residue chips

3.5.1.1. Determine the condition of the focal ecosystem before the extraction per unit area. Logging residue chips are obtained from biomass that is left behind from forest clear-cutting for other than district heating purposes. In other words, the clear-cutting was not conducted to obtain logging residues for district heating. Thus, in this case the impact of clear-cutting can be excluded from the biodiversity impact calculation and the focus must be on what is the additional biodiversity impact caused by the logging residue extraction.

The impact of forest management actions on ecosystem quality in Finland has been assessed by Mikkonen et al. (2023), Moilanen and Kotiaho (2020), and Jalkanen et al. (2023). Based on these studies, we assumed the ecological condition of a clear-cut forest site to be 0.1 habitat hectares per hectare before the extraction of the logging residue.

3.5.1.2. Determine the condition of the focal ecosystem after the extraction per unit area. When logging residue is removed from the final felling site, the amount of decaying wood is reduced by 39% (Eräjää et al., 2010). Elsewhere it has been estimated that 60% of the area's ecological value is determined by dead wood (Kotiaho et al., 2015; Kotiaho et al., 2016). Thus, by assuming that dead wood comprises 60% of the 0.1 habitat hectares per hectare of the clear-cut site, the condition of the clear-cut site after the logging residue extraction can be calculated as 0.10 hha/ha - (0.10 hha/ha × 0.60 × 0.39) = 0.077 hha/ha.




Fig. 3. Quantifying the biodiversity impact per unit area for logging residue chips. The assessed impact is logging residue removal from clear-cuts. Before the extraction the ecological condition of focal ecosystem, i.e., clear-cut area, is 0.1 habitat hectares and dead wood determines 60% of the condition (Step 5a). Logging residue removal decreases the amount of dead wood in clear-cut area by 39%, and thus the ecological condition of focal ecosystem after impact is 0.077 habitat hectares (Step 5b). Biodiversity impact per unit area is the difference in condition before and after the impact which is 0.023 hha/ha (Step 5c).

3.5.1.3. Calculate the biodiversity impact per unit area by taking the difference between the condition before and after the extraction. The negative biodiversity impact of logging residue extraction per hectare is $0.1 \text{ hha/ha} - 0.077 \text{ hha/ha} = 0.023 \text{ hha/ha}$. The steps 5a to 5c are illustrated in Fig. 3.

3.5.2. Chips from roundwood

3.5.2.1. Determine the condition of the focal ecosystem before the extraction per unit area. The condition of the young stand before the thinning has been estimated to be 0.25 habitat hectares per hectare (Moilanen and Kotiaho, 2020).

3.5.2.2. Determine the condition of the focal ecosystem after the extraction per unit area. According to Mikkonen et al. (2023), first thinning reduces the ecological condition of the site by 50%. Thus, after the first thinning, the condition of the ecosystem is $0.50 \times 0.25 \text{ hha} = 0.125 \text{ habitat hectares per hectare}$.

3.5.2.3. Calculate the biodiversity impact per unit area by taking the difference between the condition before and after the extraction. The biodiversity impact of first thinning equals $0.25 \text{ hha} - 0.125 \text{ hha} = 0.125 \text{ habitat hectares per hectare of treated area}$.

3.5.3. Peat

The calculation of the biodiversity impact of peat as a fuel for district heating is very different from that of wood fuels. To obtain the peat, the biodiversity impact occurs once when the peatland is ditched and mining is started, but the same peat extraction site can be used for approximately 15–30 years (Väyrynen et al., 2008). In other words, the fuel is extracted and thus heat obtained for a long period of time from the same site, and when the biodiversity impact of the peat-fired district heating is calculated this needs to be taken into account.

Draining the peatland by ditching and removing the top layer of soil with all the vegetation while preparing the site for production causes the most significant impacts of peat mining (Finnish Ministry of the Environment, 2015). Therefore, the biodiversity impacts at the mining site itself do not increase after initial draining and removal of vegetation.

However, draining of peat extraction sites also affects the hydrology of surrounding peatlands (Finnish Ministry of the Environment, 2015). The extent of the draining effect depends on features like topography, peat layer depth, and drainage ditch depth (Paal et al., 2016). Moreover, a common requirement for peat extraction in Finland is that the peat layer is at least 1.5 m thick. This excludes e.g., the edges of the peatland from peat production. Nevertheless, ditching does drain these areas of the peatland as well, so this area in addition to the actual production area is degraded (Kareksela et al., 2013).

It is not exactly known how large proportion of each drained peatland is suitable for peat mining. We estimated that 1/2 of the drained area is either too shallow or otherwise unsuitable for peat mining (see Kareksela et al., 2013). This means that if one hectare of peatland is being used for peat extraction, another hectare of non-mineable peatland also gets degraded. Thus, the biodiversity impact caused by peat mining should include the impact of peat extraction as well as the impact of draining the non-mineable areas.

3.5.3.1. Determine the condition of the focal ecosystem before the extraction per unit area. Since no pristine peatlands are taken into use for peat production (Ministry of Agriculture and Forestry of Finland, 2012), we assumed the condition of a peatland before preparations for peat extraction to be 30% of its natural state, or 0.3 habitat hectares per hectare. This is the average condition reported for the drained peatlands suitable for wood production or peat mining in Finland (Kotiaho et al., 2015).

The ecological condition of the non-mineable area is similar to the mineable area before peat extraction, i.e., 0.3 habitat hectares per hectare. Note that if the area would be in a better condition before the extraction, the negative biodiversity impact would be higher than what is calculated below.

3.5.3.2. Determine the condition of the focal ecosystem after the extraction per unit area. The ecological condition of a peat extraction site is only 0.01 habitat hectares per hectare (Kotiaho et al., 2015). Since peat is not extracted from the non-mineable parts of the peatland, the ecological impact in non-mineable parts is likely to be less severe. We assumed the condition of non-mineable parts to decline a further 50% due to the draining. The ecological condition of drained non-mineable site is thus $0.3 \text{ hha/ha} \times 0.5 = 0.15 \text{ habitat hectares per hectare}$.

3.5.3.3. Calculate the biodiversity impact per unit area by taking the difference between the condition before and after the extraction. The ecological condition of the peat extraction site declines from 0.3 hha/ha before peat extraction to 0.01 hha/ha after extraction, and the impact is $0.3 - 0.01 = 0.29 \text{ habitat hectares per hectare of extraction site}$. In non-mineable parts, the biodiversity impact of draining is $0.3 - 0.15 = 0.15 \text{ habitat hectares per hectare}$. In reality it is likely that the more intense drainage degrades the area over some years, but to aid the calculation we make a simplifying assumption that the biodiversity impact of draining is full from the beginning of the peat extraction. Because one non-mineable hectare gets drained per each mined hectare, we need to take into account both of the two hectares to arrive to the biodiversity impact of mining in habitat hectares per hectare: $0.29 \text{ hha/ha} + 0.15 \text{ hha/ha} = 0.44 \text{ hha/ha}$.

As the peat is extracted and thus heat obtained for a long period of time from the same site with no additional biodiversity impacts, the biodiversity impacts of peat production should be divided by the peat production years. District heat is consumed in different quantities by many organizations every year, and the division is needed to allow the partitioning of the annual biodiversity impact among the organizations based on their annual consumption. The total biodiversity impact per extraction area hectare divided for the production years of the site (22.5 years on average, Väyrynen et al., 2008) equals $0.44 \text{ hha/ha} \div 22.5 \text{ years} = 0.020 \text{ habitat hectares per year per extraction area hectare}$.

3.6. Step 6. Quantify the total biodiversity impact of each fuel type consumption

Now that we know the total consumption of the heat at the focal organization (Step 1), fuel types and the share of each with which the heat is produced (Step 2), the energy yield per volumetric unit of each fuel type (Step 3), the volumetric yield of each fuel type per unit area extracted (Step 4), and the biodiversity impact of each fuel type per unit area extracted (Step 5), we can quantify the total biodiversity impact of each fuel type.

To achieve the energy yield per extracted area, the energy yield per volumetric unit (MWh/m^3) of each fuel type from Step 3 is multiplied by the volumetric yield of each fuel type per unit area extracted (m^3/ha) from Step 4 (Eq. (2)). To achieve the biodiversity impact per energy unit (hha/MWh) for each fuel type, the biodiversity impact per unit area extracted (hha/ha) obtained in Step 5 is divided by the energy yield per area extracted (MWh/ha), obtained with Eq. (1) (Eq. (3)). To achieve the biodiversity impact of energy consumption of each fuel type, biodiversity impact per energy unit (hha/MWh) is multiplied by the total consumption of energy produced with that fuel type (Eq. (4)). Note that by using the total consumption of energy produced with each fuel type, we do not need the information about the total area used for the extraction, although it could easily be obtained by dividing the total consumption of energy with the energy yield per area (not shown).

Table 1
Results of the calculation steps with the district heating data of the University of Jyväskylä in 2019–2021.

Step	Step description	Fuel type			
		Industry by-products	Logging residue chips	Chips from roundwood	Peat
Step 1	Total energy consumption* 2019: 22201 MWh 2020: 19681 MWh 2021: 26321 MWh				
Step 2	Energy consumption (MWh)** 2019 2020 2021	6964 7403 11,272	2019 2118 2030	1308 1992 4183	11,056 7912 8554
Step 3	Energy yield per unit (MWh/m^3)	–	2.0***	2.0***	0.91****
Step 4	Volumetric yield per unit area (m^3/ha)	–	55	45	439.6
Step 5	Biodiversity impact per unit area (hha/ha)	–	0.023	0.125	0.02
Step 3 * Step 4	Energy yield per area (MWh/ha)	–	110	90	400
Step 5 / Energy yield per area	Impact factor; Biodiversity impact per energy unit (hha/MWh)	–	0.00021	0.00139	0.00005
Step 6	Total impact (hha) allocated to district heating 2019 2020 2021	– – –	0.27 0.28 0.28	1.16 1.77 3.72	0.35 0.26 0.28

* There were some between-year differences in the buildings included in the total consumption, but this makes very little difference in the total consumption (more buildings included in more recent years; share of these additional buildings was 0,5% of total energy consumption in 2020). Also note that backup fuels (coal, oil, biogas) are included in the total energy consumption, but excluded from further investigations.

** University of Jyväskylä consumption data 2019–2021.

*** Energy yield per solid cubic meter. Alakangas and Impola, 2014.

**** Energy yield per loose cubic meter. Alakangas et al., 2016.

Finally, we need to take into account the share of the biodiversity impacts to be allocated to heat, which was 64% as explained in Step 2, Eq. (1). This is done by multiplying the result of Eq. (4) by 0.64.

All of these calculations need to be done separately for each fuel type i . The results are summarized in Table 1.

$$\text{Energy yield per area}_i \left(\frac{\text{MWh}}{\text{ha}} \right) = \text{Energy yield}_i \left(\frac{\text{MWh}}{\text{m}^3} \right) \times \text{Volumetric yield}_i \left(\frac{\text{m}^3}{\text{ha}} \right) \quad (2)$$

$$\text{Biodiversity impact per energy unit}_i \left(\frac{\text{hha}}{\text{MWh}} \right) = \frac{\text{Biodiversity impact per area}_i \left(\frac{\text{hha}}{\text{ha}} \right)}{\text{Energy yield per area}_i \left(\frac{\text{MWh}}{\text{ha}} \right)} \quad (3)$$

$$\text{Biodiversity impact}_i (\text{hha}) = \text{Biodiversity impact per energy unit}_i \left(\frac{\text{hha}}{\text{MWh}} \right) \times \text{Total energy consumption}_i (\text{MWh}) \quad (4)$$

In Eq. (5) we provide the biodiversity impact of energy consumption of fuel type i directly without intermediate steps from Eqs. (2), (3), and (4).

$$\text{Biodiversity impact}_i (\text{hha}) = \frac{\text{Biodiversity impact per area}_i \left(\frac{\text{hha}}{\text{ha}} \right)}{\text{Energy yield}_i \left(\frac{\text{MWh}}{\text{m}^3} \right) \times \text{Volumetric yield}_i \left(\frac{\text{m}^3}{\text{ha}} \right)} \times \text{Total energy consumption}_i (\text{MWh}) \quad (5)$$

The last two steps of the framework combine all different fuel types.

3.7. Step 7. Quantify the total biodiversity impact of district heating

To illustrate the total biodiversity impact of heat consumption, the impacts on the same ecosystem type can be added together because the biodiversity impact is being measured as the overall impact on the ecosystem with a common currency i.e. the habitat hectare. Adding impacts over the ecosystem types is also possible to illustrate the overall impact. If such a combination is done, one should be careful in the interpretation of the result as the combination loses the information on the identity of the ecosystem types some of which may be more vulnerable than others.

The biodiversity impact of district heating consumption of the University of Jyväskylä in 2021 was 4 habitat hectares in forest ecosystems and 0.28 habitat hectares in peatlands.

3.8. Step 8. Interpret the results

When interpreting the results, it is useful to analyse both the total biodiversity impact of each fuel type, and the biodiversity impact of each fuel type per energy unit (impact factor). This are shown in Fig. 4. The total biodiversity impact indicates the actual impact of the organization on biodiversity and can be used to e.g., formulate mitigation strategies. However, the impact factors are also important, as they can be used to

compare different energy sources to choose more sustainable alternatives.

The results can also be used for evaluating biodiversity offsetting actions to reach no-net-loss or net positive impact for biodiversity. In some cases, it might be beneficial to relate the results to the size of the organization, for example report results as biodiversity impact per employee or per unit area of premises.

It is also important to scrutinize the assumptions behind the calculations. Uncertainty can be present for example in determining the ecological condition of the ecosystems, which is based on expert reviews. Furthermore, the volumetric yield of the harvest can vary case by case. Nevertheless, if the logic behind the framework is robust, uncertainties in the collection of primary data can be reduced by further research. We also focused only on the biodiversity impacts from the land use and ignored the impacts that are due to e.g., emissions from the heat production and their effects on climate change. These and other matters will be further discussed from the viewpoint of our case study in the following section.

4. Conclusions

In this paper we presented a general protocol for assessing the biodiversity impact of peat- and wood-fired district heating. With




Fig. 4. A) The biodiversity impact (hha) of district heating use by fuel type, B) the consumption of each fuel type (MWh) in the University of Jyväskylä in 2019–2021 and C) the biodiversity impact factor (hha/MWh) of each fuel type.

adequate data, the protocol can be used to assess the biodiversity impact of any biofuel-based district heating system, or in fact any other industry where biomass is extracted and utilized. We adopted the ecosystem condition-based habitat hectare approach to make a holistic biodiversity impact assessment when the condition of the focal ecosystem is known before and after the extraction of the biomass. Compared to other potential methods to evaluate biodiversity impact, our framework is more detailed and focuses on the actual condition of the focal ecosystem. The framework proposed in this paper can be used anywhere in the world, but it is crucial that all the factors and details are adapted to the local context. This includes finding out all the necessary details of local energy production, as well as where and how the different fuels are extracted, by which methods, and what is the ecological impact caused by these practices at the location in question.

Of the fuel types analyzed in this study, peat had the most intensive biodiversity impact per unit area extracted (0.44 hha/ha), followed by roundwood chips (0.125 hha/ha) and logging residue chips (0.023 hha/ha). However, it is important to note, that energy peat is extracted annually from the same site on average for 22.5 years with no additional negative biodiversity impacts, and thus the average impact per hectare per annum decreases to 0.02 hha/ha. When we are looking at the biodiversity impact per energy unit, which may in fact be more relevant in the case of energy consumption, the order of the biodiversity impacts is the opposite: largest biodiversity impact (0.00139 hha/MWh) comes from firing roundwood chips, followed by firing logging residue chips (0.00021 hha/MWh), and firing peat had the least biodiversity impact at 0.000025 hha/MWh. Roundwood chips have 56 times greater and logging residue chips 8 times greater biodiversity impact per unit of energy than peat. One aspect that is yet to be considered is the time dimension of ecosystem recovery after the impact, which we have deliberately left out from this study. It can take a very long time for some ecosystems to recover after the biomass extraction, whereas others may recover much faster. If taken into account, this might change the results significantly.

To better understand the magnitude of the biodiversity impacts of district heating, we scaled the results to assess the biodiversity impact of total wood- and peat-fired district heating consumption in Finland. In 2021 a total of 40,798,000 MWh of district heating was produced in Finland, of which 42.1% was from wood fuels and 9.5% from peat fuels (Statistics Finland, 2023). When we apply the biodiversity impact factors derived in this study (see Table 1) by taking the average biodiversity impact factor for wood fuels (0.0008 hha/MWh) and utilizing directly the 0.00005 hha/MWh factor for peat fuels, the biodiversity impact would be around 13,700 habitat hectares in forests and 200 habitat hectares in peatlands. For comparison, on average 6500 ha of forest is voluntarily protected in Finland annually (Koskela et al., 2020). In conclusion, it seems that the biodiversity impact of district heating in Finland is remarkable.

This study also provides interesting insights for the political discussion between peat and wood-fired heating in Finland and Europe, which has been further fueled by the ongoing energy crisis in Europe. As part of their climate targets, the Finnish Government 2019 and Publications of the Finnish Government (2019) has been planning to halve the energy use of peat by 2030. Peat burning comprised of 18.9% of total district heating production in Finland in 2011 and 9.5% in 2021, indicating a steady decrease (Statistics Finland, 2023). In fact, peat production has dropped even faster than expected (News, 2022). However, at the same time there has been a steady increase in the use of wood fuels in Finnish district heating, climbing up from 22.3% in 2011 to 42.1% of total district heating production in 2021 (Statistics Finland, 2023). In the light of our results, this clear shift from peat to wood firing is likely to cause a significant increase in the negative biodiversity impacts of district heating.

Interestingly, the trade-off between the climate and biodiversity impacts of wood and peat firing is truly substantial. Peat is a non-renewable fuel and as such especially detrimental to climate like any other fossil fuel (Horsburgh et al., 2022). Milled peat emits around 108 t

of CO₂ per terajoule (Statistics Finland, 2024). In contrast, wood fuels are classified as renewable and carbon neutral, even though burning as such is by no means carbon neutral (Schulze et al., 2012). The emissions from wood fuels are estimated to be around 112 t of CO₂ per terajoule (Statistics Finland, 2024). However, burning wood is classified carbon neutral only because the emissions from wood fuels are allocated to the land use sector at the point of wood extraction and the emissions are assumed to be sequestered by regrowth of the renewable biomass. The classification of wood fuels as carbon neutral has been contested by scientists as it provides no incentive to avoid burning the wood (Norton et al., 2019). In our attempts to reach the carbon neutrality targets, the classification of wood fuels as carbon neutral has given rise to strategies where peat and other non-renewables are replaced by wood fuels. For example, the district heating company Alva, the target case of this study, now provides a product called “green heat” that is deemed environmentally friendly. This product consists mostly of wood fuels (Alva, 2020), which our results here show are far from environmentally friendly. In addition to climate impacts, future policies and sustainability strategies of energy producers should focus on biodiversity impacts of energy production methods. It is already well recognized that climate change and biodiversity loss require joint solutions and seeking for synergies and avoiding trade-offs (Pörtner et al., 2021), such as revealed by our results. One of the most influential solutions would naturally be to reduce the consumption of energy.

We focused here on land use impacts on biodiversity as it is one of the drivers, if not the key driver for biodiversity loss (IPBES, 2019). However, biodiversity loss is also driven by water use, overexploitation of natural resources, climate change, pollution, and invasive alien species (IPBES, 2019) among others, and in the future, these should also be considered when determining the most sustainable practices of energy production. Relevant for the current study is that it is clear that for both peat mining (Rodhe and Svensson, 1995; Kløve, 2001) and forest use (IPCC, 2019), other significant effects, such as climate change and pollution exist in addition to the biodiversity impacts caused by the fuel extraction processes.

District heating is a necessary part of infrastructure, and it will continue to be used in the future. However, it is important to consider the biodiversity impacts when planning and using district heating. In order to find the least harmful fuel composition for district heating, more research on the biodiversity impacts of different fuels as well as other energy production systems, including non-combustion energy production, are needed.

CRediT authorship contribution statement

Veera Vainio: Writing – original draft, Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Visualization, Project administration. **Sami El Geneidy:** Writing – original draft, Conceptualization, Methodology, Writing – review & editing, Supervision. **Panu Halme:** Writing – original draft, Conceptualization, Methodology, Writing – review & editing, Supervision, Visualization. **Maiju Peura:** Writing – review & editing, Methodology, Visualization. **Janne S. Kotiaho:** Writing – original draft, Conceptualization, Methodology, Writing – review & editing, Supervision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgements

This work was funded by the Strategic Research Council at the Academy of Finland (Kotiaho 345267). We would also like to thank Ulla Helimo and the Sustainability for JYU project members for their support and comments.

References

- Äijälä, O., Koistinen, A., Sved, J., Vanhatalo, K., Väisänen, P. (Eds.), 2014. Metsänhoidon suositukset. Metsätalouden kehittämiskeskus Tapion julkaisuja.
- Alakangas, E., Impola, R., 2014. Puupolttoaineiden laatuohje: VTT-M-07608-13 – päivitys 2014. Bioenergia Ry. Retrieved from. https://publications.vtt.fi/julkaisu/t/muut/2014/VTT-M-07608-13_2014_update.pdf.
- Alakangas, E., Hurskainen, M., Laatikainen-Luntama, J., Korhonen, J., 2016. Suomessa käytettävien polttoaineiden ominaisuuksia. VTT Technical Research Centre of Finland. VTT Technol. No. 258. Retrieved from. <https://publications.vtt.fi/pdf/technology/2016/T258.pdf>.
- Alanen, A., Aapala, K., 2015. Soidensuojelutyöryhmän ehdotus soidensuojelun täydentämiseksi. Ympäristöministeriön raportteja 26/2015. Retrieved from. <http://julkaisut.valtioneuvosto.fi/handle/10138/158285>.
- Alkemade, R., van Oorschot, M., Miles, L., Nellemann, C., Bakkenes, M., ten Brink, B., 2009. GLOBIOS: A Framework to Investigate Options for Reducing Global Terrestrial Biodiversity Loss. *Ecosystems* 12, 374–390. <https://doi.org/10.1007/s10021-009-9229-5>.
- Alva, 2020. Uusiutuva ja päästötöntä lämpöä. Retrieved from. <https://www.alva.fi/blog/2020/03/03/uusiutuva-ja-paastotonta-lampoa/>.
- Anttila, P., Nivala, V., Salminen, O., Hurskainen, M., Kärki, J., Lindroos, T.J., Asikainen, A., 2018. Regional balance of forest chip supply and demand in Finland in 2030. *Silva Fenn.* 52, 1–20.
- Asikainen, A., Ilvesniemi, H., Sievänen, R., Vapaavuori, E., Muhonen, T. (Eds.), 2012. Bioenergia, ilmastomuutos ja Suomen metsät. Working Papers of the Finnish Forest Research Institute, p. 240. Retrieved from. <http://urn.fi/URN:ISBN:978-951-40-2378-1>.
- Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M., Getz, W.M., Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A., Roopnarine, P., Vermeij, G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N., Mindell, D.P., Revilla, E., Smith, A.B., 2012 Jun 6. Approaching a state shift in Earth's biosphere. *Nature* 486 (7401), 52–58. <https://doi.org/10.1038/nature11018>. 22678279.
- Börjesson, P., Hansson, J., Berndes, G., 2017. Future demand for forest-based biomass for energy purposes in Sweden. *For. Ecol. Manag.* 383, 17–26.
- Bouget, C., Lassaune, A., Jonsell, M., 2012. Effects of fuelwood harvesting on biodiversity – a review focused on the situation in Europe. *Can. J. For. Res.* 42 (8), 1421–1432. <https://doi.org/10.1139/x2012-078>.
- Bull, J.W., Taylor, I., Biggs, E., Grub, H.M., Yearley, T., Waters, H., Milner-Gulland, E.J., 2022. Analysis: the biodiversity footprint of the University of Oxford. *Nature* 604 (7906), 420–424.
- Crenna, E., Marques, A., La Notte, A., Sala, S., 2020. Biodiversity assessment of value chains: state of the art and emerging challenges. *Environ. Sci. Technol.* 54 (16), 9715–9728.
- Creutzig, F., Ravindranath, N.H., Berndes, G., Bolwig, S., Bright, R., Cherubini, F., Masera, O., 2015. Bioenergy and climate change mitigation: an assessment. *GCB Bioenergy* 7 (5), 916–944.
- Curran, M., de Baan, L., De Schryver, A.M., van Zelm, R., Hellweg, S., Koellner, T., Sonnemann, G., Huijbregts, M.A.J., 2011. Toward meaningful end points of biodiversity in life cycle assessment. *Environ. Sci. Technol.* 45 (1), 70–79. <https://doi.org/10.1021/es101444k>.
- Damiani, M., Sinkko, T., Caldeira, C., Tosches, D., Robuchon, M., Sala, S., 2023. Critical review of methods and models for biodiversity impact assessment and their applicability in the LCA context. *Environ. Impact Assess. Rev.* 101, 107134.
- Ecke, F., Löfgren, O., Sörlin, D., 2002. Population dynamics of small mammals in relation to forest age and structural habitat factors in northern Sweden. *J. Appl. Ecol.* 39 (5), 781–792.
- El Geneidy, S., Alvarez, Franco D., Baumeister, S., Halme, P., Helimo, U., Kortetmäki, T., Kotiaho, J.S., Latva-Hakuni, E., Mäkelä, M., Raippalinnä, L.-M., Vainio, V., 2021. Sustainability for JYU: Jyväskylän yliopiston ilmasto- ja luontohaitat. Wisdom Letters. University of Jyväskylä, JYU.Wisdom - School of Resource Wisdom. <http://jyx.jyu.fi/handle/123456789/75182>.
- El Geneidy, S., Baumeister, S., Peura, M., Kotiaho, J.S., 2023. Value-Transforming Financial, Carbon and Biodiversity Footprint Accounting. Manuscript. Preprint available at. <https://doi.org/10.48550/arXiv.2309.14186>.
- Ellis, E.C., Kaplan, J.O., Fuller, D.Q., Vavrus, S., Goldewijk, K.K., Verburg, P.H., 2013. Used planet: a global history. *Proc. Natl. Acad. Sci.* 2013 (110), 7978–7985.
- Eräjää, S., Halme, P., Kotiaho, J.S., Markkanen, A., Toivanen, T., 2010. The volume and composition of dead wood on traditional and forest fuel harvested ClearCuts. *Silva Fenn.* 44 (2), 203–2011.
- Eriksson, O., Finnveden, G., Ekvall, T., Björklund, A., 2007. Life cycle assessment of fuels for district heating: a comparison of waste incineration, biomass- and natural gas combustion. *Energ Policy* 35 (2), 1346–1362.
- Finnish Government 2019, Publications of the Finnish Government, 2019. 33. Programme of Prime Minister Sanna Marin's Government 10 December 2019. In: Inclusive and Competent Finland - A Socially, Economically and Ecologically Sustainable Society. Retrieved from. <http://urn.fi/URN:ISBN:978-952-287-811-3>.
- Finnish Ministry of the Environment, 2015. Turvetuotannon ympäristönsuojeluohje. Ympäristöhallinnon ohjeita 2/2015. Retrieved from. <http://julkaisut.valtioneuvosto.fi/handle/10138/155221>.
- Flyktman, M., 2012. Energia- ja ympäristöturpeen kysyntä ja tarjonta vuoteen 2020 mennessä, 3. päivitys 3/2012. VTT:n Tutkimusraportteja, VTT-R-08372-11.
- Foley, J.A., Defries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005 Jul 22. 2005. Global consequences of land use. *Science* 309 (5734), 570–574. <https://doi.org/10.1126/science.1111772>. 16040698.
- Haapalehto, T., Kotiaho, J.S., Matilainen, R., Tahvanainen, T., 2014. The effects of long-term drainage and subsequent restoration on water table level and pore water chemistry in boreal peatlands. *J. Hydrol.* 519 (Part B), 1493–1505. <https://doi.org/10.1016/j.jhydrol.2014.09.013>.
- Hedin, J., Isacson, G., Jonsell, M., Komonen, A., 2008. Forest fuel piles as ecological traps for saproxylic beetles in oak. *Scand. J. For. Res.* 23 (4), 348–357.
- Horsburgh, N., Tyler, A., Mathieson, S., Wackernagel, M., Lin, D., 2022. Biocapacity and cost-effectiveness benefits of increased peatland restoration in Scotland. *J. Environ. Manag.* 306, 114486.
- Hyvärinen, E., Juslén, A.K., Kemppainen, E., Uddström, A., Liukko, U.M., 2019. Suomen lajien uhanalaisuus 2019-Punainen kirja: The 2019 Red List of Finnish Species.
- IPBES, 2018. The IPBES Assessment Report on Land Degradation and Restoration. Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany. <https://ipbes.net/assessment-reports/ldr>.
- IPBES, 2019. Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES Secretariat, Bonn, Germany. <https://ipbes.net/global-assessment>.
- IPCC, 2019. Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems [P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Peltz, J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)]. <https://www.ipcc.ch/srccl/>.
- Jalkanen, J., Kujala, H., Nieminen, E., Ahola, A., Pekkonen, M., Salo, P., 2023. Luontotyypin ekologisen tilan arviointi ekologisessa kompensaatiossa (draft). Retrieved from. <https://boostbiodiversityofsets.fi/luontotyypin-arviointimittarit/> (Accessed on 27.9.2023).
- Jaureguiberry, P., Titeux, N., Wiemers, M., Bowler, D.E., Coscieme, L., Golden, A.S., Guerra, C.A., Jacob, U., Takahashi, Y., Settele, J., Díaz, S., Molnár, Z., Purvis, A., 2022. The direct drivers of recent global anthropogenic biodiversity loss. *Sci. Adv.* 8 (45) <https://doi.org/10.1126/sciadv.abm9982>.
- Kangas, J., Majasalmi, T., Juva, K., Kotiaho, J.S., Ahlvi, L., 2023. Suomen luonnon tila ja tulevaisuus – skenaariotarkastelu luontokadon pysäyttämiseksi vaadittavista toimita. Suomen Luontopaneelin julkaisu 4B/2023. Available at. <https://luonto-paneeli.fi/wp-content/uploads/2023/12/suomen-luontopaneelin-julkaisu-4b-2023-suomen-luonnon-tila-ja-tulevaisuus-skenaariotarkastelu.pdf>.
- Kareksela, S., Moilanen, A., Tuominen, S., Kotiaho, J.S., 2013. Use of inverse spatial conservation prioritization to avoid biological diversity loss outside protected areas. *Conserv. Biol.* 27, 1294–1303.
- Kareksela, S., Haapalehto, T., Juutinen, R., Matilainen, R., Tahvanainen, T., Kotiaho, J.S., 2015. Fighting carbon loss of degraded peatlands by jump-starting ecosystem functioning with ecological restoration. *Sci. Total Environ.* 537, 268–276.
- Karttunen, K., Laitila, J., Ranta, T., 2016. First-thinning harvesting alternatives for industrial or energy purposes based on regional Scots pine stand simulations in Finland. *Silva Fenn* 50 (2), 16. <https://doi.org/10.14214/sf.1521> article id 1521.
- Kløve, B., 2001. Characteristics of nitrogen and phosphorus loads in peat mining wastewater. *Water Res.* 35 (10), 2353–2362.
- Koistinen, A., Luuro, J.-P., Vanhatalo, K. (Eds.), 2019. Metsänhoidon suositukset energiapuun korjuuseen, työopas. Tapion julkaisuja. Retrieved from. <https://tapio.fi/wp-content/uploads/2020/10/Metsanhoidon-suositukset-energiapuun-korjuuseen-Tapio-20191230.pdf>.
- Konar, M., Todd, J., Muneepeerakul, R., Rinaldo, A., Rodriguez-Iturbe, I., 2013. Hydrology as a driver of biodiversity: controls on carrying capacity, niche formation, and dispersal. *Adv. Water Resour.* 51, 317–325. <https://doi.org/10.1016/j.advwatres.2012.02.009>.
- Kontula, T., Raunio, A., 2019. Threatened Habitat Types in Finland 2018 - Red List of Habitats Results and Basis for Assessment.
- Kortetmäki, T., Puurtinen, M., Salo, M., Aro, R., Baumeister, S., Duflo, R., Elo, M., Halme, P., Husu, H.-M., Huttunen, S., Hyvönen, K., Karkulehto, S., Kataja-aho, S., Keskinen, K.E., Kulmunkki, I., Mäkinen, T., Näyhä, A., Okkolin, M.-A., Perälä, T., JYU. Wisdom community, 2021. Planetary well-being. *Humanities and Social Sciences Communications* 8 (1). <https://doi.org/10.1057/s41599-021-00899-3>. Article 1.
- Koskela, T., Anttila, S., Simkin, J., Aapala, K., Syrjänen, K., 2020. METSO-tilannekatsaus 2019: Etelä-Suomen metsien monimuotoisuuden toimintaohjelma 2008–2025. Luonnonvara- ja biotalouden tutkimus 36/2020. Luonnonvarakeskus. Retrieved from. <http://urn.fi/URN:ISBN:978-952-326-977-4>.
- Kotiaho, J.S., Kuusela, S., Nieminen, E., Päivinen, J., 2015. Elinympäristöjen tilan edistäminen Suomessa. Suomen ympäristö. 8/2015. <http://urn.fi/URN:ISBN:978-952-11-4462-2>.
- Kotiaho, J.S., Kuusela, S., Nieminen, E., Päivinen, J., 2016. Framework for assessing and reversing ecosystem degradation. In: Reports of the Ministry of the Environment 15en/2016. <http://urn.fi/URN:ISBN:978-952-11-4603-9>.

- Lammerant, J., Starkey, M., De Horde, A., Bor, A.-M., Driesen, K., Vanderheyden, G., 2021. Assessment of Biodiversity Measurement Approaches for Business and Financial Institutions. EU Business @ Biodiversity Platform. Update Report 3. Retrieved from. https://knowledge4policy.ec.europa.eu/publication/assessment-bio-diversity-measurement-approaches-businesses-financial-institutions-update_en.
- Lauhanen, R., Ahokas, J., Esala, J., Hakonen, T., Sippola, H., Viirimäki, J., Koskinemi, E., Laurila, J., Makkonen, I., 2014. Metsätoimihenkilöiden energialaskuoppi. Seinäjoen ammattikorkeakoulun julkaisusarja C. Oppimateriaaleja 6. <https://www.theseus.fi/bitstream/handle/10024/80849/C6.pdf?sessionid=0B0997D2CC05404289A1AF5130416959?sequence=1>.
- Lenzen, M., 2014. An outlook into a possible future of footprint research. *J. Ind. Ecol.* 18 (1), 4–6. <https://doi.org/10.1111/JIEC.12080>.
- Lindqvist, M., Palme, U., Lindner, J.P., 2016. A comparison of two different biodiversity assessment methods in LCA—a case study of Swedish spruce forest. *Int. J. Life Cycle Assess.* 21 (2), 190–201. <https://doi.org/10.1007/s11367-015-1012-6>.
- Mandley, S.J., Daiglou, V., Junginger, H.M., van Vuuren, D.P., Wicke, B., 2020. EU bioenergy development to 2050. *Renew. Sust. Energ. Rev.* 127, 109858.
- Marques, A., Verones, F., Kok, M.T., Huijbregts, M.A., Pereira, H.M., 2017. How to quantify biodiversity footprints of consumption? A review of multi-regional input-output analysis and life cycle assessment. *Curr. Opin. Environ. Sustain.* 29, 75–81. <https://doi.org/10.1016/j.cosust.2018.01.005>.
- Masson-Delmotte, V., Zhai, P., Pörtner, H.O., Roberts, D., Skea, J., Shukla, P.R., Waterfield, T., 2018. IPCC special report 1.5-summary for policymakers. Global warming of 1.5° C. An IPCC special report on the impacts of global warming of 1.5° C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.
- Mazerolle, M.J., 2003. Detrimental effects of peat mining on amphibian abundance and species richness in bogs. *Biol. Conserv.* 113 (2), 215–223.
- Mazhar, A.R., Liu, S., Shukla, A., 2018. A state of art review on the district heating systems. *Renew. Sust. Energ. Rev.* 96, 420–439.
- Metsälehti, 2023. Energiapuun hinnat. Retrieved from. <https://www.metsalehti.fi/puunhinta/energiapuun-hinnat/> (Accessed on 27.9.2023).
- Mikkonen, N., Leikola, N., Lehtomäki, J., Halme, P., Moilanen, A., 2023. National high-resolution conservation prioritisation of boreal forests. *For. Ecol. Manag.* 541, 121079 <https://doi.org/10.1016/j.foreco.2023.121079>.
- Ministry of Agriculture and Forestry of Finland, 2012. Valtioneuvoston periaatepäätös soiden ja turvemaiden kestävästä ja vastuullisesta käytöstä ja suojelusta [In Finnish]. Retrieved from. https://mmm.fi/documents/1410837/1516663/MMM-119690-v5-suostrategia_valtioneuvoston_periaatepaatos_v4/005425e8-e3c4-497d-8c9f-26f343896c37.
- Moilanen, A., Kotiaho, J.S., 2018. Fifteen operationally important decisions in the planning of biodiversity offsets. *Biol. Conserv.* 227, 112–120.
- Moilanen, A., Kotiaho, J.S., 2020. Liite 18: Vapaaehtoinen ekologinen kompensatio AA Sakatti Mining Oy:n mahdolliselle Sakatin kaivokselle. Liite ympäristövaikutusten arviointiin. Retrieved from. <https://www.ymparisto.fi/sakatinkaivosyva>.
- Moilanen, A., Kotiaho, J.S., 2021. Three ways to deliver a net positive impact with biodiversity offsets. *Conserv. Biol.* 35 (5), 197–205. <https://doi.org/10.1111/cobi.13533>.
- Newbold, T., Hudson, L.N., Hill, S.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverría-Londoño, S., Edgar, M.J., Feldman, A., Garon, M., Harrison, M.L., Alhousseini, T., Ingram, D.J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D. L., Martin, C.D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H.R., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White, H.J., Ewers, R.M., Mace, G. M., Scharlemann, J.P., Purvis, A., 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520 (7545), 45–50. <https://doi.org/10.1038/nature14324>. PMID: 25832402.
- News, Yle, 2022. Peat production drops faster than expected in Finland – but may be on the way back. Retrieved from. <https://yle.fi/a/3-12409412>.
- Norton, M., Bald, A., Buda, V., Carli, B., Cudlin, P., Jones, M.B., Korhola, A., Michalski, R., Novo, F., Oszlányi, J., Santos, F.D., Schink, B., Shepherd, J., Vet, L., Walloe, L., Wijkman, A., 2019. Serious mismatches continue between science and policy in forest bioenergy. *GCB Bioenergy* 11 (11), 1256–1263. <https://doi.org/10.1111/gcbb.12643>.
- Official Statistics of Finland (OSF), 2018. Area Treated with Fellings by Region [Web Publication]. Natural Resources Institute Finland. Retrieved from. <https://statdb.luke.fi/443/PxWeb/sq/513054e6-57fb-45b5-a225-bd3cf0a7584d>.
- Official Statistics of Finland (OSF), 2021. Solid Wood Fuel Consumption in Heating and Power Plants by Region [Web Publication]. Natural Resources Institute Finland. Retrieved from. <https://statdb.luke.fi/443/PxWeb/sq/37268c1d-5e5c-4cec-83bf-6e0a48a45c7a>.
- Oy, Alva-yhtiöt, 2022. Alva – Company Presentation and Social Responsibility Report 2021. Retrieved from. <https://www.alva.fi/app/uploads/1/2022/08/Alva-yhteisku-ntavastuuraportti-2021-EN.pdf>.
- Oy, Alva-yhtiöt, 2023. Tilinpäätös ja toimintakertomus, tilikausi 1.1.2022–31.12.2022. Retrieved from. https://www.alva.fi/app/uploads/1/2023/04/Alva-yhtiöt_tilinpäätös-2022.pdf.
- Paal, J., Jürjendal, I., Suija, A., Kull, A., 2016. Impact of drainage on vegetation of transitional mires in Estonia. *Mires Peat.* 18, 1–19.
- Paiho, S., Saastamoinen, H., 2018. How to develop district heating in Finland? *Energy Policy* 122, 668–676.
- Parkes, D., Newell, G., Cheal, D., 2003. Assessing the quality of native vegetation: the ‘habitat hectares’ approach. *Ecol. Manag. Restor.* 4, S29–S38.
- Pérez-Lombard, L., Ortiz, J., Pout, C., 2008. A review on buildings energy consumption information. *Energ. Build.* 40 (3), 394–398.
- Persson, U., Werner, S., 2011. Heat distribution and the future competitiveness of district heating. *Appl. Energy* 88 (3), 568–576.
- Pörtner, H.O., Scholes, R.J., Agard, J., Archer, E., Arnett, A., Bai, X., Barnes, D., Burrows, M., Chan, L., Cheung, W.L., Diamond, S., Donatti, C., Duarte, C., Eisenhauer, N., Foden, W., Gasalla, M.A., Handa, C., Hickler, T., Hoegh-Guldberg, O., Ichii, K., Jacob, U., Inzarov, G., Kiessling, W., Leadley, P., Leemans, R., Levin, L., Lim, M., Maharaj, S., Managi, S., Marquet, P.A., McElwee, P., Midgley, G., Oberdorff, T., Obura, D., Osman, E., Pandit, R., Pascual, U., Pires, A.P.F., Popp, A., Reyes-García, V., Sankaran, M., Settele, J., Shin, Y.J., Sintayehu, D.W., Smith, P., Steiner, N., Strassburg, B., Sukumar, R., Trisos, C., Val, A.L., Wu, J., Aldrian, E., Parmesan, C., Pichs-Madruga, R., Roberts, D.C., Rogers, A.D., Diaz, S., Fischer, M., Hashimoto, S., Lavorel, S., Wu, N., Ngo, H.T., 2021. IPBES-IPCC co-sponsored workshop report on biodiversity and climate change. IPBES IPCC. <https://doi.org/10.5281/zenodo.4782538>.
- Ranius, T., Hämäläinen, A., Egnell, G., Olsson, B., Eklöf, K., Stendahl, J., Felton, A., 2018. The effects of logging residue extraction for energy on ecosystem services and biodiversity: a synthesis. *J. Environ. Manag.* 209, 409–425.
- Rodhe, H., Svensson, B., 1995. Impact on the greenhouse effect of peat mining and combustion. *Ambio* 221–225.
- Rosen, M.A., 2008. Allocating carbon dioxide emissions from cogeneration systems: descriptions of selected output-based methods. *J. Clean. Prod.* 16 (2), 171–177.
- Schulze, E.-D., Körner, C., Law, B.E., Haberl, H., Luysaert, S., 2012. Large-scale bioenergy from additional harvest of forest biomass is neither sustainable nor greenhouse gas neutral. *Glob. Change Biol. Bioenergy* 4, 611–616. <https://doi.org/10.1111/j.1757-1707.2012.01169.x>.
- Statistics Finland, 2023. Production of district heat in Finland, 2000–2021. Retrieved from. <https://pxdata.stat.fi/443/PxWeb/sq/85a92b3d-a5d2-4b2b-a56d-b755c6f9da55>.
- Statistics Finland, 2024. Fuel classification 2024. Retrieved from. https://stat.fi/media/uploads/tup/khkinv/khkaasut_polttoaineluokitus_2024.xlsx.
- Szarka, N., Eichhorn, M., Kittler, R., Bezama, A., Thrän, D., 2017. Interpreting long-term energy scenarios and the role of bioenergy in Germany. *Renew. Sust. Energ. Rev.* 68, 1222–1233.
- Tahvanainen, T., 2011. Abrupt ombrotrophication of a boreal peat mire triggered by hydrological disturbance in the catchment. *J. Ecol.* 99 (2), 404–415.
- Taylor, I., Bull, J.W., Ashton, B., Biggs, E., Clark, M., Gray, N., Milner-Gulland, E.J., 2023. Nature-positive goals for an organization’s food consumption. *Nature Food* 1–13.
- Thiffault, E., Béchard, A., Paré, D., Allen, D., 2016. Recovery rate of harvest residues for bioenergy in boreal and temperate forests: A review. *Adv. Bioenergy Sustain. Challenge* 293–316.
- Toivanen, T., Markkanen, A., Kotiaho, J.S., Halme, P., 2012. The effect of forest fuel harvesting on the fungal diversity of clear-cuts. *Biomass Bioenergy* 39, 84–93. <https://doi.org/10.1016/j.biombioe.2011.11.016>.
- Väyrynen, T., Aaltonen, R., Haavikko, H., Juntunen, M., Kalliokoski, K., Niskala, A.-L., Tukiainen, O., 2008. Turvetuotannon ympäristönsuojeluopas. Pohjois-Pohjanmaan Ympäristökeskus.
- Verkerk, P.J., Fitzgerald, J.B., Datta, P., Dees, M., Hengeveld, G.M., Lindner, M., Zudin, S., 2019. Spatial distribution of the potential forest biomass availability in Europe. *For. Ecosyst.* 6, 5. <https://doi.org/10.1186/s40663-019-0163-5>.
- Verones, F., Hellweg, S., Antón, A., Azevedo, L.B., Chaudhary, A., Cosme, N., Huijbregts, M.A., 2020. LC-IMPACT: A regionalized life cycle damage assessment method. *J. Ind. Ecol.* 24 (6), 1201–1219. <https://doi.org/10.1111/jiec.1301>.
- Viitaasaari, T., 2013. Energiapuuta päätehakkuulta -opas. Suomen metsäkeskus. Retrieved from. <https://www.metsakeskus.fi/sites/default/files/energiapuuta-paatehakkuulta-opas.pdf>.
- Weldu, Y.W., Assefa, G., 2016. Evaluating the environmental sustainability of biomass-based energy strategy: using an impact matrix framework. *Environ. Impact Assess. Rev.* 60 (9), 75–82.
- Werner, S., 2017. International review of district heating and cooling. *Energy* 137, 617–631.
- Wiedmann, T., Minx, J., 2008. A definition of ‘carbon footprint’. *Ecol. Econ. Res. Trends* 1 (2008), 1–11.
- Winter, L., Lehmann, A., Finogenova, N., Finkbeiner, M., 2017. Including biodiversity in life cycle assessment – State of the art, gaps and research needs. *Environ. Impact Assess. Rev.* 67 (11), 88–100.