

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Arya, Vedansh; Julin, Vesa

Title: Hölder continuity and Harnack estimate for non-homogeneous parabolic equations

Year: 2024

Version: Published version

Copyright: © The Author(s) 2024

Rights: CC BY 4.0

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Arya, V., & Julin, V. (2024). Hölder continuity and Harnack estimate for non-homogeneous parabolic equations. Mathematische Annalen, Early online. https://doi.org/10.1007/s00208-024-02979-6



Hölder continuity and Harnack estimate for non-homogeneous parabolic equations

Vedansh Arya¹ · Vesa Julin¹

Received: 21 May 2024 / Revised: 7 August 2024 / Accepted: 15 August 2024 © The Author(s) 2024

Abstract

In this paper we continue the study on intrinsic Harnack inequality for nonhomogeneous parabolic equations in non-divergence form initiated by the first author in Arya (Calc Var Partial Differ Equ 61:30–31, 2022). We establish a forward-in-time intrinsic Harnack inequality, which in particular implies the Hölder continuity of the solutions. We also provide a Harnack type estimate on global scale which quantifies the strong minimum principle. In the time-independent setting, this together with Arya (2022) provides an alternative proof of the generalized Harnack inequality proven by the second author in Julin (Arch Ration Mech Anal 216:673–702, 2015).

Mathematics Subject Classification 35K55 · 35B45

1 Introduction and the statement of the main results

In this paper we consider parabolic equation of the type

$$F(D^{2}u, Du, x, t) - u_{t} = 0,$$
(1.1)

where *F* is uniformly elliptic w.r.t. the Hessian and has a nonlinear growth w.r.t. the gradient. More precisely, we assume that there exist constants $0 < \lambda \le \Lambda$ such that

$$\lambda Tr(N) \le F(M+N, p, x, t) - F(M, p, x, t) \le \Lambda Tr(N)$$

for all symmetric matrices M, N with $N \ge 0$ and for every $(p, x, t) \in \mathbb{R}^n \times Q_r(x_0, t_0)$. Here $Q_r(x_0, t_0)$ is a cube centered at $(x_0, t_0) \in \mathbb{R}^{n+1}$. We assume further that F has

Vesa Julin vesa.julin@jyu.fi
 Vedansh Arya vedansh.v.arya@jyu.fi

¹ Department of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, Finland

the following growth in the gradient variable,

$$|F(0, p, x, t)| \le \phi(|p|) \tag{1.2}$$

for every $(p, x, t) \in \mathbb{R}^n \times Q_r(x_0, t_0)$, where $\phi : [0, \infty) \to [0, \infty)$ is of the form $\phi(s) = \eta(s)s$ and satisfies the following conditions:

(P1) $\phi : [0, \infty) \to [0, \infty)$ is increasing, locally Lipschitz continuous in $(0, \infty)$ and $\phi(s) \ge s$ for every $s \ge 0$. Moreover, $\eta : (0, \infty) \to [1, \infty)$ is nonincreasing on (0, 1) and nondecreasing on $[1, \infty)$;

(P2) η satisfies

$$\lim_{s \to \infty} \frac{s\eta'(s)}{\eta(s)} \log(\eta(s)) = 0;$$
(1.3)

(P3) There is a constant Λ_0 such that

$$\eta(s_1 s_2) \le \Lambda_0 \eta(s_1) \eta(s_2); \tag{1.4}$$

for every $s_1, s_2 \in (0, \infty)$.

One example of F is

$$F(D^{2}u, Du, x, t) = \operatorname{Tr}(A(x)D^{2}u) + \log(|Du|)\langle Dp(x), Du\rangle$$

where A(x) is a symmetric matrix with eigenvalues between $\lambda > 0$ and Λ , and $p \in C^1(\mathbb{R}^n)$. This operator is similar to the normalized p(x)-Laplacian, see [9].

The elliptic case is studied by the second author in [9], and there the main problem is to find a Harnack type estimate which quantifies the strong minimum principle whenever it is true and generalizes the Harnack inequality from the homogeneous setting [4, 10]. The related boundary problem is studied in [2, 12, 13]. In this paper we first discuss the assumptions (P1)-(P3). In particular, the condition (P2) in (1.3) might first seem rather resctrictive. We note that (P2) roughly states that the function η is *slowly varying*¹ in a quantitative way and, in particular, it implies the slow growth estimate on η , i.e., for every $\varepsilon > 0$ there is C_{ε} such that $\eta(s) \leq C_{\varepsilon}s^{\varepsilon}$ for $s \geq 1$ [3, Proposition 1.3.6]. Our first result (Theorem 1.1) justifies the assumption (P2) in the sense that we show that the Harnack estimate in [9] is false if the non-linearity is of the form $\eta(s) = \max\{s^{\varepsilon}, 1\}$ for any $\varepsilon > 0$.

The parabolic case is studied by the first author in [1], where the main result is the backward-in-time intrinsic Harnack inequality. The parabolic case is the main focus of this paper and we first provide the analogous forward-in-time intrinsic Harnack inequality (Theorem 1.2), and show then that this intrinsic Harnack inequality implies the Hölder continuity of the solutions (Theorem 1.3). The problem has some similarities to the p-parabolic equation which is also non-nomogeneous, and where the

¹ A positive function η is slowly varying if for all $\lambda > 0$ it holds $\lim_{s \to \infty} \frac{\eta(\lambda s)}{\eta(s)} = 1$.

concept of intrinsic Harnack inequality is developed [6, 7] (see also [11]). The difference is that in the p-parabolic equation the elliptic operator has a different scaling than the time derivative term, while in (1.1) it is the elliptic operator that does not have any homogeneity properties. Therefore, from the point of view of the scaling, the Eq. (1.1) is more similar to the reaction diffusion equation than to the p-parabolic. However, to the best our knowledge these results are not known for equations of type

$$u_t = F(D^2 u) + f(u).$$

The intrinsic Harnack inequality is a result on intrinsic scale and it is a relevant question if one can have a generalized Harnack inequality for non-negative solutions of (1.1) similar to the elliptic one in [9]. In our last result (Theorem 1.4) we provide a Harnack type estimate for non-negative solutions of (1.1) on global scale which quantifies the strong minimum principle.

1.1 Statement of the main results

In order to state our main result we first simplify the setting and, following the ideas of Caffarelli [4, 5], replace the Eq. (1.1) by two extremal inequalities which take into account the ellipticity assumption and the growth condition of the drift term. To be more precise, we assume that $u \in C(Q_2(x_0, t_0))$ is a viscosity supersolution of

$$P_{\lambda,\Lambda}^{-}(D^{2}u) - u_{t} \le \phi(|Du|) \tag{1.5}$$

and a viscosity subsolution of

$$P_{\lambda,\Lambda}^+(D^2u) - u_t \ge -\phi(|Du|),\tag{1.6}$$

where $P_{\lambda,\Lambda}^{\pm}$ denotes the extremal Pucci operators defined later in (2.1). We refer to Sect. 2 for the precise notion of viscosity sub- and supersolutions. The elliptic counterpart to (1.5) and (1.6) is defined analogously.

Throughout the paper we say that a constant *C* is *universal* if it depends only on the ellipticity constants, the nonlinearity ϕ and on the dimension *n*.

We recall that in [9] it is proven that in the elliptic case if $u \in C(B_2)$ is a nonnegative viscosity supersolution of $P_{\lambda,\Lambda}^-(D^2u) \le \phi(|Du|)$ and a viscosity subsolution $P_{\lambda,\Lambda}^+(D^2u) \ge -\phi(|Du|)$ in B_2 , then there is a universal constant C such that for $m = \inf_{B_1} u$ and $M = \sup_{B_1} u$ it holds

$$\int_{m}^{M} \frac{ds}{\phi(s)} \le C. \tag{1.7}$$

It is not difficult to see that if u is as above, then it satisfies the strong minimum principle when ϕ satisfies the so called Osgood condition [9]

$$\int_0^1 \frac{ds}{\phi(s)} = \infty. \tag{1.8}$$

🖄 Springer

Therefore it is clear that (1.7) quantifies the strong minimum principle when (1.8) holds.

Our first result justifies the assumption (P2) in (1.3).

Theorem 1.1 Assume $n \ge 3$ and fix any $\varepsilon_0 > 0$. Then there exist $\lambda, \Lambda > 0$ and a sequence of positive functions $u_k \in C^2(B_2)$ such that

$$\left|P_{\lambda,\Lambda}^{-}(D^{2}u_{k})\right| \leq |Du_{k}|^{1+\varepsilon_{0}}$$
 point-wise in B_{2} ,

while $m_k = \inf_{B_1} u_k \to 0$ as $k \to \infty$ and $M_k = \sup_{B_1} u_k \ge 1$ for all $k \in \mathbb{N}$. In particular, it holds

$$\int_{m_k}^{M_k} \frac{ds}{s^{1+\varepsilon_0}+s} \to \infty \quad \text{as } k \to \infty.$$

We note that the condition (P2) in (1.3) is an assumption only on the asymptotic behavior of the nonlinearity ϕ and plays no role whether the Osgood condition (1.8) holds or not. The point of Theorem 1.1 is that too weak information on the asymptotic growth of ϕ is the reason for the Harnack estimate to fail, even if the Harnack estimate means to quantify only the strong minimum principle.

Next we turn our attention to the parabolic setting (1.1), which is the main focus of this paper. We recall the following parabolic Harnack type inequality proven in [1] for solutions to (1.1) in a suitable intrinsic geometry corresponding to the nonlinearity. Let $u \in C(Q_2(x_0, t_0))$ be a positive viscosity supersolution of (1.5) and viscosity subsolution of (1.6). There is a universal constant C > 0 such that

$$\sup_{A_{\rho}^{-}(x_{0},t_{0})} u(x,t) \leq Cu(x_{0},t_{0}) \text{ for } \rho \leq a_{0} := \frac{u(x_{0},t_{0})}{C(\phi(u(x_{0},t_{0}))+u(x_{0},t_{0}))},$$
(1.9)

where $A_{\rho}^{-}(x_0, t_0) = \left\{ (x, t) : |x - x_0|_{\infty} \le \frac{\rho c_n}{2}, -\rho^2 + \frac{(\rho c_n)^2}{4} \le t - t_0 \le -\rho^2 + \frac{(\rho c_n)^2}{2} \right\}$ and $|\cdot|_{\infty}$ is the l^{∞} -norm. Here $c_n \le 1$ depends only on *n*. The estimate (1.9) is an intrinsic Harnack estimate as the scaling on the LHS depends on the value of the solution at the base point which is similar to [7].

Our next result is the forward-in-time counterpart of (1.9) and thus we obtain the complete intrinsic Harnack estimate for (1.1). To this aim we denote

$$A_{\rho}^{+}(x_{0}, t_{0}) := \left\{ (x, t) : |x - x_{0}|_{\infty} < \frac{\rho c_{n}}{2}, \rho^{2} - \frac{(\rho c_{n})^{2}}{2} < t - t_{0} \le \rho^{2} - \frac{(\rho c_{n})^{2}}{4} \right\},$$
(1.10)

where $c_n \leq 1$ is as above.

Theorem 1.2 Let $u \in C(Q_2)$ be a non-negative viscosity supersolution of (1.5) and viscosity subsolution of (1.6). There is a universal constant C > 0 such that if

 $u(x_0, t_0) > 0$ and $Q_{2\rho}(x_0, t_0) \cup A^+_{\rho}(x_0, t_0) \subset Q_2$ for

$$\rho \le \alpha_0 := \frac{u(x_0, t_0)}{C(\phi(u(x_0, t_0)) + u(x_0, t_0))}$$

then it holds

$$u(x_0, t_0) \le C \inf_{A_{\rho}^+(x_0, t_0)} u(x, t).$$
(1.11)

The proof of Theorem 1.2 relies on (1.9) and a careful continuity-type argument.

As in the p-parabolic case [6, 7], using the estimate (1.11) we obtain the Hölder continuity of the solutions of (1.1).

Theorem 1.3 Let $u \in C(Q_2)$ be a viscosity supersolution of (1.5) and viscosity subsolution of (1.6). Then, there exists a universal constant $\alpha \in (0, 1)$ such that u is locally α -Hölder continuous. More precisely, there is a universal constant C > 1, such that for all $(x, t), (y, s) \in Q_1$, we have

$$|u(x,t) - u(y,s)| \le C\phi(||u||_{L^{\infty}(Q_2)})(|x-y| + |t-s|)^{\alpha}.$$
(1.12)

We note that in (1.12) the exponent $\alpha \in (0, 1)$ is universal, but the Hölder norm depends on the solution u in a nonlinear way. This is necessary already in the elliptic case [14]. Our proof for Theorem 1.3 draws inspiration from the p-parabolic case [7]. However, since the elliptic operator in the equation (1.1) does not have any degree of homogeneity, the intrinsic cylinders in Theorem 1.2 may not be quantitatively monotone in size. This causes challenges which we need to overcome in order to obtain a uniform $\alpha > 0$.

Finally, we study whether we may obtain a Harnack type estimate on global scale, which quantifies the strong minimum principle similar to (1.7). Recall that the strong minimum principle is related to the Osgood condition (1.8).

Theorem 1.4 Let $u \in C(Q_4(0, 1))$ be a non-negative viscosity supersolution of (1.5) and viscosity subsolution of (1.6) in $Q_4(0, 1)$. There exist a universal constant C > 1and time levels $t_1, t_2 \ge 0$, depending on the value u(0, 0) such that $\frac{1}{C(\eta(u(0,0))+1)^2} \le t_1, t_2 \le 1$, so that it holds

$$\int_{\inf_{B_1} u(\cdot, t_1)}^{\sup_{B_1} u(\cdot, -t_2)} \frac{ds}{\phi(s)} \le C.$$
(1.13)

The parabolic Harnack estimate (1.13) takes the same form as the elliptic one (1.7), with the difference that here we have the waiting times t_1 , t_2 as usual with parabolic equations [8, 15]. Since the scaling of the Eq. (1.1) does not have any monotonicity properties, we can only give an estimate for the waiting times. However, the proof of Theorem 1.4 implies that, under the Osgood-condition (1.8), if u(0, 0) = 0 then the waiting time t_2 can be chosen zero. We state this in the following corollary, which is a quantification of the strong minimum principle at the time level t = 0.

Corollary 1.5 Let $u \in C(Q_4)$ be as in Theorem 1.4 and assume that ϕ satisfies (1.8). If u(0, 0) = 0 then

$$u(x, 0) = 0$$
 for all $x \in B_1$.

In the next section we introduce some basic notations and in Sect. 3 we prove our main results.

2 Notations and preliminaries

We denote a point in space by $x \in \mathbb{R}^n$ and in space-time by $(x, t) \in \mathbb{R}^n \times \mathbb{R}$. We denote the Euclidean norm of x in \mathbb{R}^n by |x| and the l^{∞} -norm by $|x|_{\infty} = \max\{|x_1|, |x_2|, \ldots, |x_n|\}$. We denote Df the gradient of f in x, f_t with respect to time t and $D^2 f$ denotes the Hessian matrix of f with respect to x. We denote the ball of radius r centered at x_0 by $B_r(x_0)$ and its closure by $\overline{B}_r(x_0)$. A cube of radius ρ centered at (x_0, t_0) is defined as

$$Q_{\rho}(x_0, t_0) = \{x \in \mathbb{R}^n : |x - x_0|_{\infty} < \rho\} \times (t_0 - \rho^2, t_0].$$

We denote B_r and Q_r if the ball and the cube are centered at the origin.

Let *S* be the space of real $n \times n$ symmetric matrices. We recall the definition of Pucci's extremal operators (for more detail see [5]). For $M \in S$, Pucci's extremal operators with ellipticity constant $0 < \lambda \leq \Lambda$ are defined as

$$P_{\lambda,\Lambda}^{-}(M) = \lambda \sum_{e_i > 0} e_i + \Lambda \sum_{e_i < 0} e_i,$$

$$P_{\lambda,\Lambda}^{+}(M) = \Lambda \sum_{e_i > 0} e_i + \lambda \sum_{e_i < 0} e_i,$$
(2.1)

where e_i 's are the eigenvalues of M.

We recall the definition of a viscosity supersolution of (1.5) and viscosity subsolution of (1.6).

Definition 2.1 A lower semicontinuous function $u : Q_r(x_0, t_0) \to \mathbb{R}$ is a viscosity supersolution of (1.5) in $Q_r(x_0, t_0)$ if the following holds: if $(x, t) \in Q_r(x_0, t_0)$ and $\varphi \in C^2(Q_r(x_0, t_0))$ are such that $\varphi \le u$ and $\varphi(x, t) = u(x, t)$ then

$$P^{-}_{\lambda,\Lambda}(D^{2}\varphi(x,t)) - \varphi_{t}(x,t) \leq \phi(|D\varphi(x,t)|).$$

An upper semicontinuous function $u : Q_r(x_0, t_0) \to \mathbb{R}$ is a viscosity subsolution of (1.6) in $Q_r(x_0, t_0)$ if the following holds: if $(x, t) \in Q_r(x_0, t_0)$ and $\varphi \in C^2(Q_r(x_0, t_0))$ are such that $\varphi \ge u$ and $\varphi(x, t) = u(x, t)$ then

$$P_{\lambda,\Lambda}^+(D^2\varphi(x,t)) - \varphi_t(x,t) \ge -\phi(|D\varphi(x,t)|).$$

Note that the inequalities (1.5) and (1.6) are not homogeneous. We thus need the following rescaling lemma from [1, Lemma 2.4] and [9, Lemma 4.4].

Lemma 2.2 Let $u \in C(Q_r(x_0, t_0))$ be a viscosity supersolution of (1.5) (subsolution of (1.6)) in $Q_r(x_0, t_0)$. There exists a universal constant $L_2 \ge \Lambda_0$ such that if $A \in (0, \infty)$ then for every $r \le r_A$, where

$$r_A = \frac{A}{L_2(\phi(A) + A)} = \frac{1}{L_2(\eta(A) + 1)}$$
(2.2)

the rescaled function

$$u_r(x,t) := \frac{u(rx, r^2t)}{A},$$

is a supersolution of (1.5) (subsolution of (1.6)) in its domain.

3 Proofs of the main results

In this section we give the proofs of the four theorems.

Proof of Theorem 1.1 Without loss of generality we assume that n = 3. The case n > 3 then follows by adding dummy variables. In the following we write $(x, z) \in \mathbb{R}^2 \times \mathbb{R}$ for a point in \mathbb{R}^3 .

Fix a small $\varepsilon_0 > 0$ and choose $q = \frac{4}{\varepsilon_0}$. For $k \ge k_0(\varepsilon_0)$ denote $r = k^{-\varepsilon_0}$. We first define the functions u_k in $(\mathbb{R}^2 \setminus B_r) \times \mathbb{R}$ as

$$u_k(x,z) = \frac{1}{k}|x|^{-q}$$
 for $(x,z) \in (\mathbb{R}^2 \setminus B_r) \times \mathbb{R}$.

By direct calculation we have

$$\partial_{x_i x_j}^2 u_k(x, z) = \frac{q(q+2)}{k} |x|^{-q-4} x_i x_j - \frac{q}{k} |x|^{-q-2} \delta_{ij}.$$

The matrix $D^2 u_k(x, z)$ has eigenvalues $-\frac{q}{k}|x|^{-q-2}$, $\frac{q(q+1)}{k}|x|^{-q-2}$ and zero. Therefore, by choosing $\lambda = 1$ and $\Lambda = q + 1$, we have

$$P_{\lambda,\Lambda}^{-}(D^{2}u_{k}) = -(q+1)\frac{q}{k}|x|^{-q-2} + \frac{q(q+1)}{k}|x|^{-q-2} = 0.$$

We proceed by defining u_k in $B_r \times \mathbb{R}$ as

$$u_k(x, z) = \mathcal{A} - \mathcal{B}|x|^2 + \mathcal{C}|x|^4 + \mathcal{D}(z+2)\rho(|x|),$$

where $\rho \in C_c^{\infty}([0, r))$ is a cut-off function with $\rho \equiv 1$ in [0, r/2], $\rho' \leq 0$ in [0, r) and $|\rho'| \leq \frac{4}{r}$. Here, the coefficients $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are chosen so that u_k is a twice differentiable function. By a direct calculations this leads to

$$\mathcal{A} = \frac{8+6q+q^2}{8k}r^{-q}, \quad \mathcal{B} = \frac{q}{4k}(q+4)r^{-q-2}, \text{ and } C = \frac{q}{8k}(q+2)r^{-q-4}.$$
(3.1)

We then choose

$$\mathcal{D} = \frac{r^{-q}}{k}.$$
(3.2)

This choice will be clear later in the argument. By direct calculation we immediately verify that in $B_r \times (-2, 2)$ it holds

$$|P_{\lambda,\Lambda}^{-}(D^{2}u_{k})| \le C(\varepsilon_{0})\frac{r^{-q-2}}{k}.$$
(3.3)

Notice that in $B_{r/2} \times (-2, 2)$, using $\rho \equiv 1$, we have $|Du_k| \geq |\partial_z u_k| = \mathcal{D}$. Hence, we find that $|Du_k|^{1+\varepsilon_0} \geq k^{\varepsilon_0} \frac{r^{-q-2}}{k}$. Here, we used $r = k^{-\varepsilon_0}$ and $q = \frac{4}{\varepsilon_0}$. Consequently, for k large enough, we conclude that by (3.3) in $B_{r/2} \times (-2, 2)$ it holds

$$|P_{\lambda,\Lambda}^{-}(D^{2}u_{k})| \leq |Du_{k}|^{1+\varepsilon_{0}},$$

when k is large enough.

We are left to find a lower bound for $|Du_k|$ in $(B_r \setminus B_{r/2}) \times (-2, 2)$. Trivially, we have

$$|Du_k|^2 \ge |D_x u_k|^2 = |x|^2 \left(-2\mathcal{B} + 4\mathcal{C}|x|^2 + \mathcal{D}(z+2)\frac{\rho'}{|x|}\right)^2.$$

By the choices of \mathcal{B} and \mathcal{C} in (3.1) it holds

$$-2\mathcal{B} + 4C|x|^{2} \le -2\mathcal{B} + 4Cr^{2} = -\frac{q}{k}r^{-q-2},$$

in $B_r \times (-2, 2)$. Moreover, since $\rho' \le 0$ and $\frac{r}{2} \le |x| \le r$ in $(B_r \setminus B_{r/2}) \times (-2, 2)$, we obtain

$$|Du_k|^2 \ge |x|^2 \left(-2\mathscr{B} + 4\mathscr{C}|x|^2\right)^2 \ge \frac{q^2}{4} \frac{r^{-2q-2}}{k^2}$$

in $(B_r \setminus B_{r/2}) \times (-2, 2)$. Finally, we recall that $q = \frac{4}{\varepsilon_0}$ and $r = k^{-\varepsilon_0}$ to conclude that

$$|Du_k|^{1+\varepsilon_0} \ge k^{2\varepsilon_0+\varepsilon_0^2} \frac{r^{-q-2}}{k}$$

Deringer

in $(B_r \setminus B_{r/2}) \times (-2, 2)$. Therefore, by (3.3) we have for k large enough

$$|P^{-}_{\lambda,\Lambda}(D^2u_k)| \le |Du_k|^{1+\varepsilon_0}$$

in $(B_r \setminus B_{r/2}) \times (-2, 2)$ as well. Thus, we have established that, in B_2 ,

$$|P^{-}_{\lambda,\Lambda}(D^2u_k)| \le |Du_k|^{1+\varepsilon_0}$$

Finally we notice that $\inf_{B_1} u_k = \frac{1}{k}$ and $\sup_{B_1} u_k > 1$. Thus

$$\int_{\inf_{B_1} u_k}^{\sup_{B_1} u_k} \frac{ds}{s^{1+\varepsilon_0}+s} \ge \frac{1}{2} \int_{1/k}^1 \frac{ds}{s} = \frac{1}{2} \log(k) \to \infty$$

as $k \to \infty$. This completes the proof.

Proof of Theorem 1.2 Without loss of generality we may assume that u > 0. (Otherwise, we define $u + \varepsilon$ and let $\varepsilon \to 0$.) Let $(x_0, t_0) \in Q_2$ with $Q_{2\rho_0} \cup A^+_{\rho_0}(x_0, t_0) \subset Q_2$ and fix $\rho_0 \le \alpha_0$, where

$$\alpha_0 = \frac{a_0}{L_2(\eta(\frac{1}{2C}) + 1)}.$$
(3.4)

Here $A_{\rho_0}^+(x_0, t_0)$ is defined in (1.10), $a_0 = \frac{1}{C(\eta(u(x_0, t_0))+1)}$ and $C, L_2 > 1$ are from (1.9) and from Lemma 2.2 respectively. Since *u* is continuous there exists $\rho > 0$ such that

$$u(x_0, t_0) < 2C \inf_{A^+_{a_0\rho}(x_0, t_0)} u(x, t),$$

where *C* and a_0 are as above. In order to shorten the notation we denote $A^+_{a_0\rho}(x_0, t_0)$ by $A^+_{a_0\rho}$. We choose the smallest ρ , denoted by ρ_s , such that $a_0\rho_s \leq \rho_0$ and

$$u(x_0, t_0) = 2C \inf_{\substack{A_{a_0, \rho_s}^+}} u(x, t).$$
(3.5)

If such a ρ_s does not exist or if $a_0\rho_s = \rho_0$, then the claim of the theorem is trivially true. Let us then assume that $a_0\rho_s < \rho_0$. From (3.5) we deduce that there exists a point (x_s, t_s) in the closure of $A_{a_0\rho_s}^+$ such that

$$u(x_0, t_0) = 2Cu(x_s, t_s).$$

We define

$$v(x,t) = 2Cu(\rho_s x + x_s, \rho_s^2 t + t_s).$$

Next we recall the definition of α_0 in (3.4). Then $a_0\rho_s \leq \rho_0 \leq \alpha_0$ implies $\rho_s \leq \frac{1}{L_2(\eta(\frac{1}{2C})+1)}$. Therefore Lemma 2.2 implies that v is a positive viscosity supersolution of (1.5) and viscosity subsolution of (1.6) in Q_2 with $v(0, 0) = u(x_0, t_0)$. Thus, we may apply (1.9) to find

$$\sup_{A_{a_0}^-} v(x,t) \le C v(0,0).$$

Recall that $A_{a_0}^- = \left\{ (x, t) : |x|_{\infty} \le \frac{a_0 c_n}{2}, -a_0^2 + \frac{(a_0 c_n)^2}{4} \le t \le -a_0^2 + \frac{(a_0 c_n)^2}{2} \right\}$. It is easy to see that $\left(\frac{x_0 - x_s}{\rho_s}, \frac{t_0 - t_s}{\rho_s^2} \right) \in A_{a_0}^-$. Thus, in particular, we have

$$2Cu(x_0, t_0) \le Cu(x_0, t_0),$$

which is a contradiction as $u(x_0, t_0) > 0$. Hence, we have $a_0 \rho_s = \rho_0$ and

$$u(x_0, t_0) \le 2C \inf_{A_{\rho_0}^+} u(x, t).$$

This completes the proof of the theorem.

Let us then prove Theorem 1.3. As usual we use the above Harnack estimate to deduce that the oscillation has an algebraic decay as we reduce the size of the cubes. The difficulty is that we need to match the decay estimate to the size of the associated cubes in order to apply the intrinsic Harnack inequality.

Proof of Theorem 1.3 Let *C* be from Theorem 1.2. We let $\delta = 1 - \frac{1}{4C}$, $\rho_0 = 1$ and $\omega_0 := \sup_{Q_1} u - \inf_{Q_1} u$. Note that since C > 1 then $\delta > \frac{1}{2}$. We also recall the notation $r_A := \frac{1}{L_2(\eta(A)+1)}$ for a number A > 0 from Lemma 2.2. For $k \ge 1$ we define

$$\rho_k = \left(\frac{r_\delta c_n}{4C\eta(4)}\right)^k r_{\omega_0/4}.$$

We claim that for every $k \ge 0$ it holds

$$\operatorname{osc}_{\mathcal{Q}_{\rho_k}} u \le \delta^k \omega_0, \tag{3.6}$$

where $\operatorname{osc}_{Q_{\rho}} u := \sup_{Q_{\rho}} u - \inf_{Q_{\rho}} u$.

We argue by contradiction. Note that (3.6) is trivially true for k = 0 and therefore the contradiction assumption implies that there is $k \ge 0$ such that

$$\operatorname{osc}_{Q_{\rho_k}} u \le \delta^k \omega_0 \tag{3.7}$$

and

$$\operatorname{osc}_{\mathcal{Q}_{\rho_{k+1}}} u > \delta^{k+1} \omega_0. \tag{3.8}$$

🖉 Springer

We denote $M_k = \sup_{Q_{\rho_k}} u$, $m_k = \inf_{Q_{\rho_k}} u$ and define also $t_k = \left(1 - \frac{c_n^2}{4}\right) \left(\frac{r_{\delta}\rho_k}{2C\eta(4)}\right)^2$. We claim that one of the following holds

(i) $M_k - u(0, -t_k) \ge \frac{\delta^k \omega_0}{4}$ (ii) $u(0, -t_k) - m_k \ge \frac{\delta^k \omega_0}{4}$

Indeed, if both are false then $\operatorname{osc}_{Q_{\rho_k}} u \leq \delta^k \omega_0/2$. Note that from (3.8), we have $\delta^{k+1}\omega_0 < \operatorname{osc}_{Q_{\rho_k}} u$. Hence, we obtain $\delta < 1/2$, which is a contradiction. Hence either (i) or (ii) holds.

Assume first that the condition (i) above holds. We define a function $v: Q_2 \to \mathbb{R}$ as

$$v(x,t) = \frac{M_k - u(r_\delta \rho_k x, (r_\delta \rho_k)^2 t - t_k)}{\delta^k \omega_0 / 4}.$$
(3.9)

Let us denote $b_0 := \frac{1}{2C\eta(4)}$. We note that by the choices of ρ_k , t_k and b_0 it holds

$$A_{b_0 r_\delta \rho_k}^+(0, -t_k) = Q_{\rho_{k+1}} \subset Q_2,$$
(3.10)

where A_{ρ}^{+} is defined in (1.10). Let us show that v is a supersolution of (1.5) and subsolution of (1.6) in Q_2 .

Indeed, by the definition of ρ_k it holds

$$r_{\delta}\rho_{k} \leq r_{\delta}^{k+1}r_{\omega_{0}/4} = \frac{1}{L_{2}^{k+2}}\frac{1}{(\eta(\delta)+1)^{k+1}}\frac{1}{\eta(\omega_{0}/4)+1}$$

On the other hand, by the condition (P3) in (1.4) on η it holds

$$\eta(\delta^k \omega_0/4) + 1 \le \Lambda_0^{k+1} (\eta(\delta) + 1)^k (\eta(\omega_0/4) + 1).$$

Therefore, since $\Lambda_0 \leq L_2$ it holds

$$r_{\delta}\rho_k \le \frac{1}{L_2(\eta(\delta^k\omega_0/4)+1)} = r_{\delta^k\omega_0/4}$$

and by Lemma 2.2, v defined in (3.9) is a supersolution of (1.5) and subsolution of (1.6).

Since $M_k = \sup_{Q_{\rho_k}} u$, we conclude that v is non-negative in Q_2 , by possibly enlarging the constant L_2 if necessary. Moreover, the condition (i) and (3.7) imply $1 \le v(0,0) \le 4$. Therefore since η is increasing in $[1,\infty)$ and $\eta \ge 1$, we have $b_0 = \frac{1}{2C\eta(4)} \le \alpha_0 = \frac{1}{C(\eta(v(0,0))+1)}$. Therefore by Theorem 1.2, we have

$$\frac{1}{C}v(0,0) \le \inf_{A_{b_0}^+} v(x,t).$$

Using $v(0, 0) \ge 1$, we have

$$\frac{\delta^k \omega_0}{4C} \leq M_k - \sup_{A_{b_0}^+} u(r_\delta \rho_k x, (r_\delta \rho_k)^2 t - t_k).$$

Recalling (3.10) the above yields

 $\sup_{\mathcal{Q}_{\rho_{k+1}}} u - m_k \le M_k - m_k - \frac{\delta^k \omega_0}{4C} = \operatorname{osc}_{\mathcal{Q}_{\rho_k}} u - \frac{\delta^k \omega_0}{4C} \le \delta^k \omega_0 - \frac{\delta^k \omega_0}{4C} = \delta^{k+1} \omega_0.$

Hence, we obtain

$$\operatorname{osc}_{Q_{\rho_{k+1}}} u \leq \delta^{k+1} \omega_0$$

which contradicts (3.8).

If the condition (ii) holds, we define

$$v(x,t) = \frac{u(r_{\delta}\rho_k x, (r_{\delta}\rho_k)^2 t - t_k) - m_k}{\delta^k \omega_0/4}.$$

By repeating the above argument leads again to a contradiction. (We leave the details for the reader). Therefore we finally have the estimate (3.6).

Let us now prove the Hölder continuity. Given $r \leq 1$, choose k such that

$$\left(\frac{r_{\delta}c_n}{4C\eta(4)}\right)^{k+1} \le r \le \left(\frac{r_{\delta}c_n}{4C\eta(4)}\right)^k$$

and use (3.6) to conclude

$$\operatorname{osc}_{Q_r} u\left(r_{\omega_0/4}x, r_{\omega_0/4}^2t\right) \leq \frac{1}{\delta}r^{\alpha}\omega_0,$$

where $\alpha = \min\left\{\frac{1}{2}, \frac{\log \delta}{\log\left(\frac{r_{\delta}c_n}{4C\eta(4)}\right)}\right\}$ is a universal constant. Therefore, we have for all $(x, t) \in Q_1$

$$\left| u\left(r_{\omega_0/4}x, r_{\omega_0/4}^2 t \right) - u(0,0) \right| \le \frac{1}{\delta} (|x| + |t|^{1/2})^{\alpha} \omega_0$$

Thus, for $(x, t) \in Q_{r_{\omega_0/4}}$ we have

$$|u(x,t) - u(0,0)| \le \frac{1}{r_{\omega_0/4}^{\alpha}\delta} (|x| + |t|^{1/2})^{\alpha} \omega_0 \le \tilde{C}\phi(\omega_0)(|x| + |t|^{1/2})^{\alpha},$$

Deringer

where in the second inequality we have used the fact that $\phi(s) = \eta(s)s$ for $\eta \ge 1$ and $\alpha < 1$. Here $\tilde{C} > 1$ is a universal constant. On the other hand, it is easy to see that for all $(x, t) \in Q_1 \setminus Q_{r_{\alpha\alpha}/4}$ it holds

$$|u(x,t) - u(0,0)| \le \omega_0 \frac{r_{\omega_0/4}^{\alpha}}{r_{\omega_0/4}^{\alpha}} \le \tilde{C}\phi(\omega_0)(|x| + |t|^{1/2})^{\alpha}.$$

Therefore, we conclude that for all $(x, t) \in Q_1$, it holds

$$|u(x,t) - u(0,0)| \le \tilde{C}\phi(||u||_{\infty})(|x| + |t|^{1/2})^{\alpha}.$$

This implies the claim by using the fact that the equations (1.5) and (1.6) are translation invariant together with a standard covering argument.

Proof of Theorem 1.4 Without loss of generality we may assume that u > 0. (Otherwise, we define $u + \varepsilon$ and let $\varepsilon \to 0$.)

We define a sequence of radii $r_i > 0$ and time levels $t_i < 0$ as follows. Set first $M_0 = u(0, 0), \rho_1 = \frac{1}{C(\eta(M_0)+1)}$ and choose the radius $r_1 = \frac{\rho_1 c_n}{2}$ and the time $t_1 = \left(-1 + \frac{c_n^2}{4}\right)\rho_1^2$. Here C > 1 and $c_n \le 1$ are from (1.9). We proceed by defining r_i iteratively such that if r_i is defined, we let

$$\tilde{M}_i = \sup_{x \in B_{r_i}} u(x, t_i) \text{ and } M_i = \max\{M_{i-1}, \tilde{M}_i\},$$
(3.11)

set $\rho_{i+1} = \frac{1}{C(\eta(M_i)+1)}$ and define

$$r_{i+1} = r_i + \frac{\rho_{i+1}c_n}{2}$$
 and $t_{i+1} = t_i + \left(-1 + \frac{c_n^2}{4}\right)\rho_{i+1}^2$.

We first observe that there exists $K \in \mathbb{N}$ such that $r_K \ge 1$. Indeed, if $r_i < 1$ for all $i \in \mathbb{N}$ then the definition of r_i yields

$$\frac{c_n}{2C}\sum_{i=0}^{\infty}\frac{1}{\eta(M_i)+1} \le 1$$

and $t_i \ge -1$ for all *i*. But then necessarily $\rho_i \to 0$ as $i \to \infty$, which means that $\eta(M_i) \to \infty$ as $i \to \infty$. This in turn implies $M_i \to \infty$, which is a contradiction since *u* is continuous. We define *K* to be the first index for which $r_K \ge 1$ and note that $r_K \le 2$.

We proceed by claiming that for all i = 0, 1, ..., K - 1 it holds

$$M_{i+1} \le CM_i. \tag{3.12}$$

If $M_{i+1} = M_i$ then (3.12) is trivially true. Let us then assume that $M_{i+1} > M_i$. We choose $x_{i+1} \in \overline{B}_{r_{i+1}}$ such that $M_{i+1} = u(x_{i+1}, t_{i+1})$ and let y_i be the closest point to

 x_{i+1} in \overline{B}_{r_i} . Then it holds $u(y_i, t_i) \leq M_i$. We define

$$f(r) := u\left(x_{i+1} + \frac{r}{\rho_{i+1}}(y_i - x_{i+1}), t_{i+1} + \left(1 - \frac{c_n^2}{4}\right)r^2\right)$$

and notice that $f(0) = u(x_{i+1}, t_{i+1}) = M_{i+1} > M_i \ge u(y_i, t_i) = f(\rho_{i+1})$. By continuity of *u* there exists $0 < \rho_0 \le \rho_{i+1}$ such that $f(\rho_0) = M_i$. We set

$$(x_0, t_0) = \left(x_{i+1} + \frac{\rho_0}{\rho_{i+1}}(y_i - x_{i+1}), t_{i+1} - \left(-1 + \frac{c_n^2}{4}\right)\rho_0^2\right)$$

and apply (1.9) to have

$$\sup_{A_{\rho_0}^-(x_0,t_0)} u(x,t) \le CM_i.$$

Notice that $(x_{i+1}, t_{i+1}) \in A^-_{\rho_0}(x_0, t_0)$ and therefore it holds $u(x_{i+1}, t_{i+1}) = M_{i+1} \le CM_i$, which proves (3.12).

We use (3.12) to estimate

$$\int_{M_0}^{M_K} \frac{ds}{\phi(s) + s} \le \sum_{i=0}^{K-1} \int_{M_i}^{M_{i+1}} \frac{ds}{\phi(s) + s} \le \sum_{i=0}^{K-1} \int_{M_i}^{CM_i} \frac{ds}{\phi(s) + s}$$

Since ϕ is increasing and of the form $\phi(s) = \eta(s)s$ we obtain

$$\begin{split} \sum_{i=0}^{K-1} \int_{M_i}^{CM_i} \frac{ds}{\phi(s)+s} &\leq \sum_{i=0}^{K-1} \int_{M_i}^{CM_i} \frac{ds}{\phi(M_i)+M_i} \\ &= \sum_{i=0}^{K-1} \frac{(C-1)M_i}{\phi(M_i)+M_i} = (C-1) \sum_{i=0}^{K-1} \frac{1}{\eta(M_i)+1} \\ &= C(C-1) \sum_{i=0}^{K-1} \rho_{i+1} \leq \frac{2C^2}{c_n} r_K. \end{split}$$

Since $r_K \leq 2$ and $\phi(s) \geq s$, we conclude

$$\int_{M_0}^{M_K} \frac{ds}{\phi(s)} \le \tilde{C},\tag{3.13}$$

for a universal constant \tilde{C} .

Deringer

Note that in (3.13) we have $M_0 = u(0, 0)$ and $M_K \ge \sup_{x \in B_1} u(x, t_K)$. We let $t_2 = -t_K$ and need yet to estimate $|t_K|$. By the definition of t_K and ρ_i it holds

$$|t_K| = \left(1 - \frac{c_n^2}{4}\right) \sum_{i=1}^K \rho_i^2 \le \sum_{i=1}^K \rho_i^2.$$
(3.14)

Then by increasing *C* if necessary such that $C^{-1} \leq \frac{c_n}{4}$, we have $\rho_i = \frac{1}{C(\eta(M_i)+1)} \leq \frac{c_n}{4}$. Then since $r_K \leq 2$ we have

$$|t_K| \le \sum_{i=1}^K \rho_i^2 \le \frac{c_n}{4} \sum_{i=1}^K \rho_i = \frac{r_K}{2} \le 1.$$

On the other hand, trivially it holds

$$|t_K| \ge |t_1| \ge \frac{1}{2C^2(\eta(u(0,0))+1)^2}.$$

Consequently, we obtain

$$\frac{1}{\tilde{C}(\eta(u(0,0))+1)^2} \le t_2 \le 1,$$

for a universal constant \tilde{C} .

Similarly, we prove

$$\int_{\inf_{B_1} u(\cdot,t_1)}^{u(0,0)} \frac{ds}{\phi(s)} \le \tilde{C}$$

for a universal constant \tilde{C} and a time level $t_1 > 0$ with

$$\frac{1}{\tilde{C}(\eta(u(0,0))+1)^2} \le t_1 \le 1.$$

This completes the proof.

Corollary 1.5 follows from Theorem 1.4 but we include the argument for the reader's convenience.

Proof of Corollary 1.5 Let $u \in C(Q_4(0, 1))$ be as in the statement and define $u_{\varepsilon} = u + \varepsilon$. We adopt the notation from the proof of Theorem 1.4 and denote $M_i(\varepsilon)$ as in (3.11) for u_{ε} and $\rho_{i+1} = \frac{1}{C(\eta(M_i(\varepsilon))+1)}$. Since $u_{\varepsilon}(0, 0) = \varepsilon$ we have by Theorem 1.4 that

$$\int_{\varepsilon}^{M_{K}(\varepsilon)} \frac{ds}{\phi(s)} \leq C$$

Deringer

for $M_K(\varepsilon) \ge \sup_{x \in B_1} u_{\varepsilon}(x, t_K(\varepsilon))$. Here $t_K(\varepsilon) = -t_2$. But since ϕ satisfies (1.8) then $M_K(\varepsilon) \to 0$ as $\varepsilon \to 0$. We need yet to prove that we may choose $t_K(\varepsilon)$ such that $t_K(\varepsilon) \to 0$ as $\varepsilon \to 0$.

We divide the proof in two cases. Assume first

$$\lim_{s\to 0}\eta(s)<\infty.$$

Then there is \tilde{C} such that $1 \leq \eta(s) \leq \tilde{C}$ for all $s \in (0, 1]$. Therefore the equations (1.5) and (1.6) reduce to the linear case and we have the classical Harnack inequality by [8, 15] and the claim follows.

Assume then

$$\lim_{s \to 0} \eta(s) = \infty.$$

The definition (3.11) of $M_i(\varepsilon)$ implies $M_i(\varepsilon) \le M_K(\varepsilon)$. Since $M_K(\varepsilon) \to 0$ as $\varepsilon \to 0$, then for every $\delta > 0$ it holds $\rho_{i+1} = \frac{1}{C(\eta(M_i(\varepsilon))+1)} < \delta$ for all $i \le K(\varepsilon) - 1$ when ε is small. By the definition of $r_{K(\varepsilon)}$ it holds

$$\sum_{i=1}^{K(\varepsilon)} \rho_i = \frac{2}{c_n} r_{K(\varepsilon)} \le \frac{4}{c_n}$$

Therefore by (3.14) we have

$$|t_K(\varepsilon)| \le \sum_{i=1}^{K(\varepsilon)} \rho_i^2 \le \delta \sum_{i=1}^{K(\varepsilon)} \rho_i \le \frac{4\delta}{c_n}.$$

The claim follows by letting first $\varepsilon \to 0$ and then $\delta \to 0$.

Acknowledgements The authors were supported by the Academy of Finland grant 314227. The authors have no non-financial competing interests to declare that are relevant to the concent of this article.

Funding Open Access funding provided by University of Jyväskylä (JYU).

Data availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Arya, V.: An Intrinsic Harnack inequality for some non-homogeneous parabolic equations in nondivergence form. Calc. Var. Partial Differ. Equ. 61, 30–31 (2022)
- Avelin, B., Julin, V.: A Carleson type inequality for fully nonlinear elliptic equations with non-Lipschitz drift term. J. Funct. Anal. 272, 3176–3215 (2017)
- Bingham, N.H., Goldie, C.M., Teugels, J.L.: Regular Variation. Encyclopedia of Mathematics and Its Applications, vol. 27. Cambridge University Press, Cambridge (1989)
- Caffarelli, L.: Interior a priori estimates for solutions of fully nonlinear equations. Ann. Math. (2) 130, 189–213 (1989)
- Caffarelli, L., Cabre, X.: Fully Nonlinear Elliptic Equations. American Mathematical Society Colloquium Publications, vol. 43. American Mathematical Society, Providence (1995)
- DiBenedetto, E., Gianazza, U., Vespri, V.: Harnack estimates for quasi-linear degenerate parabolic differential equations. Acta Math. 200, 181–209 (2008)
- DiBenedetto, E., Gianazza, U., Vespri, V.: Harnack's Inequality for Degenerate and Singular Parabolic Equations. Springer Monographs in Mathematics, p. xiv+278. Springer, New York (2012). (ISBN: 978-1-4614-1583-1)
- Imbert, C., Silvestre, L.: An Introduction to Fully Nonlinear Parabolic Equations. In: An Introduction to the K\u00e4hler-Ricci Flow. Lecture Notes in Mathematics, vol 2086. pp. 7–88, Springer, Cham (2013)
- 9. Julin, V.: Generalized Harnack inequality for Nonhomogeneous elliptic equations. Arch. Ration. Mech. Anal. 216, 673–702 (2015)
- Krylov, N., Safonov, M.: A proöperty of the solutions of parabolic equations with measurable coefficients. Izv. Akad. Nauk SSSR Ser. Mat. 44, 161–175 (1980). ((Russian))
- Kuusi, T.: Harnack estimates for weak supersolutions to nonlinear degenerate parabolic equations. Ann. Sc. Norm. Super. Pisa Cl. Sci. 7(5), 673–716 (2008)
- Lundström, N.: Growth of subsolutions to fully nonlinear equations in halfspaces. J. Differ. Equ. 320, 143–173 (2022)
- Lundström, N., Olofsson, M., Toivanen, O.: Strong maximum principle and boundary estimates for nonhomogeneous elliptic equations. Potential Anal. 60, 425–443 (2024)
- 14. Trudinger, N.S.: Comparison principles and pointwise estimates for viscosity solutions of nonlinear elliptic equations. Rev. Mat. Iberoam. **4**, 453–468 (1988)
- Wang, L.: On the regularity theory of fully nonlinear parabolic equations. I. Commun. Pure Appl. Math. 45(1), 27–76 (1992)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.