

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

- Author(s): Wasternack, Janos; Schröder, Hendrik V.; Witte, J. Felix; Ilisson, Mihkel; Hupatz, Henrik; Hille, Julian F.; Gaedke, Marius; Valkonen, Arto M.; Sobottka, Sebastian; Krappe, Alexander; Schubert, Mario; Paulus, Beate; Rissanen, Kari; Sarkar, Biprajit; Eigler, Siegfried; Resch-Genger, Ute; Schalley, Christoph A.
- **Title:** Switchable protection and exposure of a sensitive squaraine dye within a redox active rotaxane

Year: 2024

Version: Published version

Copyright: © The Author(s) 2024

Rights: CC BY 4.0

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Wasternack, J., Schröder, H. V., Witte, J. F., Ilisson, M., Hupatz, H., Hille, J. F., Gaedke, M., Valkonen, A. M., Sobottka, S., Krappe, A., Schubert, M., Paulus, B., Rissanen, K., Sarkar, B., Eigler, S., Resch-Genger, U., & Schalley, C. A. (2024). Switchable protection and exposure of a sensitive squaraine dye within a redox active rotaxane. Communications Chemistry, 7, Article 229. https://doi.org/10.1038/s42004-024-01312-1

0

https://doi.org/10.1038/s42004-024-01312-1

Switchable protection and exposure of a sensitive squaraine dye within a redox active rotaxane

Check for updates

Janos Wasternack ¹, Hendrik V. Schröder ¹, J. Felix Witte ², Mihkel Ilisson ¹, Henrik Hupatz ¹, Julian F. Hille¹, Marius Gaedke¹, Arto M. Valkonen ³, Sebastian Sobottka⁴, Alexander Krappe ⁵, Mario Schubert ⁶, Beate Paulus², Kari Rissanen ³, Biprajit Sarkar^{4,7}, Siegfried Eigler ⁵, Ute Resch-Genger ⁸ & Christoph A. Schalley ¹

In nature, molecular environments in proteins can sterically protect and stabilize reactive species such as organic radicals through non-covalent interactions. Here, we report a near-infrared fluorescent rotaxane in which the stabilization of a chemically labile squaraine fluorophore by the coordination of a tetralactam macrocycle can be controlled chemically and electrochemically. The rotaxane can be switched between two co-conformations in which the wheel either stabilizes or exposes the fluorophore. Coordination by the wheel affects the squaraine's stability across four redox states and renders the radical anion significantly more stable—by a factor of 6.7—than without protection by a mechanically bonded wheel. Furthermore, the fluorescence properties can be tuned by the redox reactions in a stepwise manner. Mechanically interlocked molecules provide an excellent scaffold to stabilize and selectively expose reactive species in a co-conformational switching process controlled by external stimuli.

From RNA enclosed in capsid shells^{1,2} to reactive molecular centers and active sites³ buried deep within shielding proteins, confinement in protective environments is a fundamental strategy in nature to mitigate undesired side reactions of labile species^{4,5}. In specific molecular environments within proteins, for instance, reactive molecules such as organic chromophores, radicals, or ions are not only sterically shielded, but are also stabilized through non-covalent interactions. Prime examples are β -barrels hosting labile chromophores in fluorescent proteins⁶ or organic radical ions stabilized through hydrogen bonding in flavoenzymes⁷ and photosynthetic reaction centers⁸.

It remains a formidable challenge for synthetic chemists to stabilize reactive species (e.g., organic radicals) without interfering with the molecules' functions to enable applications in biomedical imaging⁹, sensing¹⁰, or proximity labeling¹¹. In contrast to classical covalent protecting group strategies, encapsulation by a receptor offers a supramolecular approach for the stabilization of reactive guests by non-covalent interactions (Fig. 1a). A host molecule, such as a macrocycle, confines a reactive guest and acts as a dynamically bound 'supramolecular protecting group'. The protection strength is determined by energetic stabilization upon binding inside the host and steric shielding. The abundance of free, unprotected guest is significantly decreased upon binding inside a confined space¹²⁻¹⁶.

An alternative approach to increase stability involves the incorporation of reactive sites into mechanically interlocked molecules (MIMs) as seen for synthetic¹⁷⁻²¹ and natural²²⁻²⁶ examples like the lasso peptide microcin J25²⁷ (Fig. 1b). An advantage compared to a non-interlocked complex is that the mechanical bond prevents dissociation into subcomponents and thus conformationally locks the stabilizing structure. For example, some mechanically interlocked peptides display mechanical bonds with a low degree of conformational freedom, equipping them with an unusually high physicochemical stability compared to their non-interlocked counterparts^{22,27}. A synthetic example was reported by SMITH et al., who applied protection within MIMs for sensitive squaraine dyes. Squaraine (Sq) dyes are important near infrared (NIR) fluorescent dyes with broad application in bioimaging^{28–31}. NIR dyes offer low absorption interference by biological tissues, deeper tissue penetration, reduced scattering, and among them, Sq are of particular interest due to their highly intense NIR absorption

¹Institut für Chemie und Biochemie, Freie Universität Berlin, Arnimallee 20, 14195 Berlin, Germany. ²Institut für Chemie und Biochemie, Freie Universität Berlin, Arnimallee 22, 14195 Berlin, Germany. ³University of Jyvaskyla, Department of Chemistry, Survontie 9 B, 40014 Jyväskylä, Finland. ⁴Institut für Chemie und Biochemie, Freie Universität Berlin, Freie Universität Berlin, Freie Universität Berlin, Freie Universität Berlin, Germany. ⁶Institut für Chemie und Biochemie, Freie Universität Berlin, Germany. ⁶Institut für Chemie und Biochemie, Freie Universität Berlin, Germany. ⁶Institut für Chemie und Biochemie, Freie Universität Berlin, Germany. ⁶Institut für Chemie und Biochemie, Freie Universität Berlin, Altensteinstraße 23A, 14195 Berlin, Germany. ⁶Institut für Chemie und Biochemie, Freie Universität Berlin, Takustraße 3, 14195 Berlin, Germany. ⁷Institut für Anorganische Chemie, Universität Stuttgart, Pfaffenwaldring 55, 70569 Stuttgart, Germany. ⁸Bundesanstalt für Materialforschung und -prüfung (BAM), Biophotonics, Richard Willstätter Straße 11, 12489 Berlin, Germany. ^Ke-mail: c.schalley@fu-berlin.de



Fig. 1 | **Different protecting strategies. a** Encapsulation of a labile guest by a host molecule results in an inclusion complex in which the guest is sterically shielded from unwanted side reactions with a compound X. **b** Lasso peptide microcin J25²⁷ as

example of drastically increased physicochemical stability by mechanical bonding. c Electrochemical switching of protection by a macrocycle encircling a labile squaraine dye in a [2]rotaxane.

and emission bands³¹⁻³³. However, Sq suffer from their vulnerability to nucleophilic attack at the dye's electrophilic C_4O_2 core, resulting in color fading²⁸. Their enclosure by HUNTER-VÖGTLE type tetralactam macrocycles encircling the Sq core through NH ··· O hydrogen bonds is a useful means of protection against nucleophilic attack²⁸.

A particularly interesting aspect of MIMs is that co-conformational flexibility can be intentionally designed into a molecular structure, enabling controlled large-amplitude motions of their subcomponents. This property has made MIMs preferred scaffolds for the construction of artificial molecular switches and machines³⁴⁻³⁸ mimicking dynamic conformational changes found in biological nanomachines. In SMITH's Sq rotaxanes, addition of chloride salts leads to the displacement of the macrocycle from the Sq station, thus switching off the protection²⁸⁻³⁰.

To make such switchable protection multi-stimuli responsive is the aim of this work. Besides switching by chloride addition and removal, we demonstrate that electrochemical switching can also be realized in this Sq rotaxane. We show that the energetic stabilization and steric shielding of the reactive Sq fluorophore within a conformationally dynamic MIM can be altered by electrochemical and chemical stimuli. These external stimuli either increase the stabilization by the wheel or induce a co-conformational change, resulting in the controllable exposure of this reactive site (Fig. 1c). Furthermore, the rotaxane exists in four oxidation states, which can be utilized for tuning of the optoelectronic properties of those dyes.

Results and discussion

Preparation and structure of Rot

[2]Rotaxane Rot comprises a HUNTER-VÖGTLE type tetralactam macrocycle TLM and a threaded and mechanically interlocked bis(aminothienyl)

squaraine axle. This macrocycle was chosen because the pyridyl group facilitates binding and preorganisation during the synthesis of Rot and also improves solubility of the macrocycle³⁹. Wheel TLM and axle Ax form a threaded complex, pseudo[2]rotaxane psRot in dichloromethane with an association constant of $K_a = 520 \pm 160 \text{ M}^{-1}$ at 293 K as determined by ¹H NMR titration (SI section S2 Fig. S17). Slow evaporation of dichloromethane solutions of Ax and of psRot yielded single crystals suitable for XRD analysis. The 1:1 complex stoichiometry was confirmed by solid-state XRD structures of the psRot and Ax (Figs. 2 and 4g). The structure of psRot is similar to reported Sq tetralactam rotaxanes (SI section S4 Fig. S25)^{28,30,40}. The Sq bound by four NH \cdots O hydrogen bonds (2.21 Å) and the bond lengths within the neutral Sq within psRot are very close to the ones in free Ax. Rot was obtained in 32% yield from the solution of psRot in an end-capping approach via copper-catalyzed azide-alkyne HUISGEN cycloaddition of a sterically demanding trityl stopper (Figs. 3b and 8). The most stable coconformation ^{Sq}Rot⁰ involves the wheel encircling the squaraine core by forming NH ... O hydrogen bonds.

Chloride switching of neutral Rot

Addition of chloride salts to a dichloromethane solution of **Rot** results in a significantly increased fluorescence intensity, likely caused by shuttling of the wheel to the vicinity of the triazol group, thereby exposing the squaraine core (Figs. S21–S23). This shuttling behavior has been observed in similar tetralactam rotaxanes^{28,40}. The fluorescence quantum yields as well as the maxima of absorption and emission of the fluorescent states of **Ax** and **Rot** were measured (SI section S3, Table S1). **Ax**⁰ displays fluorescence with an emission maximum of $\lambda_{max}(\mathbf{Ax}^0) = 678$ nm and an absolutely measured fluorescence quantum yield of $\Phi(\mathbf{Ax}^0) = 6.7\%$ and **Rot**⁰ displays



Fig. 2 | Single Crystal XRD structure of psRot with selection of bond lengths (thermal displacement parameters at 50% probability level). The perspective is indicated by the point of view icon next to the schematic depiction. The macrocycle is positionally disordered in the crystal and the pyridine dicarboxylic amide and isophthalic diamide can replace each other in their positions.

fluorescence with an emission maximum of $\lambda_{max}(\mathbf{Rot}^0) = 683$ nm and a quantum yield of $\Phi(\mathbf{Rot}^0) = 4.0\%$. The lower quantum yield of \mathbf{Rot}^0 could be attributed to lower absorption of the shielded Sq or partial absorption of the emitted light by the wheel. Addition of 50 equiv. Bu₄NCl to a solution of \mathbf{Ax}^0 leads to a decreased quantum yield of $\Phi(\mathbf{Ax}^0 \supset Cl^-) = 5.9\%$, while under the same conditions the quantum yield of \mathbf{Rot}^0 increases to $\Phi(\mathbf{Rot}^0 \supset Cl^-) = 4.2\%$. The increased fluorescence intensity in Triaz $\mathbf{Rot}^0 \supset Cl^-$ thus likely results from increased absorbance and reduced quenching by the wheel of the deshielded squaraine. The observed properties and the chloride induced translocation of the wheel are in good alignment with similar systems^{28,40}.

The chloride affinity of **Rot**⁰ was determined by ¹H NMR $(K_a = 84 \pm 12 \text{ M}^{-1})$ and fluorescence titrations $(K_a = 90 \pm 20 \text{ M}^{-1})$ using *n*-Bu₄NCl (Figs. S19–S22). Figure 3 shows the ¹H NMR spectra of **Rot** before and after addition of 59 equiv. *n*-Bu₄NCl. Binding of chloride induces the co-conformational change to ^{Triaz}**Rot**⁰ \supset Cl⁻ in which the wheel **TLM** binds a chloride ion and the triazole station, as indicated by characteristic shifts of the NH signals ε : $\Delta\delta(\varepsilon) = 0.2$ ppm and σ : $\Delta\delta(\sigma) = -1.7$ ppm, and the aromatic signals χ : $\Delta\delta(\chi) = 0.3$ ppm and σ : $\Delta\delta(\sigma) = 0.2$ ppm. A binding mode of chloride in similar tetralactam - triazol rotaxanes through CH ---Cl⁻--HN hydrogen bonds between the triazole's CH group and the NH groups of the tetralactam has been reported^{41,42}.

Quantum chemical investigation of ^{Triaz}**Rot**⁰ \supset Cl⁻ by MD simulations using CREST, indicate hydrogen bonding between the amide NH protons of the isophthalamide and chloride, while the pyridyl bis amide participates in hydrogen bonding towards the triazol group. Weak hydrogen bonds between the triazol proton m and chloride are indicated by the comparatively low shift of m: $\Delta\delta(m) = 0.1$ ppm. This is in line with the calculated binding motif in which the triazol is tilted away from the bound chloride anion (SI section S9 Fig. S42). The presence of only one set of broadened signals suggests that translational exchange of the wheel between the two triazoles is faster than the NMR time scale. Precipitation of the chloride by addition of Na[B(Ph-3,5-(CF₃)₂)₄] leads to back shuttling of the wheel onto the Sq. Reversibility of the shuttling was again confirmed by fluorescence spectroscopy (Fig. S23).

Redox switching of Rot

We hypothesized that a similar switching between co-conformations might be realized by electrochemistry. LEIGH and co-workers reported an electrochemically switchable hydrogen-bonded molecular shuttle⁴³. We recently developed a series of tetrathiafulvalene rotaxanes, which are electrochemically switchable between different co-conformations^{44–53}. (Quasi) reversible redox reactions have been reported for some Sq derivatives⁵⁴.

Cyclic voltammetry in 1,2-dichloroethane (Fig. 4b and Table S2) reveals that Ax^0 undergoes an irreversible one-electron reduction to the radical anion $Ax^{\bullet-}$ ($E_{pc}^{-1} = -0.99$ V against the decamethylferrocene/ decamethylferrocenium couple) as well as two reversible one-electron oxidations, to the radical cation $Ax^{\bullet+}$ ($E_{1/2}^1 = 0.45$ V) and dication Ax^{2+} ($E_{1/2}^2 = 0.95$ V). These three redox reactions are most likely localized on the thiophene-squaraine π -system⁵⁴.

Scan rate-dependent plots of the anodic peak potential (SI section S5 Figs. S26 and S29) indicate that the reduction of the axle $(Ax^0 \rightarrow Ax^{\bullet-})$ is followed by an irreversible chemical reaction step in a so-called EC_i mechanism. For Sq^{•-}, Quantum chemical calculations at various levels of density functional theory (DFT) (SI section S9) indicate elongated O-C1bonds and an increase of electron density on the oxygen atoms (Fig. 4a and h, Table S4). In $Rot^{\bullet-}$, both effects result in stabilization of the NH \cdots O hydrogen bonds between $Sq^{\bullet-}$ and wheel by 42 kJ/mol (Table S7). In addition to the electronic stabilization of the wheel- $Sq^{\bullet-}$ interaction, the steric shielding of Sq^{•-} by TLM contributes to an observed increase in lifetime. In contrast to the (quasi)irreversible reduction of Ax^0 , the reduction of Rot⁰ is reversible on the CV timescale (scan rate of 100 mV/s), further supporting the significant stabilization of the radical anion Sq^{•-} by rotaxanation. The lifetime of the radical anion Rot^{•-} was investigated by CVbased digital simulations (SI section S5 Fig. S29). Fitting the experimental voltammograms at different scan rates according to an EC_i mechanism yields an average radical lifetime of 1.7 s ($Ax^{\bullet-}$) and 10.8 s ($Rot^{\bullet-}$). However, the anodic peak (Epa) of Rot was overlayed by incipient solvent/ electrolyte decomposition. Thus, the life-time ratio $\tau_{Rot}/\tau_{Ax} > 6.7 \pm 2.8$ is considered to be a lower boundary of the stabilization effect caused by rotaxanation. Organic radicals are typically highly reactive species with short lifetimes⁵⁵. They were categorised by INGOLD according to their lifetimes as transient radicals (lifetimes < 1.44 ms) and persistent radicals $(lifetimes > 1.44 ms)^{5657}$. For example the benzyl radical is transient with a lifetime below 1 ms^{57,58}. The drastic impact of thermodynamic and steric factors on radical stability are illustrated by the existence of bench stable radicals like TEMPO. Supramolecular encapsulation is a common means of stabilisation of transient and persistent radicals4,43,55,59-64. A prominent example being the drastic stabilisation of a persistent tetrazine radical anion with a lifetime of 2 h through encapsulation by two cyanostar macrocycles leading to a complex with a 360-fold lifetime of 30 d⁶⁴. Our case thus represents the stabilisation of a more reactive persistent radical.

In contrast to the strengthening of the wheel–axle interaction in **Rot** by the reduction, oxidation reduces the electron density of Sq as indicated by calculated electrostatic potential surface maps (ESPs), resulting in a weakened wheel–axle interaction in **Rot**^{\bullet +} and **Rot**²⁺ (Fig. 4a and d and S43, Tables S4 and S7).

The one-electron oxidation potential of \mathbf{Rot}^0 is indeed anodically shifted by $\Delta E_{1/2}^{-1} = +90 \text{ mV}$ compared to free \mathbf{Ax}^0 , indicating that the Sq-wheel interaction energetically hampers the first oxidation (Fig. 4e, f). We investigated the co-conformational equilibrium between ^{Sq}Rot and ^{Triaz}Rot in four oxidation states by exploring the co-conformational landscapes using the CREST code^{65,66} by GRIMME and subsequent evaluation of energetic stabilities employing various density functional approximations (Fig. 5 and Table S7). Whereas the neutral Sq station is energetically favored over the triazole station by 29 kJ/mol, the first oxidation renders the triazole station slightly more favorable than the Sq^{•+} station by 3.5 kJ/mol. Hence, one-electron oxidation results in molecular shuttling (^{Sq}Rot⁰ \rightarrow TriazRot⁺)



Fig. 3 | Structure of rotaxane Rot and chloride-induced shuttling. a Equilibrium between Ax, TLM and psRot & schematic depictions of these species. b Molecular structure of rotaxane Rot with schematic depiction. c Partial ¹H NMR spectrum of ^{Sq}Rot⁰. d Partial ¹H NMR spectrum of ^{Triaz}Rot⁰ \supset Cl⁻ after addition of 59 equiv. n-Bu₄NCl.

and a co-conformational distribution in which the wheel preferably adopts a distal position to Sq (Fig. 5).

Gradual oxidation of Rot^0 in an electron paramagnetic resonance (EPR) spectroelectrochemical experiment (dichloromethane, 298 K) gave an isotropic signal at g = 2.003 (Fig. 4c) confirming its transition into a stable

radical cation $Sq^{\bullet+}$. A low temperature spectrum at 223 K reveals the underlying hyperfine structure of the signal partially. The best fit, which is in excellent agreement with the experimental spectrum, was obtained with a simulation using a *g*-value of 2.004 and hyperfine coupling to two equivalent ¹⁴N nuclei and three sets of equivalent ¹H nuclei totaling 14 hydrogen nuclei



Fig. 4 | Opto-electrochemical investigation of the electronic and co-conformational properties of Ax and Rot. a Electrostatic potential surface maps of Ax in four different oxidation states. b Cyclic voltammograms (100 mV/s) of Ax and Rot. The second scan cycle is shown. The additional peaks at approximately -0.1 V in the second scan cycle, the i_{pa} / i_{pc} ratio, and the non-linear behavior of i_{pa} at different scan rates (inset) indicate an EC_i mechanism for the one-electron reduction of Ax. c EPR spectra of Ax⁺⁺ and Rot⁺⁺ (d) Spin density plots the radical species Ax⁺⁺, isovalue: 0.02 a.u. e Differential pulse voltammograms (10 mV/s scan rate, 25 mV modulation amplitude, 50 ms modulation time, 5 mV step potential, 0.5 s interval

time) of **Ax**, **Rot**, and **Rot** after addition of excess n-Bu₄NCl. The second oxidation reaction of the rotaxane in presence of chloride appears irreversible in both the DPV and CV. **f** Energy diagram, illustrating the stabilizing and destabilizing effects of the macrocycle and the presence of chloride on the first oxidation potentials of the Sq in **Ax** & **Rot**. Energies were calculated from (^a) redox potentials: $\Delta G = n \cdot F \cdot E$, (^b) binding constants: $\Delta G = -R \cdot T \cdot \ln(K_a)$ and (^c) according to HESS' law $\Delta G^c = \Sigma \Delta G^{a\&b}$. **g** The SCXRD-structure of **Ax** (thermal displace parameters at 50% probability level). **h** Annotation of discussed bond lengths (from DFT calculations) and rotational barriers within the bis(aminothienyl)squaraine core.



Fig. 5 | Structural investigation of the most stable co-conformational space of Rot. a Calculated most stable co-conformations of Rot in four different oxidation states with schematic depictions of each state. b Co-conformational electrochemical

and chemical equilibrium between six states of **Rot**. Only the most stable coconformations of each state are depicted.

(four ¹H thienyl, two ¹⁴N amine, eight ¹H N-C¹H₂) (Table S3). Supported by a spin density plot derived from DFT calculations, the experimental data indicate that the radical in Sq^{•+} is delocalized over the thiophene-Sq system (Fig. 4d and Fig. S43). The low temperature EPR spectrum of singly oxidized **Rot**^{•+} shows very similar spectral features indicating virtually no electronic effect of the wheel on the electronic coupling of the axle-centered radical cation. Similar findings have been observed for other hydrogen-bonded radicals⁴ and could result from weakened wheel–Sq interactions in **Rot**^{•+} and/or subsequent co-conformational changes as indicated by the electrochemical experiments.

The second one-electron oxidation ($Rot^{\bullet+}/Rot^{2+}$) occurs at a similar potential compared to that of the free $Ax^{\bullet+}$ ($\Delta E_{1/2}^2 = +10$ mV), indicating that hydrogen bonding between Sq and the wheel in **Rot**⁺ is significantly weakened compared to the neutral state (Fig. 4a, e, and f and 5). Consequently, the second oxidation of the deshielded Sq energetically resembles the free axle more than the first oxidation, as no further conformational equilibria contribute to the stabilization or destabilization of the system. Further evidence was derived from differential pulse voltammetry utilizing Rot⁰, both in the absence and presence of one equivalent *n*-Bu₄NCl (Fig. 4e). As the chloride-binding wheel in complex ^{Triaz}Rot⁰⊃Cl⁻ adopts a distal position to Sq, the first oxidation reaction displays a potential $(E_{1/2}^{1} = 0.43 \text{ V})$ similar to free **Ax⁰**. This illustrates that the increased oxidation potential in Rot⁰ is indicative of the wheels position: The first oxidation potential of Rot^0 is higher than that of free Ax^0 , because the stabilization energy in Rot⁰ arising from rotaxanation and the intramolecular wheel-Sq hydrogen bonds, needs to be compensated upon oxidation. As the chloride-addition already leads to a co-conformation $\bar{\text{Triaz}} Rot^0 \supset Cl^-$ in which the Sq is deshielded, the binding energy of the macrocycle is compensated for by the interaction between the wheel, triazole and chloride. The subsequent oxidation of the deshielded Sq thus requires a similar potential as the free axle. Further, the slightly lower oxidation potential of **Rot**⁰ in the presence of chloride suggests a stabilization of the oxidized $Rot^{\bullet+} \supset Cl^-$ by the strongly coordinating chloride anion⁶⁷. Further oxidation to $Rot^{2+} \supset Cl^{-}$ is irreversible, likely caused by a nucleophilic attack at the now sterically deprotected Sq core²⁸.

Theoretical calculations show doubly oxidized Rot^{2+} to have a pronounced energetic bias towards the triazole station of 18 kJ/mol (Fig. 5 and Table S7). The reduced electron density and thus, significantly lower H-bond accepting capability of Sq^{2+} is illustrated by electrostatic potential surface maps of the stoppered axle as found within **Rot**. The negative charge accumulation at Sq within **Rot**[•] and **Rot**⁰ and conversely, the lack of negative charge at Sq in **Rot**^{•+} and **Rot**²⁺ are depicted in Fig. S43. Thus, the H-bond accepting ability of Sq decreases drastically from Sq^{•-} to Sq²⁺, while more negative charge is retained at the triazole groups in the oxidized states. In accordance with the electrochemical and quantum chemical data the interaction between Sq and the wheel are already weak enough in **Rot**^{•+} that the wheel adopts a position close to the triazole motif. Quantum chemical investigation of this binding motif suggests H-bonding between two NH protons of the wheel and the electron rich N atoms of the triazol unit (Fig. S42).

To investigate the oxidation induced shuttling (^{Sq}Rot⁰ \Rightarrow ^{Triaz}Rot²⁺) by 1D and 2D NMR experiments, we developed a reliable method to obtain chemically stable solutions of Ax^{2+} and Rot^{2+} . Initial attempts using NOPF₆ as oxidant in dichloromethane- d_2 resulted in partial precipitation. In acetonitrile, the resulting dye solutions decompose within minutes. The solubility and lifetime in dichloromethane- d_2 of the Sq²⁺ species could be drastically increased (time scale of days) by addition of the weakly coordinating anion [Al(O-C(CF₃)₃)₄]⁻ ([pf]⁻) as Li[pf] during the oxidation with NOPF₆. Rot²⁺ did not display a stabilizing effect compared to the dicationic Ax^{2+} , as expected for the Sq being exposed in this state (see below for stability study). Oxidation reactions for both, Ax and Rot, are reversible, as demonstrated by using tin powder as reductant in an UV-Vis titration (SI section S7 Fig. S34).

The ¹H NMR spectrum of Ax^{2+} displays multiple sets of signals which can be attributed to rotational isomers of Ax^{2+} (Supporting Information Section S1, S6 and S7). Quantum chemical calculations reveal a four-fold increase of rotational barriers in Ax^{2+} compared to Ax^{0} (SI section S6 Table S4 and section S9 Table S5). At room temperature these rotations are free in Ax^{0} and hindered in Ax^{2+} leading to the observed spectral behavior. The much lower rotational barriers of Ax^{0} and Rot^{0} cause decoalescence of most signals only below -30 °C (SI section S8). Based on the calculated barriers for Ax^{0} , the rotation around the bonds between the Sq and the thiophene units should be frozen first, due to their higher barrier.

NOE cross peaks between proton signals of one of triazoles and methyl groups of the macrocycle indicate the wheel to be located at the triazole in **Rot**²⁺ (Figs. S6–S10). Multiple sets of signals and extensive peak broadening



Fig. 6 | UV-Vis (solid) spectra of Rot in three oxidation states and fluorescence (dashed) spectra of Rot⁰ and Rot²⁺. The inset photographs on each curve depict a vial containing dichloromethane solution of each species, the counterion is [pf]⁻.

in the 1D and 2D NMR spectra reveal a slow shuttling motion of the wheel between the two triazoles. Additionally, the non-bound side of the axle likely displays rotational isomerism like Ax^{2+} .

The stability of both oxidized states, $Rot^{\bullet+}$ and Rot^{2+} , on a laboratory time scale allowed us to study their optical properties (Fig. 6 and S34 and SI section S3). The first oxidation of Rot⁰ is accompanied by a hypsochromic shift of the most intense absorbance band, **Rot**⁰ ($\lambda_{max} = 665 \text{ nm}$) to **Rot**^{\bullet +} $(\lambda_{max} = 601 \text{ nm})$. The strongest band is further shifted by the second oxidation from **Rot**^{•+} ($\lambda_{max} = 601 \text{ nm}$) to **Rot**²⁺ ($\lambda_{max} = 509 \text{ nm}$). The emission band of $\mathbf{Rot}^{\mathbf{0}}$ ($\lambda_{\max} = 683 \text{ nm}, \Phi(\mathbf{Rot}^{\mathbf{0}}) = 4.0\%$) is close to the bands of similar reported bis-(aminothienyl)squaraine rotaxanes⁶⁸. Due to the disproportionation equilibrium from Rot⁺⁺ into Rot⁰ and Rot²⁺, strong emission bands of the neutral and doubly charged species limit the detection of potential fluorescence of the radical cation. Dicationic Rot²⁺ $(\lambda_{\text{max}} = 570 \text{ nm}, \Phi(\text{Rot}^{2+}) = 0.6\%)$ shows weak fluorescence, a photograph of the luminescence is depicted as inset in Fig. 6. In line with the ^{Triaz}Rot²⁺ co-conformation, the quantum yield is almost equal to that of Ax^{2+} $(\lambda_{\text{max}} = 548 \text{ nm}, \Phi(\mathbf{Ax}^{2+}) = 0.5\%)$ (SI section S3 Fig. S24 and Table S1). The change of absorption properties was supported by time-dependent DFT calculations, suggesting that charge-transfer effects upon excitation play a role in Ax^0 and Ax^{0+} . The difference density of the bright state displays a charge transfer from Sq to the neighboring thiophene moieties (Table S6).

To further elucidate the protection and deprotection in the neutral and +2 oxidation states, bleaching of the squaraine core in Ax^0 , Rot^0 , Ax^{2+} and Rot^{2+} by attack of nucleophiles was investigated using UV-Vis spectroscopy (SI section S10 Fig. 43). Tetramethylammonium hydroxide in dichloromethane proved effective in bleaching Ax^{2+} and Rot^{2+} , while Rot^0 was unaffected (Fig. S44b–d). While the half-life of Ax^{2+} (≈ 10 min) and Rot^{2+} (≈ 5 min) under these conditions is comparable, Rot^0 was stable (Fig. S44a). Because the squaraine core in the neutral rotaxane is shielded by the macrocycle from nucleophilic attack, no significant bleaching occurs. In line with the suggested co conformation $^{Triaz}Rot^{2+}$, the deshielded Sq²⁺ core bleaches similarly quickly as in the free Ax^{2+} .

The shielding effect of the macrocycle has been demonstrated for similar systems²⁸. As demonstrated above, low concentrations of hydroxide are not sufficient to bleach **Rot**⁰. At higher concentrations of OH⁻, the possibility of shuttling by binding of hydroxide in a similar motive as with chloride, cannot be excluded. Thus, for the comparison of the stability of **Ax**⁰ and **Rot**⁰, sterically demanding potassium 'butoxide was used (Fig. S44f, g). To increase the solubility of the nucleophile, 5% of 'BuOH were added to the solvent. In this mixture, free **Ax**⁰ was bleached (\approx 12 min), while **Rot**⁰ was affected only very little, indicating the expected protection of the squaraine core within the bound wheel (Fig. S44e).

Summary and conclusion

In this proof-of-concept study, we show that a co-conformationally dynamic rotaxane can serve as scaffold to either stabilize or expose highperforming dye molecules used for bioimaging applications, in this case a near-infrared squaraine dye. [2]Rotaxane Rot can be reversibly switched between four different redox states, resulting in two different co-conformations, in which the squaraine dye is either shielded (oxidation state 1 and 0) or exposed (oxidation state +1 and +2). Shielding results in an increase of the dye's lifetime, whereas the exposed states display physicochemical and optoelectronic properties similar to the non-complexed dye. At the same time, the squaraine-centered redox reactions allow for a stepwise tuning of the emission between 570 and 683 nm of the fluorescent Rot⁰ and Rot²⁺ states. A potential application involves the transport of labile fluorophores-stabilized by mechanical bonding-into a location where the co-conformational equilibrium of the rotaxane is triggered leading to the exposure of the fluorophore or a different reactive species. Such a system could be useful for applications in bioimaging (e.g. for multiplexing), sensing, and proximity labeling.

Methods

General

NMR experiments were performed on JEOL ECX 400, JEOL ECP 500, Bruker AVANCE 500, JEOL ECZ 600, and Bruker AVANCE 700 instruments. Residual solvent signals were used as the internal standards. All shifts are reported in ppm and NMR multiplicities are abbreviated as s (singlet), d (doublet), t (triplet), m (multiplet) and br (broad).

High-resolution ESI mass spectra were recorded on an Agilent 6210 ESI-TOF mass spectrometer. HPLC grade solvents were used for sample preparation and the samples were introduced into the ion source with a flow rate of $2-4 \,\mu$ L/min. UV/Vis spectra were recorded on a Varian Cary 50 Bio spectrometer with a xenon lamp. Fluorescence spectra were recorded on Infinite® M Nano⁺ (Tecan Deutschland GmbH, Crailsheim, Germany) and PerkinElmer Fl 6500 spectrometers. Suprasil glass cuvettes with pathlengths of 1 cm were used. Photoluminescence quantum yields (Φ) were measured with an integrating sphere setup from Hamamatsu (Quantaurus-QY C11347-11). All measurements were performed at room temperature using 10 mm × 10 mm long neck quartz cuvettes.

All reagents and solvents were obtained from commercial sources and used without further purification. Lithium tetrakis(perfluoro-^{tert}butoxy) aluminate was provided by Prof. Dr. Ingo Krossing and Malte Sellin (University of Freiburg, Germany). Dry solvents were purchased from Acros Organics. Deuterated solvents were purchased from Eurisotop and Deutero. Deuterated dichloromethane was dried by vigorous stirring with calcium hydride for 2 weeks at room temperature.

N-Ethyl-*N*-(2-(prop-2-yn-1-yloxy)ethyl)thiophen-2-amine⁶⁹, 4-tritylphenylazide⁴² and tetralactam macrocycle **TLM**⁷⁰ were synthesized according to literature procedures. Thin-layer chromatography was performed on silica gel coated plates with fluorescent indicator F254 (Macherey-Nagel). For column chromatography, silica gel (0.04-0.063 mm, Macherey-Nagel) was used.

Synthesis

Bis(aminothienyl)squaraine Axle Ax⁶⁸

Bis(aminothienyl)squaraine axle **Ax** was prepared according to a modified literature procedure⁶⁸ (Fig. 7): A mixture of toluene (750 mL) and *n*-butanol (250 mL) was azeotropically dried by refluxing on a DEAN-STARK trap (oil bath 145 °C) for 2 h. The collected water was drained and the solvent mixture was cooled under N₂-atmosphere. *N*-Ethyl-*N*-(2-(prop-2-yn-1-yloxy)ethyl)thiophen-2-amine (9.7 g, 47 mmol, 2.0 equiv.) and squaric acid (2.7 g, 24 mmol, 1.0 equiv.) were added. A color change from yellow to deep green to deep blue to black was observed. The reaction mixture was heated to reflux under N₂ on the DEAN-STARK trap (oil bath 145 °C) for 8 h. The solvents were removed under reduced pressure, and the metallic green residue was purified by column chromatography using dichloromethane-acetone (0% \rightarrow 30% acetone) as



Fig. 7 | Adapted procedure for the preparation of squaraine Ax.55.



Fig. 8 | Capping-synthesis of Rot by Cu^{II} catalyzed azide alkyne HUISGEN cycloaddition of trityl phenyl azide with the pseudo[2]rotaxane of Ax and TLM in dichloromethane. TBTA: tris((1-benzyl-4-triazolyl)methyl)amine.

eluent. Analytically pure Ax was obtained as a metallic green solid (6.50 g, 13 mmol, 56% yield), the analytical data are consistent with the reported values (Figs. S1, S2, and S16)⁶⁸.

¹**H NMR** (600 MHz, CD₂Cl₂): δ = 7.88 (d, *J* = 4.7 Hz, 2 H, d), 6.31 (d, *J* = 4.7 Hz, 2 H, e), 4.17 (d, *J* = 2.4 Hz, 4 H, k), 3.79 (t, *J* = 5.4 Hz, 4 H, j), 3.69 (t, *J* = 5.4 Hz, 4 H, i), 3.61 (q, *J* = 7.2 Hz, 4 H, g), 2.50 (t, *J* = 2.4 Hz, 2H, m), 1.30 (t, *J* = 7.2 Hz, 6 H, h) ppm.

¹³C{¹H} NMR (151 MHz, CD₂Cl₂): δ = 178.2, 171.5, 170.0, 138.1, 115.6, 109.3, 79.6, 75.1, 67.4, 58.9, 54.2, 50.4, 12.2 ppm.

HRMS (ESI): m/z calculated for $C_{26}H_{28}N_2O_4S_2$ [M]^{\bullet +} 496.1501, found 496.1513.

Squaraine-TLM Rotaxane Rot

Rotaxane **Rot** was prepared in the following manner (Fig. 8): A 25-ml dried Schlenk flask was charged with tetralactam macrocycle **TLM** (190 mg, 0.20 mmol, 1.0 equiv.) and **Ax** (200 mg, 0.40 mmol, 2.0 equiv.) under N₂ counterflow. Dry dichloromethane (8 mL) was added, the flask was sealed, and the mixture was stirred for 1 h at room temperature. After that, 4-trityl-phenylazide (350 mg, 0.96 mmol, 4.8 equiv.), Cu(ACN)₄BF₄ (25 mg, 0.08 mmol, 0.4 equiv.) and tris((1-benzyl-4-triazolyl)methyl)amine (TBTA) (43 mg, 0.08 mmol, 0.4 equiv.) were added under N₂ counterflow. The flask was sealed again and stirred overnight at room temperature. The mixture was purified by silica column chromatography using step gradient elution

 $(0.8\% \rightarrow 2.6\%$ methanol in dichloromethane). Rot eluted with $1.3\% \rightarrow 1.5\%$ methanol content.

Analytically pure **Rot** was obtained as a deep blue crystalline solid (140 mg, 0.06 mmol, 32% based on **TLM**) (Fig. S11, 12, and 16).

¹**H NMR** (600 MHz, CD₂Cl₂): δ = 10.34 (b, *NH*, 2 H, ε), 9.18 (b, 1 H, **χ**), 8.76 (b, *NH*, 2 H, **o**), 8.42 (d, *J* = 7.8 Hz, 2 H, **β**), 8.31 (d, *J* = 1.4 Hz, 2 H, **σ**), 8.09 (t, *J* = 7.8 Hz, 1 H, **α**), 7.98 (s, 2 H, **m**), 7.64 (d, *J* = 8.8 Hz, 4 H, **o**), 7.51 (d, *J* = 4.7 Hz, 2 H, **d**), 7.45 (d, *J* = 8.8 Hz, 4 H, **p**), 7.25-7.30 (m, 24 H, **t & u**), 7.20-7.23 (m, 6 H, **v**), 6.83 & 6.82 (2x s, 8 H, **ι&x**²), 6.17 (d, *J* = 4.7 Hz, 2 H, **e**), 4.62 (s, 4 H, **k**), 3.72 (t, *J* = 5.4 Hz, 4H, **j**), 3.58 (t, *J* = 5.4 Hz, 4H, **i**), 3.49 (q, *J* = 7.2 Hz, 4 H, **g**), 2.20–2.37 (m, 8 H, **µ**), 2.07 & 2.06 (2x s, 24 H, **0&d**²), 1.61 (m, 8 H, **v**), 1.49 (m, 4 H, **ξ**), 1.39 (s, 9 H, **φ**), 1.19 (t, *J* = 7.2 Hz, 6 H, **h**) ppm.

¹³C{¹H} NMR (175 MHz, CD₂Cl₂): δ = 179.2, 169.9, 167.7, 164.7, 162.6, 153.7, 149.7, 148.4, 147.9, 147.4, 146.7, 145.2, 139.3, 137.9, 135.4, 135.2, 135.0, 134.5, 132.7, 131.8, 131.6, 131.3, 128.8, 128.2, 126.6, 125.7, 125.4, 125.1, 123.3, 121.4, 120.0, 114.1, 109.3, 67.7, 65.3, 64.8, 53.8, 50.3, 45.2, 35.6, 34.7, 31.5, 30.1, 26.8, 23.4, 19.1, 19.0, 12.1 ppm.

HRMS (ESI): m/z calculated for $C_{139}H_{137}N_{13}O_8S_2$ [M]^{$\bullet+$} 2180.0154, found 2180.0108.

Electrochemical switching

Cyclic voltammetry (CV) (Figs. S26 and S27) and differential pulse voltammetry (DPV) (Fig. S27 and Table S2) were performed on an Autolab PGSTAT302N potentiostat using a three-electrode configuration: a freshly polished glassy carbon working electrode, a platinum wire counter electrode, and a silver wire pseudoreference electrode. All measurements were conducted at least three times and with a broad range of different scan rates (10–2000 mV/s) (Figs. S26 and S29). The decamethylferrocene/decamethylferrocenium ([FeCp₂*]^{+/0}) couple was used as internal reference. Dry and nitrogen-purged 1,2-dichloroethane (DCE) as solvent, *n*-Bu₄NPF₆ (0.1 M) as electrolyte, and an analyte concentration of 1.0 mM were used for all measurements. Binding of chloride to **Rot**⁰ in the presence of Bu₄NPF₆ was confirmed by–ESI HRMS (Fig. S28). The error of redox potentials derived by CV and DPV measurements is estimated to be smaller than ± 10 mV.

Digital simulations (Fig. S29) were performed with the DigiElch 8 Professional software package (ElchSoft GbR) using the Butler-Volmer equation. The surface area of the working electrode was set to 0.05 cm², and the starting concentration of analytes was set to 1.0 mM. The chargetransfer coefficients α were set to the initial value of 0.5, and the heterogeneous rate constants k_s were estimated by the peak separation and set between 0.01 and 0.1 cm/s. The diffusion coefficient was set to its default value of 1×10^{-5} cm²/s. The fitting model for the experimental CV data is based on an (EE)EC_i mechanism for the direction of a cathodic sweep (starting from the +2 oxidation state). The reaction kinetics, i. e. the rate constant of the chemical reaction step k_{fb} of **Ax**^{\bullet} and **Rot**^{\bullet} were evaluated according to an irreversible (pseudo)first-order reaction based on four different scan rates (25, 100, 250, and 1000 mV/s).

Quantum chemical calculations

While the available crystal structure of Ax could be used as a starting point for the investigation of Ax, for Rot a search of the conformational landscape was undertaken using Stefan Grimme's GFN2-xTB⁶⁵ and CREST⁶⁶ codes. All subsequent structure optimizations were performed using the Turbomole program package⁷¹ (version 7.6) employing the PBEh-3c⁷² composite method in combination with the COSMO⁷³ solvent model. Single point electronic energies were evaluated at the PBE0-D3(BJ)74,75, B3LYP-D3(BJ)76, BHLYP-D3(BJ)⁷⁷, and ω B97X-D3⁷⁸ levels of density functional theory, all in addition with the SMD⁷⁹ solvent model with keyword "dichloromethane", utilizing the ORCA program suite (version 5.0.4)⁸⁰. A def2-QZVP⁸¹ basis set was used for Ax, a def2-TZVP basis set for Rot and the stoppered axle. Optical transitions were calculated for Ax using the same density functional approximations, but with a def2-TZVP basis set. Transition states for the two rotational barriers in Ax were obtained by manually displacing the respective dihedral angle and subsequent transition state optimization following the lowest eigenvalue of the Hessian ("imaginary frequency"). Negative eigenvalues above -20 cm^{-1} were neglected when assessing the quality of the transition state structures after a run had converged. Transition state runs were performed with charges 0 and +2. All other calculations were done with charges -1, 0, +1, and +2, respectively.

Data availability

Crystallographic data for the structures reported in this article have been deposited at the Cambridge Crystallographic Data Centre, under deposition numbers CCDC 2332801 (**Ax**) and 2332802 (**psRot**). Copies of the data can be obtained free of charge via https://www.ccdc.cam.ac.uk/structures/. All other relevant data generated and analyzed during this study, which include experimental, spectroscopic, crystallographic and computational data, are included in this article and the Supplementary Data 1 file. XRD structure files for **Ax** (Dataset 1, Checkcif Dataset 3), and for **psRot** (Dataset 2, Checkcif Dataset 4) are attached. All calculated structures are contained in Dataset 5.

Received: 21 February 2024; Accepted: 19 September 2024; Published online: 04 October 2024

References

 Duda, R. L. Protein chainmail: catenated protein in viral capsids. *Cell* 94, 55–60 (1998).

- 2. Mateu, M. G. Assembly, stability and dynamics of virus capsids. *Arch. Biochem. Biophys.* **531**, 65–79 (2013).
- Petsko, G. A. & Ringe, D. Protein Structure and Function, Vol. 195 (Oxford Univ. Press, London, 2009).
- Benson, C. R. et al. Extreme stabilization and redox switching of organic anions and radical anions by large-cavity, CH hydrogenbonding cyanostar macrocycles. *J. Am. Chem. Soc.* 138, 15057–15065 (2016).
- Mal, P., Breiner, B., Rissanen, K. & Nitschke, J. R. White phosphorus is air-stable within a self-assembled tetrahedral capsule. *Science* 324, 1697–1699 (2009).
- Pakhomov, A. A. & Martynov, V. I. GFP family: structural insights into spectral tuning. *Chem. Biol.* 15, 755–764 (2008).
- Walsh, C. T. & Wencewicz, T. A. Flavoenzymes: versatile catalysts in biosynthetic pathways. *Nat. Prod. Rep.* 30, 175–200 (2013).
- 8. Allen, J. P. & Williams, J. C. Photosynthetic reaction centers. *FEBS Lett.* **438**, 5–9 (1998).
- Hong, G., Antaris, A. L. & Dai, H. Near-infrared fluorophores for biomedical imaging. *Nat. Biomed. Eng.* 1, 1–22 (2017).
- Lee, M. H., Kim, J. S. & Sessler, J. L. Small molecule-based ratiometric fluorescence probes for cations, anions, and biomolecules. *Chem. Soc. Rev.* 44, 4185–4191 (2015).
- Seath, C. P., Trowbridge, A. D., Muir, T. W. & MacMillan, D. W. C. Reactive intermediates for interactome mapping. *Chem. Soc. Rev.* 50, 2911–2926 (2021).
- Körner, S. K., Tucci, F. C., Rudkevich, D. M., Heinz, T. & Rebek, J. J. A self-assembled cylindrical capsule: new supramolecular phenomena through encapsulation. *Chem. Eur. J.* 6, 187–195 (2000).
- Breiner, B., Clegg, J. K. & Nitschke, J. R. Reactivity modulation in container molecules. *Chem. Sci.* 2, 51–56 (2011).
- Galan, A. & Ballester, P. Stabilization of reactive species by supramolecular encapsulation. *Chem. Soc. Rev.* 45, 1720–1737 (2016).
- Iwasawa, T., Hooley, R. J. & Rebek, J. Stabilization of labile carbonyl addition intermediates by a synthetic receptor. *Science* **317**, 493–496 (2007).
- Yoshizawa, M., Klosterman, J. K. & Fujita, M. Functional molecular flasks: new properties and reactions within discrete, self-assembled hosts. *Angew. Chem. Int. Ed.* 48, 3418–3438 (2009).
- Eelkema, R., Maeda, K., Odell, B. & Anderson, H. L. Radical cation stabilization in a cucurbituril oligoaniline rotaxane. *J. Am. Chem. Soc.* 129, 12384–12385 (2007).
- Li, H. et al. Mechanical bond-induced radical stabilization. J. Am. Chem. Soc. 135, 456–467 (2013).
- Movsisyan, L. D. et al. Polyyne rotaxanes: stabilization by encapsulation. J. Am. Chem. Soc. 138, 1366–1376 (2016).
- Patrick, C. W. et al. Masked alkynes for synthesis of threaded carbon chains. *Nat. Chem.* 16, 193–200 (2024).
- 21. Sun, J. et al. Mechanical-bond-protected, air-stable radicals. J. Am. Chem. Soc. 139, 12704–12709 (2017).
- 22. Hegemann, J. D. Factors governing the thermal stability of lasso peptides. *Chembiochem* **21**, 7–18 (2020).
- Maksimov, M. O., Pan, S. J. & James Link, A. Lasso peptides: structure, function, biosynthesis, and engineering. *Nat. Prod. Rep.* 29, 996–1006 (2012).
- Zimmermann, M., Hegemann, J. D., Xie, X. & Marahiel, M. A. The astexin-1 lasso peptides: biosynthesis, stability, and structural studies. *Chem. Biol.* 20, 558–569 (2013).
- Allen, C. D. & Link, A. J. Self-assembly of catenanes from lasso peptides. J. Am. Chem. Soc. 138, 14214–14217 (2016).
- Schröder, H. V., Stadlmeier, M., Wühr, M. & Link, A. J. The shuttling cascade in lasso peptide benenodin-1 is controlled by non-covalent interactions. *Chem. Eur. J.* 28, e202103615 (2022).
- Bayro, M. J. et al. Structure of antibacterial peptide microcin J25: a 21-residue lariat protoknot. *J. Am. Chem. Soc.* **125**, 12382–12383 (2003).

- Gassensmith, J. J., Baumes, J. M. & Smith, B. D. Discovery and early development of squaraine rotaxanes. *Chem. Commun.* 14, 6329–6338 (2009).
- Collins, C. G., Peck, E. M., Kramer, P. J. & Smith, B. D. Squaraine rotaxane shuttle as a ratiometric deep-red optical chloride sensor. *Chem. Sci.* 4, 2557 (2013).
- Arunkumar, E., Forbes, C. C., Noll, B. C. & Smith, B. D. Squarainederived rotaxanes: sterically protected fluorescent near-IR dyes. *J. Am. Chem. Soc.* **127**, 3288–3289 (2005).
- Escobedo, J. O., Rusin, O., Lim, S. & Strongin, R. M. NIR dyes for bioimaging applications. COCHBI 14, 64–70 (2010).
- 32. He, J. et al. Squaraine dyes for photovoltaic and biomedical applications. *Adv. Funct. Mater.* **31**, 2008201 (2021).
- Ilina, K. et al. Squaraine dyes: molecular design for different applications and remaining challenges. *Bioconjug. Chem.* **31**, 194–213 (2020).
- Erbas-Cakmak, S., Leigh, D. A., McTernan, C. T. & Nussbaumer, A. L. Artificial molecular machines. *Chem. Rev.* 115, 10081–10206 (2015).
- Heard, A. W. & Goldup, S. M. Simplicity in the design, operation, and applications of mechanically interlocked molecular machines. ACS Cent. Sci. 6, 117–128 (2020).
- Pease, A. R. et al. Switching devices based on interlocked molecules. Acc. Chem. Res. 34, 433–444 (2001).
- Sauvage, J.-P. From chemical topology to molecular machines (Nobel Lecture). Angew. Chem. Int. Ed. 56, 11080–11093 (2017).
- Stoddart, J. F. Mechanically interlocked molecules (MIMs)-molecular shuttles, switches, and machines (Nobel Lecture). *Angew. Chem. Int. Ed.* 56, 11094–11125 (2017).
- Kaufmann, L. et al. Substituent effects on axle binding in amide pseudorotaxanes: comparison of NMR titration and ITC data with DFT calculations. Org. Biomol. Chem. 10, 5954–5964 (2012).
- 40. Gassensmith, J. J. et al. Squaraine rotaxane as a reversible optical chloride sensor. *Chem. Eur. J.* **16**, 2916–2921 (2010).
- Castellano, R. Progress toward understanding the nature and function of C-H...O interactions. COC 8, 845–865 (2004).
- Dzyuba, E. V. et al. CH-O hydrogen bonds in "clicked" diketopiperazinebased amide rotaxanes. Org. Lett. 13, 4838–4841 (2011).
- 43. Altieri, A. et al. Electrochemically switchable hydrogen-bonded molecular shuttles. *J. Am. Chem. Soc.* **125**, 8644–8654 (2003).
- Gaedke, M. et al. Dual-stimuli pseudorotaxane switches under kinetic control. Org. Chem. Front. 8, 3659–3667 (2021).
- Gaedke, M. et al. Sequence-sorted redox-switchable hetero[3] rotaxanes. Org. Chem. Front. 9, 64–74 (2021).
- Gaedke, M. et al. Chiroptical inversion of a planar chiral redoxswitchable rotaxane. *Chem. Sci.* **10**, 10003–10009 (2019).
- Hupatz, H. et al. Thermodynamic and electrochemical study of tailormade crown ethers for redox-switchable (pseudo)rotaxanes. *Beilstein J. Org. Chem.* 16, 2576–2588 (2020).
- Schröder, H. V. et al. A divalent pentastable redox-switchable donoracceptor rotaxane. *Chem. Eur. J.* 23, 2960–2967 (2017).
- Schröder, H. V. et al. Switchable synchronisation of pirouetting motions in a redox-active 3rotaxane. *Nanoscale* **10**, 21425–21433 (2018).
- Schröder, H. V. & Schalley, C. A. Tetrathiafulvalene a redoxswitchable building block to control motion in mechanically interlocked molecules. *Beilstein J. Org. Chem.* 14, 2163–2185 (2018).
- Schröder, H. V. et al. Impact of mechanical bonding on the redoxswitching of tetrathiafulvalene in crown ether-ammonium 2rotaxanes. *Chem. Sci.* 8, 6300–6306 (2017).
- Schröder, H. V. et al. Accordion-like motion in electrochemically switchable crown ether/ammonium oligorotaxanes. *Angew. Chem. Int. Ed.* 58, 3496–3500 (2019).
- Schröder, H. V., Wollschläger, J. M. & Schalley, C. A. Redoxcontrolled self-inclusion of a lasso-type pseudo1rotaxane. *Chem. comm.* 53, 9218–9221 (2017).
- 54. Maltese, V. et al. Electro-optical properties of neutral and radical ion thienosquaraines. *Chem. Eur. J.* **22**, 10179–10186 (2016).

- 55. Barnes, J. C. et al. A radically configurable six-state compound. *Science* **339**, 429–433 (2013).
- Griller, D. & Ingold, K. U. Persistent carbon-centered radicals. Acc. Chem. Res. 9, 13–19 (1976).
- Sowndarya, S. V. S., St John, P. C. & Paton, R. S. A quantitative metric for organic radical stability and persistence using thermodynamic and kinetic features. *Chem. Sci.* 12, 13158–13166 (2021).
- Forbes, M. D. E. Carbon-Centered Free Radicals and Radical Cations. Structure, Reactivity, and Dynamics Vol. 392 (Wiley, Hoboken, NJ, 2010).
- Fioravanti, G. et al. Three state redox-active molecular shuttle that switches in solution and on a surface. *J. Am. Chem. Soc.* 130, 2593–2601 (2008).
- Supur, M., El-Khouly, M. E., Seok, J. H., Kay, K.-Y. & Fukuzumi, S. Elongation of lifetime of the charge-separated state of ferrocenenaphthalenediimide-60fullerene triad via stepwise electron transfer. *J. Phys. Chem. A* **115**, 14430–14437 (2011).
- Tang, B., Zhao, J., Xu, J.-F. & Zhang, X. Tuning the stability of organic radicals: from covalent approaches to non-covalent approaches. *Chem. Sci.* **11**, 1192–1204 (2020).
- 62. Spruell, J. M. et al. Highly stable tetrathiafulvalene radical dimers in [3] catenanes. *Nat. Chem.* **2**, 870–879 (2010).
- Song, Q., Li, F., Wang, Z. & Zhang, X. A supramolecular strategy for tuning the energy level of naphthalenediimide: Promoted formation of radical anions with extraordinary stability. *Chem. Sci.* 6, 3342–3346 (2015).
- Ziganshina, A. Y., Ko, Y. H., Jeon, W. S. & Kim, K. Stable pi-dimer of a tetrathiafulvalene cation radical encapsulated in the cavity of cucurbit8uril. *Chem. Commun.* 7, 806–7 (2004).
- Grimme, S., Bannwarth, C. & Shushkov, P. A robust and accurate tight-binding quantum chemical method for structures, vibrational frequencies, and noncovalent interactions of large molecular systems parametrized for all spd-block elements (Z = 1–86). *J. Chem. Theory Comput.* 13, 1989–2009 (2017).
- Pracht, P., Bohle, F. & Grimme, S. Automated exploration of the lowenergy chemical space with fast quantum chemical methods. *Phys. Chem. Chem. Phys.* 22, 7169–7192 (2020).
- Witte, J. F., Wasternack, J., Wei, S., Schalley, C. A. & Paulus, B. The interplay of weakly coordinating anions and the mechanical bond: a systematic study of the explicit influence of counterions on the properties of (Pseudo)rotaxanes. *Molecules (Basel, Switzerland)* 28, 3077 (2023).
- Peck, E. M. et al. Rapid macrocycle threading by a fluorescent dyepolymer conjugate in water with nanomolar affinity. *J. Am. Chem. Soc.* 137, 8668–8671 (2015).
- Spence, G. T., Hartland, G. V. & Smith, B. D. Activated photothermal heating using croconaine dyes. *Chem. Sci.* 4, 4240 (2013).
- Braun, O., Hünten, A. & Vögtle, F. Template synthesis of rotaxanes with carbamate-linked axles. J. Prakt. Chem. 341, 542–547 (1999).
- Ahlrichs, R., Bär, M., Häser, M., Horn, H. & Kölmel, C. Electronic structure calculations on workstation computers: the program system turbomole. *Chem. Phys. Lett.* **162**, 165–169 (1989).
- Grimme, S., Brandenburg, J. G., Bannwarth, C. & Hansen, A. Consistent structures and interactions by density functional theory with small atomic orbital basis sets. *J. Chem. Phys.* 143, 54107 (2015).
- Klamt, A. & Schüürmann, G. COSMO: a new approach to dielectric screening in solvents with explicit expressions for the screening energy and its gradient. *J. Chem. Soc. Perkin Trans.* 2, 799–805 (1993).
- Perdew, J. P., Ernzerhof, M. & Burke, K. Rationale for mixing exact exchange with density functional approximations. *J. Chem. Phys.* 105, 9982–9985 (1996).
- Grimme, S., Antony, J., Ehrlich, S. & Krieg, H. A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu. *J. Chem. Phys.* 132, 154104 (2010).

- Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. J. Chem. Phys. 98, 5648–5652 (1993).
- Becke, A. D. A new mixing of Hartree–Fock and local densityfunctional theories. J. Chem. Phys. 98, 1372–1377 (1993).
- Chai, J.-D. & Head-Gordon, M. Systematic optimization of long-range corrected hybrid density functionals. J. Chem. Phys. 128, 84106 (2008).
- Marenich, A. V., Cramer, C. J. & Truhlar, D. G. Universal solvation model based on solute electron density and on a continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions. J. Phys. Chem. B 113, 6378–6396 (2009).
- Neese, F. Software update: the ORCA program system, version 4.0. WIREs Comput. Mol. Sci. 8, e1327 (2018).
- Weigend, F. & Ahlrichs, R. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: design and assessment of accuracy. *Phys. Chem. Chem. Phys.* 7, 3297–3305 (2005).

Acknowledgements

We would like to thank Prof. Dr. Ingo Krossing and Malte Sellin from the University of Freiburg for supplying us with a generous sample of lithium tetrakis(perfluoro-^{tert}butoxy)aluminate. We want to thank the Deutsche Forschungsgemeinschaft (DFG; grants PA 1360/16-1 and SCHA 893/14-1) for funding this research. This work was also supported by the Estonian Research Council grant PUTJD911. We are grateful for the assistance of the Core Facility BioSupraMoI. The computing facilities of Freie Universität Berlin (Zentrale Einrichtung für Datenverarbeitung) and the Alliance for National High-Performance Computing (Verbund für Nationales Hochleistungsrechnen - NHR) are gratefully acknowledged for providing computational resources. We thank Jana Hildebrandt and the NMR department of the Humboldt-Universität zu Berlin for the VT-NMR measurements and Sebastian Müller for help with the synthesis of **Ax**.

Author contributions

J. Wasternack: Conceptualization, formal analysis, investigation, data curation. writing-original draft preparation. writing-review and editing. visualization, supervision. H. V. Schröder: Conceptualization, formal analysis, investigation, writing-original draft preparation, writing-review and editing, visualization, supervision. J. F. Witte: Formal analysis, investigation, writing-review and editing, visualization. M. Ilisson: Conceptualization, formal analysis, investigation, data curation, writingoriginal draft preparation, writing-review and editing, visualization, supervision. H. Hupatz: Conceptualization, formal analysis, investigation, data curation, writing-review and editing, supervision. J. Hille: Investigation, writing-review and editing, supervision. M. Gaedke: Formal analysis, investigation, writing-review and editing. A. M. Valkonen: Formal analysis, investigation, writing-review and editing. S. Sobottka: Formal analysis, investigation, writing-review and editing. A. Krappe: Investigation, writing-review and editing. M. Schubert: Formal analysis, investigation, writing-review and editing. B. Paulus: Resources, writing-review and

editing, supervision, project administration, funding acquisition. K. Rissanen: Resources, writing—review and editing, supervision, project administration, funding acquisition. B. Sarkar: Resources, writing—review and editing, supervision, project administration, funding acquisition. S. Eigler: Resources, writing—review and editing, supervision, project administration, funding acquisition. U. Resch-Genger: Resources, writing review and editing, supervision, project administration, funding acquisition. C. A. Schalley: Resources, writing—review and editing, supervision, project administration, funding acquisition. All authors have read and agreed to the published version of the manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at https://doi.org/10.1038/s42004-024-01312-1.

Correspondence and requests for materials should be addressed to Christoph A. Schalley.

Peer review information *Communications Chemistry* thanks the anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at http://www.nature.com/reprints

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2024