

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Vähäsarja, Kati; Salmela, Sanna; Villberg, Jari; Rintala, Pauli; Vanhala, Mauno; Saaristo, Timo; Peltonen, Markku; Keinänen-Kiukaanniemi, Sirkka; Korpi-Hyövälti, Eeva; Kujala, Urho M.; Moilanen, Leena; Niskanen, Leo; Oksa, Heikki; Poskiparta, Marita

Title: Perceived need to increase physical activity levels among adults at high risk of type 2 diabetes. A cross-sectional analysis within a community-based diabetes prevention project FIN-D2D

Year: 2012

Version:

Please cite the original version:

Vähäsarja, K., Salmela, S., Villberg, J., Rintala, P., Vanhala, M., Saaristo, T., Peltonen, M., Keinänen-Kiukaanniemi, S., Korpi-Hyövälti, E., Kujala, U., Moilanen, L., Niskanen, L., Oksa, H. & Poskiparta, M. (2012). Perceived need to increase physical activity levels among adults at high risk of type 2 diabetes. A cross-sectional analysis within a community-based diabetes prevention project FIN-D2D. *BMC Public Health*, 12 (514). doi:10.1186/1471-2458-12-514

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

RESEARCH ARTICLE

Open Access

Perceived need to increase physical activity levels among adults at high risk of type 2 diabetes. A cross-sectional analysis within a community-based diabetes prevention project FIN-D2D

Kati Vähäsarja^{1*}, Sanna Salmela¹, Jari Villberg¹, Pauli Rintala², Mauno Vanhala^{3,4,5}, Timo Saaristo^{6,7}, Markku Peltonen⁸, Sirkka Keinänen-Kiukaanniemi^{9,10,11}, Eeva Korpi-Hyövälti¹², Urho M Kujala¹, Leena Moilanen¹³, Leo Niskanen^{14,15}, Heikki Oksa^{7,16} and Marita Poskiparta¹

Abstract

Background: Increased physical activity is a cornerstone of type 2 diabetes prevention. The perception of a need to change is considered essential in behaviour change processes. However, the existing literature on individuals' perceived need to change health behaviour is limited. In order to improve understanding of diabetes prevention through increased physical activity levels (PAL), we assessed factors associated with perceiving a need to increase PAL among adults at high risk of diabetes.

Methods: Opportunistic screening was used within a primary-care based lifestyle intervention covering 10 149 men and women at high risk of type 2 diabetes. Data were obtained at baseline visits. The explored determinants were demographic, anthropometric/clinical, behavioural and psychosocial characteristics, along with four categories of PAL awareness. Logistic regression was used in the analysis.

Results: 74% of men (n = 2 577) and 76% of women (n = 4 551) perceived a need to increase their PAL. The participants most likely to perceive this need were inactive, had a larger waist circumference, rated their PAL as insufficient, and were at the contemplation stage of change. Smoking, elevated blood pressure, dyslipidaemia, and a family history of diabetes were not associated with this perception. The likelihood was also greater among women with less perceived fitness and less education. Demographic factors other than education did not determine participants' perceived need to increase PAL. PAL overestimators were less likely to perceive the need to increase their PAL than realistic inactive participants.

Conclusions: Subjective rather than objective health factors appear to determine the perception of a need to increase PAL among adults at high risk of diabetes. Client perceptions need to be evaluated in health counselling in order to facilitate a change in PAL. Practical descriptions of the associations between metabolic risk factors, PAL, and diabetes are needed in order to make the risk factors concrete for at-risk individuals.

Keywords: Physical activity, Exercise, Behaviour change, Perception, Awareness, Type 2 diabetes, Prevention, FIN-D2D, Lifestyle intervention

* Correspondence: kati.vahasarja@jyu.fi

¹Department of Health Sciences, University of Jyväskylä, Jyväskylä 40014, Finland

Full list of author information is available at the end of the article

Background

Extensive literature has indicated that type 2 diabetes can be prevented or delayed by increased levels of physical activity (PAL) [1-8]. People with diabetes are claimed to be least ready to increase their physical activity levels (PAL), to a greater degree than people without diabetes or people with heart disease, elevated blood pressure, or dyslipidaemia [9]. However, inclinations toward PAL behaviour change among the high-risk population remain imperfectly understood. It has been suggested [10,11] that perceived risk will increase protective behaviour. Indeed, the vast majority of participants at risk of diabetes have been reported as intending to increase their PAL [12], or as having tried to increase their PAL during the past twelve months [13]. However, it has been reported elsewhere that diabetes risk perception is not associated with physical activity intention among adults at risk of type 2 diabetes [14], and that persons with a higher diabetes risk perception do not have a greater intention than others to increase their PAL [15].

An individual's perceived need to change behaviour is regarded as a pre-step toward an intention [16-18]. However, only a few studies have assessed the perceived need to change health behaviour: three studies found perceived need to significantly predict the intention to improve diet [19-21], whereas no association was found between the perceived need and an intention to exercise [22]. The authors examined the perceived need to change health behaviour as a continuum within the *theory of planned behaviour* [23], taking the view that perceived need reflects perceived benefits, but that the perceptions of need and benefits are distinct. It was assumed that a person may have a positive attitude toward a behaviour, but not necessarily feel a need to change the behaviour if (i) the costs outweigh the benefits, (ii) the outcome of the behaviour is not valued highly by the person, or (iii) if the outcome is believed to be achievable by other means.

The concept of perceived need has been also associated with the *transtheoretical theory* (TTM) [17,24]. TTM suggests that individuals go through five distinct stages according to their motivational readiness to change behaviour. Individuals at the *precontemplation* stage are not regularly physically active, do not view their behaviour as a problem, and may therefore not perceive the need to change their behaviour. Once an individual becomes aware of his/her problem behaviour he/she may shift to the *contemplation* stage and continue to the *preparation*, *action*, and *maintenance* stages.

The third approach emphasising the role of perceived need in the change process is a counselling approach called *motivational interviewing* (MI) [18,25]. MI focuses primarily on enhancing the intrinsic motivation to

change risk behaviour among individuals at the precontemplation and contemplation stages (who are reluctant or hesitant about the change). MI proposes that motivation to change increases in parallel with the individual's own arguments regarding his/her perceived *need*, *desire*, *ability*, and *reasons* for change. These arguments strongly predict commitment, which in turn predicts change [26-28].

TTM [29-32] and MI [33,34] have both been widely applied to physical activity, and to type 2 diabetes prevention [35,36]. Even though these approaches regard a person's perceived need to change as an essential preparatory step for the intention and commitment to behaviour change, the concept has rarely been examined. Similarly, the existing literature on the correlates of the pre-contemplation stage is limited, and has tended to focus mainly on psychosocial aspects [15,37,38].

Perceiving the need to increase PAL is crucial for a population at high risk of type 2 diabetes. Thus, it is important to explore which factors determine this need. We examined demographic, anthropometric/clinical, behavioural, and psychosocial factors as determinants of this need perception among adults at high risk of type 2 diabetes. Since previous studies have shown misperception of one's own physical activity level to be a common phenomenon, and also to be associated with physical activity intention [37,39-43], we further examined the perceived need to increase PAL across four categories of physical activity awareness groups.

Methods

Study design and participants

FIN-D2D, an implementation project within a national programme for the prevention of type 2 diabetes, was conducted by the Finnish Diabetes Association in five Finnish hospital districts covering a population of 1.5 million [44,45]. The collaborating bodies were the National Public Health Institute, the Ministry of Social Affairs and Health and the FIN-D2D Study Group. The specific aims were to improve the screening of people at risk of diabetes and to detect undiagnosed diabetes. The programme included intensified interventions among high-risk individuals as part of normal clinical practice (for the full study protocol see [46,47]).

Opportunistic screening was conducted in health care centres and pharmacies, and at various public venues and events (e.g. health fairs), in order to screen individuals at high risk of type 2 diabetes. A modified Finnish diabetes risk score (FINDRISC) was used for the screening process [48]. Individuals were referred to primary care for lifestyle intervention on a voluntary basis if they met any of the following criteria (i) a FINDRISC test score ≥ 15 , (ii) a history of gestational diabetes, (iii) a history of impaired glucose tolerance or impaired fasting

Table 1 Characteristics of the study population

	Men (n = 2 577)	Women (n = 4 551)
	%	%
Age (years)		
<45	14	13
45-54	32	30
55-64	20	21
≥65	34	35
Marital status***		
Married/cohabiting	77	72
Other	23	28
Educational level		
Low	41	39
Intermediate	50	53
High	9	9
Occupational status***		
Manual work	34	9
Non-manual work	25	44
Retired	34	35
Not employed	7	13
Family history of diabetes ^{1***}	57	66
Elevated blood pressure ^{2***}	88	82
Dyslipidaemia ^{3***}	30	40
Obesity ^{4***}	57	63
Waist circumference ^{5***}		
Normal	8	3
Elevated risk	22	10
High risk	70	87
Currently smoking ^{6***}	23	15
Physically inactive ⁷	65	65
Self-rated fitness*		
Low	22	25
Satisfactory	50	48
High	28	27
Stage of change***		
Precontemplation	11	7
Contemplation	29	24
Preparation	39	46
Action	12	14
Maintenance	9	9
Physical activity level perceived as sufficient*	33	30
Perceives the need to increase physical activity level	74	76

¹ At least one of the first-degree relatives had diabetes (type 1 or 2).
² Systolic blood pressure ≥ 130 mmHg and/or diastolic blood pressure ≥ 85 mmHg.
³ HDL < 1.03 mmol/l (men), < 1.29 mmol/l (women); LDL ≥ 3 mmol/l or/and triglycerides ≥ 1.7 mmol/l, or medication for these lipid abnormalities.
⁴ Body mass index (BMI) ≥ 30 kg/m².
⁵ Normal: < 94 cm (m), < 80 cm (w); elevated risk: 94-101 cm (m), 80-87 cm (w); high risk: ≥ 102 cm (m), ≥ 88 cm (w).
⁶ Includes occasional and regular smokers.
⁷ Engaging in physical exercise < 30 minutes three times a week.
* $p < 0.05$ (for the statistical difference between the sexes).
** $p < 0.01$.
*** $p < 0.001$.

glucose, or (iv) a history of coronary heart disease. After identification, consenting high-risk individuals attended health check-ups conducted in primary health care units as part of standard care. Individuals received written information on the FIN-D2D and their participation in the project within normal primary care activities.

The present study analysed the baseline data from the FIN-D2D intervention, collected in 400 primary health care centres and occupational health clinics between 2004 and 2008. Altogether, 10 149 individuals aged 18–87 participated in the baseline assessments. Out of these, 9 984 individuals finally met the criteria for being at high risk. Glucose tolerance [49] was assessed with an oral glucose tolerance test (OGTT) among 8 353 of the participants. Individuals with previously-diagnosed type 2 diabetes, and screen-detected type 2 diabetes were excluded from the analysis, leaving 7 128 individuals in the analysis. The participant characteristics are presented in Table 1.

Measurements

The participants completed a questionnaire (questionnaire 1, issues regarding information on e.g. health behaviour and health status) and underwent health examinations. After these assessments the participants discussed with a nurse their health examination results, current health status, and health behaviour, in relation to current health recommendations and the risk of diabetes.

Perceived need to increase PAL

During the nurse-participant discussion, the nurse asked participants whether they perceived a need to change their health behaviours (to increase PAL, make changes in diet, quit smoking, reduce alcohol consumption, or lose weight in general). The nurses recorded the participants' behaviour change targets within the structured questionnaire (response options "no need perceived"/"need perceived", for each item separately) according to the participant's perception (questionnaire 2). The evaluation form (based on TTM and MI) was developed especially for the FIN-D2D intervention [46].

Demographic factors

The participants were classified into four age-group categories (see Table 1). The highest level of education was asked, with subsequent regrouping into three educational levels. The occupational status category originally included seven response-options, but these were subsequently regrouped into four categories. Marital status was also assessed.

Anthropometric and clinical risk factors

A family history of diabetes was assessed via a self-report. If at least one of the first-degree relatives (father, mother, or sibling) had diabetes (type 1 or type 2), family history was considered to be positive. Height and weight (usual light clothing, no shoes) were measured by the nurse for calculation of BMI (kg/m^2). The participants were classified into two groups by BMI ($<30 \text{ kg}/\text{m}^2$ and $\geq 30 \text{ kg}/\text{m}^2$). Waist circumference was measured to the nearest centimetre and classified into three groups: normal (men $<94 \text{ cm}$, women $<80 \text{ cm}$), elevated risk (men $94\text{--}101 \text{ cm}$, women $80\text{--}87 \text{ cm}$), and high risk (men $\geq 102 \text{ cm}$, women $\geq 88 \text{ cm}$). Blood pressure (mmHg, to the nearest 1mmHg) was measured twice from the right arm in sitting position (with at least a 1-min interval). The mean reading was recorded. Blood pressure was classified into two groups: normal, and elevated (systolic $\geq 130 \text{ mmHg}$ and/or diastolic $\geq 85 \text{ mmHg}$) blood pressure. Plasma lipids and lipoproteins were determined locally from fasting venous plasma samples using enzymatic methods. Participants with impaired values of HDL ($<1.03 \text{ mmol}/\text{l}$ in men, $<1.29 \text{ mmol}/\text{l}$ in women), LDL ($\geq 3 \text{ mmol}/\text{l}$), or triglycerides ($\geq 1.7 \text{ mmol}/\text{l}$), or medication for these lipid abnormalities were combined into a group called “dyslipidaemia”. The cut-off points used in the analysis were all based on international definitions [50,51].

Behavioural factors

Self-reported PAL was assessed with following questions (i) “How many times a week do you engage in leisure-time physical exercise causing you at least moderate sweating or breathlessness?” (ii) “For how long do you usually engage in leisure-time physical exercise at a time?”. The respondents were also asked to describe with structured descriptions their leisure-time physical activity during a typical week. Here the response options ranged from inactivity through light physical activity, and further to moderate/vigorous physical activity. The participants were classified as active if they did at least 30 minutes of physical exercise three times a week, and if they described the intensity of their usual leisure physical activity as at least moderate. All other participants were classified as inactive. Current smoking

(including regular and occasional smoking) was assessed and dichotomised (*yes/no*).

Psychosocial factors

Participants rated their physical fitness via five response options ranging from *very high* to *very low*. Extremes were combined with the nearest options. Participants also rated their readiness to increase their PAL level via a 5-item question. Here the response options covered stages of change from the precontemplation to the maintenance stage. Furthermore, participants rated their current PAL as sufficient or as insufficient for maintaining their health and physical fitness (*yes/no*).

Physical activity awareness

PAL awareness was assessed by comparing the participants' self-perceived sufficiency of their PAL with their self-reported PAL. Participants were classified into four physical activity awareness categories; *realistic active* (active by self-reported PAL, and perceiving their own PAL as sufficient), *overestimators* (inactive by self-report, but perceiving their own PAL as sufficient), *realistic inactive* (inactive by self-report, and perceiving their own PAL as insufficient) and *underestimators* (active by self-report, but perceiving their own PAL as insufficient).

Statistical methods

SPSS for Windows (14.1) was used for the statistical analysis. Descriptive statistics were used to describe the sample. Pearson's Chi Square tests were used to analyse differences between the groups and for a preliminary assessment of the associated factors. A multivariable logistic regression model was used to evaluate the association between a perceived need to increase PAL and demographic factors, anthropometric and clinical risk factors, behavioural factors, and psychosocial factors (Table 2). The results are presented as adjusted odds ratios (OR) and 95% confidence intervals (CI). Medication for a risk condition was taken as equivalent to having a risk condition. The category of lowest risk with regard to type 2 diabetes was used as a reference group for all variables. A separate logistic regression was used to determine the associations of physical activity awareness groups with the perceived need to increase PAL (Table 3).

Ethical considerations

The Ministry of Social Affairs and Health in Finland gave permission to the National Institute for Health and Welfare (formerly National Public Health Institute) to collect the data from health care units for evaluation purposes. In addition, the Institutional Review Board (IRB) of the National Institute for Health and Welfare approved the study. As lifestyle interventions were conducted as part of normal routine in primary health care

Table 2 Multivariable logistic regression model for perceived need to increase physical activity levels (PAL) among adults at high risk of diabetes (n = 7 128) by selected variables

	Men			Women		
	OR	(95% CI)	p	OR	(95% CI)	p
Age (years)						
<45	1.00			1.00		
45-54	1.05	(0.65-1.66)	0.892	0.79	(0.55-1.15)	0.213
55-64	0.68	(0.41-1.12)	0.132	0.76	(0.51-1.14)	0.187
≥65	0.90	(0.51-1.58)	0.714	0.69	(0.44-1.08)	0.101
Marital status						
Married/cohabiting	1.00			1.00		
Other	0.95	(0.68-1.32)	0.771	0.95	(0.75-1.20)	0.665
Educational level						
High	1.00			1.00		
Intermediate	0.63	(0.35-1.13)	0.118	1.46	(1.01-2.09)	0.042
Low	0.60	(0.33-1.09)	0.092	1.49	(1.01-2.19)	0.043
Occupational status						
Non-manual work	1.00			1.00		
Manual work	0.91	(0.62-1.33)	0.628	0.78	(0.54-1.15)	0.209
Retired	0.70	(0.45-1.10)	0.119	0.84	(0.62-1.16)	0.295
Not employed	0.75	(0.42-1.35)	0.343	0.90	(0.64-1.27)	0.543
Family history of diabetes ¹						
No	1.00			1.00		
Yes	0.98	(0.75-1.28)	0.865	1.09	(0.88-1.36)	0.422
Elevated blood pressure ²						
No	1.00			1.00		
Yes	0.95	(0.62-1.46)	0.821	0.84	(0.64-1.10)	0.208
Dyslipidaemia ³						
No	1.00			1.00		
Yes	1.14	(0.85-1.55)	0.384	1.11	(0.90-1.38)	0.324
Body mass index (kg/m ²)						
<30	1.00			1.00		
≥30	1.31	(0.94-1.83)	0.117	1.21	(0.95-1.55)	0.117
Waist circumference ⁴						
Normal	1.00			1.00		
Elevated	1.34	(0.80-2.25)	0.273	1.37	(0.73-2.56)	0.322
High risk	1.83	(1.07-3.13)	0.028	2.72	(1.50-4.93)	<0.001
Smoking						
No	1.00			1.00		
Yes ⁵	1.05	(0.75-1.47)	0.783	0.77	(0.58-1.04)	0.084
Physically active ⁶						
Yes	1.00			1.00		
No	1.46	(1.07-1.98)	0.017	1.57	(1.57-1.98)	<0.001
Self-rated fitness						
High	1.00			1.00		
Satisfactory	1.04	(0.75-1.42)	0.833	1.30	(1.02-1.66)	0.034

Table 2 Multivariable logistic regression model for perceived need to increase physical activity levels (PAL) among adults at high risk of diabetes (n = 7 128) by selected variables (Continued)

Low	1.42	(0.90-2.22)	0.128	1.65	(1.18-2.33)	0.004
Stage of change						
Maintenance	1.00			1.00		
Action	2.17	(1.25-3.78)	0.006	2.29	(1.54-2.41)	<0.001
Preparation	2.54	(1.51-4.28)	<0.001	2.24	(1.56-3.23)	<0.001
Contemplation	2.68	(1.55-4.65)	<0.001	2.91	(1.90-4.46)	<0.001
Precontemplation	0.95	(0.53-1.69)	0.856	1.42	(0.87-2.31)	0.163
PAL perceived as sufficient						
Yes	1.00			1.00		
No	2.40	(1.72-3.37)	<0.001	2.95	(2.27-3.83)	<0.001

Adjusted odds ratios (OR), and their 95% confidence intervals (CI), and *p*-values

¹ At least one of the first-degree relatives had diabetes (type 1 or 2).

² Systolic blood pressure ≥ 130 mmHg and/or diastolic blood pressure ≥ 85 mmHg.

³ HDL < 1.03 mmol/l (men), < 1.29 mmol/l (women); LDL ≥ 3 mmol/l or/and triglycerides ≥ 1.7 mmol/l, or medication for these lipid abnormalities.

⁴ Normal: < 94 cm (m), < 80 cm (w); elevated risk: 94-101 cm (m), 80-87 cm (w); high risk: ≥ 102 cm (m), ≥ 88 cm (w).

⁵ Includes occasional and regular smokers.

⁶ Engaging in physical exercise < 30 minutes three times a week.

units, informed consent was not collected from the participants; however, participants did receive written information on the FIN-D2D. The FIN-D2D project was not a scientific study under the legislation of the Medical Research Act in Finland, but rather an *implementation study within routine primary care in Finland*. Therefore, it would not in fact have been possible to collect statements of consent. Furthermore, the FIN-D2D data collection system was established by the participating hospital districts as part of normal patient records within primary health care.

Results

The mean age of the participants was 55.4 (SD ± 10.23) years, and the mean BMI 31.3 kg/m² (SD ± 4.7 kg/m²) in men, and 32.1 (SD ± 5.4 kg/m²) in women. The majority (64%) of the 7 128 participants were women. The basic characteristics of the participants are presented in Table 1. Smoking and elevated blood pressure were significantly more prevalent among men, whereas a family history of diabetes, dyslipidaemia, and obesity were all more prevalent among women. In total, 65% of the participants were classified as physically inactive. Women rated their fitness as low more often than men. However, compared to women, men were significantly more often (40% for men vs. 31% for women, $p < 0.001$) in the early stages of change (precontemplation and contemplation), and men also perceived their PAL as sufficient significantly more often than women (33% for men vs. 30% for women, $p < 0.05$).

In total, 74% of the men and 76% of the women perceived a need to increase their PAL with no significant difference between the sexes (Table 1). As shown in Table 2,

a lower education level increased the likelihood of perceiving the need to increase PAL among women (intermediate education level: OR 1.46 [95% CI 1.01-2.09]; low education level: OR 1.49 [95% CI 1.01-2.19]). The opposite association appeared to exist among men, but did not reach statistical significance. Other demographic factors did not make any contribution. The need to increase PAL was significantly more often perceived by men and women with high-risk waist circumference than by those with normal waist circumference. The other anthropometric or clinical risk factors did not emerge as significant determinants. Physically inactive men and women were more likely than physically active participants to perceive the need to increase their PAL. In women, lower self-rated fitness also increased the likelihood of perceiving the need to increase PAL. Furthermore, men and women at the contemplation, preparation and action stages – but not at the

Table 3 Perceived need to increase physical activity levels (PAL) by physical activity awareness categories, odds ratios (OR), 95% confidence intervals (CI), and *p*-values by sex

	Men			Women		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Realistic active ¹	1.00			1.00		
Overestimators ²	1.51	(1.06-2.17)	0.024	1.79	(1.35-2.37)	<0.001
Realistic inactive ³	7.14	(5.38-9.48)	<0.001	7.86	(6.37-9.70)	<0.001
Underestimators ⁴	4.23	(2.92-6.14)	<0.001	4.68	(3.57-6.13)	<0.001

¹ Defined as physically active (≥ 30 min 3 times a week) and perceiving PAL as sufficient.

² Defined as physically inactive (< 30 min 3 times a week), but perceiving PAL as sufficient.

³ Defined as inactive and perceiving PAL as insufficient.

⁴ Defined as active, but perceiving PAL as insufficient.

precontemplation stage – were significantly more likely to perceive the need to increase their PAL than those at the maintenance stage of change; the perceived need to increase PAL was most likely among the participants at the contemplation stage and least likely among the participants at the precontemplation stage. The need to increase PAL was perceived more often by participants (both men and women) who rated their PAL as insufficient than by those who perceived their PAL as sufficient.

Out of the total population, 21% of men and 20% of women were classified as realistic active, 52% men and 55% women as realistic inactive, 13% men and 10% women as overestimators, and 14% men and 15% women as underestimators (not presented in the tables). In comparison with the realistic active participants, with regard to the three other awareness groups, the realistic inactive participants were the most likely to perceive the need to increase their PAL (Table 3). Overestimators had a significantly lower likelihood of perceiving a need to increase their PAL than realistic inactive participants.

Discussion

The vast majority of the study participants (74% of men and 76% of women) perceived a need to increase their PAL. This reflects the high value placed on physical activity among the type 2 diabetes at-risk population, as indicated also by previous studies [12,13]. Considering the importance of increased physical activity in diabetes prevention, and the high prevalence of inactivity among the study population, the finding is promising, bearing in mind also that expression of a perceived need to change is a key component of a client's "change talk", predicting a commitment to change health behaviour [27,28]. Through professional advice [52] and effective counselling techniques e.g. [53] those perceiving the need to increase PAL could be encouraged to make a genuine commitment to changing their behaviour.

The present study showed that people at the contemplation stage of change, who perceived their PAL as sufficient, and who were classified as physically inactive, were more likely than others to perceive a need to increase their PAL. In addition, women who gave lower ratings to their personal physical fitness were more likely than others to perceive this need. A previous study [22] found general health and well-being, physical fitness, and weight control to be the most important determinants of a perceived need to exercise. Along similar lines, the present paper indicated a significant association between increased waist circumference and a perceived need to increase PAL in both sexes. By contrast, other type 2 diabetes risk factors did not contribute to this perception. These findings indicate the groups which are least likely to increase their PAL, and which should therefore be given particular attention within health care, in terms of endeavours towards

diabetes prevention. Increased PAL would be of great benefit to individuals with a family history of diabetes, elevated blood pressure, or dyslipidaemia [8], yet the benefit does not appear to be recognised by these people. Waist circumference is an objective measure which is highly tangible for the individual, and which inevitably reflects the person's current PAL in a manner readily apparent to that individual.

Our results suggest that among the high-risk population the need to increase PAL is strongly determined by the individual's subjective perceptions. This finding is in line with the principles of the motivational interviewing approach to counselling, which emphasise the importance of listening to and reflecting individuals' perceptions if one is to evoke the motivation to change. However, clients' perceptions may easily be overlooked in a busy primary health care practice, especially if health professionals view the client as reluctant to change [54]. We would argue that if the perceptions of clients are neglected in health counselling, an important opportunity to support the client's motivation for behaviour change will remain unutilised.

Our results support previous findings on PAL awareness indicating that overestimation of one's PAL may be an obstacle to behaviour change [37,39-43]. In demonstrating a lack of any intention to change, the overestimators differed only slightly from realistic active persons, but there was a large difference between the overestimators and the realistic inactive persons. If the overestimators could be led to view their PAL realistically, they could be expected to see more clearly the need for change. Such efforts could also help to prevent incorrect tailoring in physical activity counselling [55].

As far as we know, no other studies have so far simultaneously assessed such a variety of determinants of the perceived need to increase PAL. One could expect that factors associated with a failure to perceive a need for increased PAL would be in line with determinants of the pre-contemplation stage of change. However, we found no evidence to support this presumption [37,38,56]. It appears that the pre-contemplation stage can best be regarded as a complex entity with its own set of determinants. It seems to be the case that some individuals at the precontemplation stage simply do not perceive the need to change their behaviours [57]. However, others perceive the need to increase their PAL in principle, yet – for whatever reason – do not consider changing their behaviour.

A previous study by Payne et al. [22] conducted among the general population (n = 286) found that 94% of the participants perceived the need to exercise (note that the authors did not assess the need to *increase exercise*, merely the need to *exercise* in general). It is true that the authors did not find a significant association between the perceived need and the intention or behaviour; yet the concept of

perceived need requires further examination. Payne et al. assessed short-range intention only (the intention to exercise next week, and actual exercise behaviour a week later). Here it is worth bearing in mind that e.g. in the Diabetes Prevention Programme (DPP) [12], those who were at the contemplation stage at the baseline actually continued to increase their PAL until the 3-year (final) follow-up assessments; by contrast, other groups decreased their PAL. In fact, even six months can be regarded as a short period for changing physical activity behaviour [58-60]. The causal relationships between the perceived need to change, intentions, and behaviour should be assessed in future studies.

The study has some limitations. The study population is considered to represent the Finnish high-risk population [61]. Nevertheless, as opportunistic screening was used in the programme, there is a possibility of selection bias. The study participants were aware of their risk due to health examinations, and this aspect limits comparisons with the unaware high-risk population outside the programme. Furthermore, the assessment of the perceived need to change health behaviours was based on an evaluation form developed for the FIN-D2D intervention. The nurses were instructed to record not their own, but merely the participant's perception; however, it is possible that a persuasive style on the part of the nurse could have influenced the responses (though attempts were made to avoid such a bias through careful instruction of the nurses). Furthermore, the self-reporting method of measuring physical activity limits the assessment of PAL and PAL awareness, due to issues of recall and response bias (e.g. social desirability, inaccurate memory) and the inability of the respondent to estimate the frequency, intensity, and duration of physical activity [62]. It was due to these considerations that we did not assess lifestyle physical activity, which is often sporadic, and even more difficult to recall and report than planned, discretionary, leisure-time exercise [63]. Here it should be noted that with regard to cost, staff training, participant burden, and time, self-reporting was the only feasible method for measuring PAL in such a large implementation programme [64]. It should also be noted that the cut-off point for physically active and inactive participants was not determined according to the CDC/ACSM physical activity recommendations [65], due to the limitations of the questionnaire. However, it has been suggested [7] that the amount of physical activity needed to prevent diabetes should in fact be set at lower than the recommendations for the general adult population given by CDC/ACSM.

Conclusions

The present study shows that the vast majority of high risk individuals perceive the need to increase their PAL. However, apart from waist circumference, the perception

of need appears not to be determined by the objective diabetes risk factors. Instead, the perception appears to be strongly determined by factors that are based on individuals' subjective impressions. This finding, taken together with the association between physical activity awareness and a perceived need to increase PAL, highlights the importance of listening to and reflecting clients' perceptions (which may not always be realistic), rather than focusing mainly on objective health measures. It is further important to increase clients' awareness by *concretely* defining the association of physical activity with all the metabolic risk factors – and with the development of diabetes. This study did not assess the predictors of behaviour change, but it does demonstrate factors worth recognising in future interventions whose aim is to facilitate the intention and commitment to change physical activity behaviour among individuals at high risk of type 2 diabetes.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KV conceived of and designed the study, interpreted the data and drafted the manuscript. SS, JV, MV, PR and MEP participated in the design of the study. JV processed the data and performed the statistical analyses. All authors participated in interpretation of the results and critically revised subsequent versions of the paper for important intellectual content. TS, MV, MP, LN, LM, EK-H, SK-K, HO, and MEP were all in charge of the FIN-D2D concept and design, and the acquisition of the data. All the authors read and approved the final manuscript.

Acknowledgements

This work was supported by the Ministry of Social Affairs and Health; the Social Insurance Institution of Finland; the Diabetes Research Foundation of Finland; the Ministry of Education and Culture [grant numbers 121/722/2004, 45/627/2007 to KV] and the Juho Vainio Foundation [to KV]. FIN-D2D was supported by hospital districts of Pirkanmaa, Southern Ostrobothnia, North Ostrobothnia, Central Finland and Northern Savo; the Finnish National Public Health Institute; the Finnish Diabetes Association; the Ministry of Social Affairs and Health in Finland, and Finland's Slot Machine Association, in cooperation with the FIN-D2D Study Group, and the Steering Committee (consisting of Huttunen J, Kesäniemi A, Kiuru S, Niskanen L, Oksa H, Pihlajamäki J, Puolakka J, Puska P, Saaristo T, Vanhala M, and Uusitupa M); the Academy of Finland [129293]; and the Commission of the European Communities, Directorate C – Public Health [2004310]. We acknowledge the help of Donald Adamson (University of Jyväskylä) in editing the manuscript.

Author details

¹Department of Health Sciences, University of Jyväskylä, Jyväskylä 40014, Finland. ²Department of Sport Sciences, University of Jyväskylä, Jyväskylä 40014, Finland. ³Family Practice Unit, Central Hospital Central Finland, Jyväskylä 40620, Finland. ⁴Family Practice Unit, Kuopio University Hospital, Kuopio 70210, Finland. ⁵University of Eastern Finland, Kuopio 70211, Finland. ⁶Finnish Diabetes Association, Tampere 33680, Finland. ⁷Pirkanmaa Hospital District, Tampere 33680, Finland. ⁸National Institute for Health and Welfare, Helsinki 00271, Finland. ⁹Institute of Health Sciences, University of Oulu, Oulu 90014, Finland. ¹⁰Unit of General Practice, Oulu University Hospital, Oulu 90220, Finland. ¹¹Health Centre of Oulu, Oulu 90100, Finland. ¹²Department of Internal Medicine, Seinäjoki Central Hospital, Seinäjoki 60220, Finland. ¹³Department of Medicine, Kuopio University Hospital, Kuopio 70210, Finland. ¹⁴Department of Medicine, Central Hospital Central Finland, Jyväskylä 40014, Finland. ¹⁵University of Eastern Finland, Kuopio 70211, Finland. ¹⁶Tampere University Hospital, Tampere 33521, Finland.

Received: 14 December 2011 Accepted: 21 June 2012
Published: 10 July 2012

References

1. Laaksonen DE, Lindstrom J, Lakka TA, Eriksson JG, Niskanen L, Wilkstrom K, Aunola S, Keinanen-Kiukkaanniemi S, Laakso M, Valle TT, Ilanne-Parikka P, Louheranta A, Hamalainen H, Rastas M, Salminen V, Cepaitis Z, Hakumaki M, Kaikkonen H, Harkonen P, Sundvall J, Tuomilehto J, Uusitupa M, Finnish diabetes prevention study: **Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study.** *Diabetes* 2005, **54**:158–165.
2. Bassuk SS, Manson JE: **Epidemiological evidence for the role of physical activity in reducing risk of type 2 diabetes and cardiovascular disease.** *J Appl Physiol* 2005, **99**:1193–1204.
3. Warburton DE, Nicol CW, Bredin SS: **Health benefits of physical activity: the evidence.** *CMAJ* 2006, **174**:801–809.
4. Hu G, Qiao Q, Silventoinen K, Eriksson JG, Jousilahti P, Lindstrom J, Valle TT, Nissinen A, Tuomilehto J: **Occupational, commuting, and leisure-time physical activity in relation to risk for Type 2 diabetes in middle-aged Finnish men and women.** *Diabetologia* 2003, **46**:322–329.
5. Borodulin K, Tuomilehto J, Peltonen M, Lakka TA, Sundvall J, Jousilahti P: **Association of leisure time physical activity and abdominal obesity with fasting serum insulin and 2-h postchallenge plasma glucose levels.** *Diabet Med* 2006, **23**:1025–1028.
6. Healy GN, Dunstan DW, Salmon J, Cerin E, Shaw JE, Zimmet PZ, Owen N: **Objectively measured light-intensity physical activity is independently associated with 2-h plasma glucose.** *Diabetes Care* 2007, **30**:1384–1389.
7. Waller K, Kaprio J, Lehtovirta M, Silventoinen K, Koskenvuo M, Kujala UM: **Leisure-time physical activity and type 2 diabetes during a 28 year follow-up in twins.** *Diabetologia* 2010, **53**:2531–2537.
8. LaMonte MJ, Blair SN, Church TS: **Physical activity and diabetes prevention.** *J Appl Physiol* 2005, **99**:1205–1213.
9. Boyle RG, O'Connor PJ, Pronk NP, Tan A: **Stages of change for physical activity, diet, and smoking among HMO members with chronic conditions.** *Am J Health Promot* 1998, **12**:170–175.
10. Brewer NT, Chapman GB, Gibbons FX, Gerard M, McCaul KD, Weinstein ND: **Meta-analysis of the relationship between risk perception and health behavior: The example of vaccination.** *Health Psychol* 2007, **26**:136–145.
11. Kim C, Mc Ewen LN, Piette JD, Goewey J, Ferrara A, Walker E: **Risk perception for diabetes among women with histories of gestational diabetes mellitus.** *Diabetes Care* 2007, **30**:2281–2286.
12. Delahanty LM, Conroy MB, Nathan DM, Diabetes Prevention Program Research Group: **Psychological predictors of physical activity in the diabetes prevention program.** *J Am Diet Assoc* 2006, **106**:698–705.
13. Centers for Disease Control and Prevention (CDC): **Self-reported prediabetes and risk-reduction activities—United States, 2006.** *MMWR Morb Mortal Wkly Rep* 2008, **57**:1203–1205.
14. Blue CL: **Does the theory of planned behavior identify diabetes-related cognitions for intention to be physically active and eat a healthy diet?** *Public Health Nurs* 2007, **24**:141–150.
15. Hivert MF, Warner AS, Shrader P, Grant RW, Meigs JB: **Diabetes risk perception and intention to adopt healthy lifestyles among primary care patients.** *Diabetes Care* 2009, **32**:1820–1822.
16. Prochaska JO, Norcross JC: **Systems of psychotherapy. A Transtheoretical analysis.** 7th edition. Belmont, CA: Brooks/Cole; 2010.
17. Miller WR, Rose GS: **Toward a theory of motivational interviewing.** *Am Psychol* 2009, **64**:527–537.
18. Hettema J, Steele J, Miller WR: **Motivational interviewing.** *Annu Rev Clin Psychol* 2005, **1**:91–111.
19. Paisley CM, Sparks P: **Expectations of reducing fat intake: the role of perceived need within the theory of planned behaviour.** *Psychol Health* 1998, **13**:341–353.
20. Povey R, Conner M, Sparks P, James R, Shepherd R: **Application of the theory of planned behaviour to two dietary behaviours: roles of perceived control and self-efficacy.** *Br J Health Psychol* 2000, **5**:121–139.
21. Raats MM, Sparks P, James R, Shepherd R: **Understanding dietary change: Perceptions of ten dietary changes.** *Proc Nutr Soc* 1996, **55**:1A–77A.
22. Payne N, Jones F, Harris PR: **The role of perceived need within the theory of planned behaviour: a comparison of exercise and healthy eating.** *Br J Health Psychol* 2004, **9**(Pt 4):489–504.
23. Ajzen I: **The theory of planned behaviour.** *Organ Behav Hum Decis Process* 1991, **50**:179–211.
24. Prochaska JO, DiClemente CC: **Stages and processes of self-change of smoking: toward an integrative model of change.** *J Consult Clin Psychol* 1983, **51**:390–395.
25. Miller WR, Rollnick S: **Motivational interviewing: preparing people for change.** 2nd edition. New York: Guilford Press; 2002.
26. Amrhein PC, Miller WR, Yahne CE, Palmer M, Fulcher L: **Client commitment language during motivational interviewing predicts drug use outcomes.** *J Consult Clin Psychol* 2003, **71**:862–878.
27. Moyers TB, Martin T, Christopher PJ, Houck JM, Tonigan JS, Amrhein PC: **Client language as a mediator of motivational interviewing efficacy: where is the evidence?** *Alcohol Clin Exp Res* 2007, **31**(10 Suppl):405–475.
28. Apodaca TR, Longabaugh R: **Mechanisms of change in motivational interviewing: a review and preliminary evaluation of the evidence.** *Addiction* 2009, **104**:705–715.
29. Hardcastle S, Taylor A, Bailey M, Castle R: **A randomised controlled trial on the effectiveness of a primary health care based counselling intervention on physical activity, diet and CHD risk factors.** *Patient Educ Couns* 2008, **70**:31–39.
30. Bennett JA, Lyons KS, Winters-Stone K, Nail LM, Scherer J: **Motivational interviewing to increase physical activity in long-term cancer survivors: a randomized controlled trial.** *Nurs Res* 2007, **56**:18–27.
31. Brodie DA, Inoue A: **Motivational interviewing to promote physical activity for people with chronic heart failure.** *J Adv Nurs* 2005, **50**:518–527.
32. Harland J, White M, Drinkwater C, Chinn D, Farr L, Howel D: **The Newcastle exercise project: a randomised controlled trial of methods to promote physical activity in primary care.** *Br Med J* 1999, **319**:828–832.
33. Hutchison AJ, Breckon JD, Johnston LH: **Physical activity behavior change interventions based on the transtheoretical model: a systematic review.** *Health Educ Behav* 2009, **36**:829–845.
34. Spencer L, Adams TB, Malone S, Roy L, Yost E: **Applying the transtheoretical model to exercise: a systematic and comprehensive review of the literature.** *Health Promot Pract* 2006, **7**:428–443.
35. Carino JL, Coke L, Gulanic M: **Using motivational interviewing to reduce diabetes risk.** *Prog Cardiovasc Nurs* 2004, **19**:149–154.
36. Penn L, White M, Oldroyd J, Walker M, Alberti KG, Mathers JC: **Prevention of type 2 diabetes in adults with impaired glucose tolerance: the European Diabetes Prevention RCT in Newcastle upon Tyne, UK.** *BMC Public Health* 2009, **9**:342.
37. Ronda G, Van Assema P, Brug J: **Stages of change, psychological factors and awareness of physical activity levels in The Netherlands.** *Health Promot Int* 2001, **16**:305–314.
38. Garber CE, Allsworth JE, Marcus BH, Hesser J, Lapane KL: **Correlates of the stages of change for physical activity in a population survey.** *Am J Public Health* 2008, **98**:897–904.
39. De Ridder C, Lechner L: **Overschatting van het eigen bewegingsgedrag: gevolgen voor determinanten en stage of change.** *Tijdschrift Sociale Gezondheidszorg* 2004, **79**:346–355.
40. van Sluijs EM, Griffin SJ, van Poppel MN: **A cross-sectional study of awareness of physical activity: associations with personal, behavioral and psychosocial factors.** *Int J Behav Nutr Phys Act* 2007, **4**:53.
41. Watkinson C, van Sluijs EM, Sutton S, Hardeman W, Corder K, Griffin SJ: **Overestimation of physical activity level is associated with lower BMI: a cross-sectional analysis.** *Int J Behav Nutr Phys Act* 2010, **7**:68.
42. Lechner L, Bolman C, Van Dijke M: **Factors related to misperception of physical activity in The Netherlands and implications for health promotion programmes.** *Health Promot Int* 2006, **21**:104–112.
43. Corder K, van Sluijs EM, McMinn AM, Ekelund U, Cassidy A, Griffin SJ: **Perception versus reality awareness of physical activity levels of British children.** *Am J Prev Med* 2010, **38**:1–8.
44. Finnish Diabetes Association: **Programme for the Prevention of Type 2 Diabetes in Finland 2003–2010.** Gummerus Printing, Jyväskylä: Finnish Diabetes Association; 2003.
45. Finnish Diabetes Association: **Dehkon 2D-hanke (D2D) 2003–2007. Loppuraportti.** In *Finnish*. Kirjapaino Hermes Oy: Finnish Diabetes Association; 2009.
46. Saariisto T, Peltonen M, Keinanen-Kiukkaanniemi S, Vanhala M, Saltevo J, Niskanen L, Oksa H, Korpi-Hyövälti E, Tuomilehto J, FIN-D2D Study Group: **National type 2 diabetes prevention programme in Finland: FIN-D2D.** *Int J Circumpolar Health* 2007, **66**:101–112.

47. Saaristo T, Moilanen L, Korpi-Hyovalti E, Vanhala M, Saltevo J, Niskanen L, Jokelainen J, Peltonen M, Oksa H, Tuomilehto J, Uusitupa M, Keinänen-Kiukaanniemi S: **Lifestyle intervention for prevention of type 2 diabetes in primary health care: one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D).** *Diabetes Care* 2010, **33**:2146–2151.
48. Lindstrom J, Tuomilehto J: **The diabetes risk score: a practical tool to predict type 2 diabetes risk.** *Diabetes Care* 2003, **26**:725–731.
49. World Health Organisation: *Definition, Diagnosis and Classification of Diabetes Mellitus and Its complications. Report of a Who Consultation. Part 1: Diagnosis and Classification of Diabetes Mellitus*; 1999. 99.2.
50. Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, Dallongeville J, De Backer G, Ebrahim S, Gjelsvik B, Herrmann-Lingen C, Hoes A, Humphries S, Knapp M, Perk J, Priori SG, Pyörälä K, Reiner Z, Ruilope L, Sans-Menendez S, Op Reimer WS, Weissberg P, Wood D, Yarnell J, Zamorano JL, Walma E, Fitzgerald T, Cooney MT, Dudina A, Vahanian A, et al: **European guidelines on cardiovascular disease prevention in clinical practice: full text. Fourth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts).** *Eur J Cardiovasc Prev Rehabil* 2007, **14**(Suppl 2):S1–S113.
51. Alberti KG, Zimmet P, Shaw J: **International Diabetes Federation: a consensus on Type 2 diabetes prevention.** *Diabet Med* 2007, **24**:451–463.
52. Geiss LS, James C, Gregg EW, Albright A, Williamson DF, Cowie CC: **Diabetes risk reduction behaviors among U.S. adults with prediabetes.** *Am J Prev Med* 2010, **38**:403–409.
53. Williams SL, French DP: **What are the most effective intervention techniques for changing physical activity self-efficacy and physical activity behaviour—and are they the same?** *Health Educ Res* 2011, **26**:308–322.
54. Jallinoja P, Absetz P, Kuronen R, Nissinen A, Talja M, Uutela A, Patja K: **The dilemma of patient responsibility for lifestyle change: perceptions among primary care physicians and nurses.** *Scand J Prim Health Care* 2007, **25**:244–249.
55. Noar SM, Benac CN, Harris MS: **Does tailoring matter? Meta-analytic review of tailored print health behavior change interventions.** *Psychol Bull* 2007, **133**:673–693.
56. Lorentzen C, Ommundsen Y, Holme I: **Psychosocial correlates of stages of change in physical activity in an adult community sample DOI:dx.doi.org.** *Eur J Sport Sci* 2007, **7**:93–106.
57. Miilunpalo S, Nupponen R, Laitakari J, Marttila J, Parosnen O: **Stages of change in two modes of health-enhancing physical activity: methodological aspects and promotional implications.** *Health Educ Res* 2000, **15**:435–448.
58. Marcus BH, Forsyth LH, Stone EJ, Dubbert PM, McKenzie TL, Dunn AL, Blair SN: **Physical activity behaviour change: issues in adoption and maintenance.** *Health Psychol* 2000, **19**(Suppl 1):32–41.
59. Marcus BH, Williams DM, Dubbert PM, Sallis JF, King AC, Yancey AK, Franklin BA, Buchner D, Daniels SR, Claytor RP, American Heart Association Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), American Heart Association Council on Cardiovascular Disease in the Young, Interdisciplinary Working Group on Quality of Care and Outcomes Research: **Physical activity intervention studies: what we know and what we need to know: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity); Council on Cardiovascular Disease in the Young; and the Interdisciplinary Working Group on Quality of Care and Outcomes Research.** *Circulation* 2006, **114**:2739–2752.
60. Foster C, Hillsdon M, Thorogood M: **Interventions for promoting physical activity.** *Cochrane Database Syst Rev* 2009, **1**. doi:10.1002/14651858.CD003180.pub2.
61. Saaristo T: *Assessment of risk and prevention of type 2 diabetes in primary health care.* 2011. PhD thesis. Acta Universitatis Ouluensis D1144, University of Oulu.
62. Shephard RJ: **Limits to the measurement of habitual physical activity by questionnaires.** *Br J Sports Med* 2003, **37**:197–206. discussion 206.
63. Troiano RP, Pette Gabriel KK, Welk GJ, Owen N, Sternfeld B: **Reported physical activity and sedentary behaviour: why do you ask?** *J Phys Act Health* 2012, **9**(Suppl 1):S68–S72.
64. Hooker SP, Fulton J, Mudd LM: **Practice-based evidence: A novel concept for self-report physical activity measurement.** *J Phys Act Health* 2012, **9**(Suppl 1):S85–S87.
65. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, Macera CA, Heath W, Thompson PD, Bauman A: **Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association.** *Med Sci Sports Exerc* 2007, **39**:1423–1434.

doi:10.1186/1471-2458-12-514

Cite this article as: Vähäsarja et al.: Perceived need to increase physical activity levels among adults at high risk of type 2 diabetes. A cross-sectional analysis within a community-based diabetes prevention project FIN-D2D. *BMC Public Health* 2012 **12**:514.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

