

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Stenroth, Lauri; Cronin, Neil; Peltonen, Jussi; Korhonen, Marko; Sipilä, Sarianna; Finni Juutinen, Taija

Title: Triceps surae muscle-tendon properties in older endurance- and sprint-trained athletes

Year: 2016

Version:

Please cite the original version:

Stenroth, L., Cronin, N., Peltonen, J., Korhonen, M., Sipilä, S., & Finni Juutinen, T. (2016). Triceps surae muscle-tendon properties in older endurance- and sprint-trained athletes. *Journal of Applied Physiology*, 120(1), 63-69.
<https://doi.org/10.1152/jappphysiol.00511.2015>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 **Triceps surae muscle-tendon properties in older endurance- and sprint-**
2 **trained athletes**

3 Stenroth L.^{1,2}, Cronin N.J.¹, Peltonen J.¹, Korhonen M.T.², Sipilä S.² and Finni, T.¹

4 ¹University of Jyväskylä, Neuromuscular Research Center, Department of Biology of Physical
5 Activity, ²University of Jyväskylä, Gerontology Research Center and Department of Health
6 Sciences

7 Conception and design: LS, NJC, SS and TF; Performed experiments: LS; Analyzed data: LS
8 and JP; Interpreted results: LS and MTK; Prepared figures: LS; Drafted manuscript: LS; Edited
9 and revised manuscript: LS, NJC, JP, MTK, SS and TF; Approved final version: LS, NJC, JP,
10 MTK, SS and TF.

11 Running head: Muscle-tendon properties in older athletes

12 Corresponding author: Lauri Stenroth, University of Jyväskylä, Department of Biology of
13 Physical Activity, 40014 Jyväskylä, Finland. Email: lauri.stenroth@jyu.fi.

14 **Abstract**

15 Previous studies have shown that aging is associated with alterations in muscle architecture and
16 tendon properties. However, the possible influence of different types of regular exercise loading
17 on muscle architecture and tendon properties in older adults is poorly understood. To address
18 this, triceps surae muscle-tendon properties were examined in older male endurance (OE, n=10,
19 age=74.0±2.8) and sprint runners (OS, n=10, age=74.4±2.8) with an average of 42 years of
20 regular training experience, and compared to age-matched (OC, n=33, age=74.8±3.6) and young
21 untrained controls (YC, n=18, age=23.7±2.0).

22 Compared to YC, Achilles tendon cross-sectional area (CSA) was 22% (p=0.022), 45 %
23 (p=0.001) and 71% (p<0.001) larger in OC, OE and OS, respectively. Among older groups, OS
24 had significantly larger tendon CSA compared to OC (p=0.033). No significant between-group
25 differences were observed in Achilles tendon stiffness. In older groups, Young's modulus was
26 31-44% and maximal tendon stress 44-55% lower than in YC (p≤0.001). OE showed shorter
27 soleus fascicle length than both OC (p<0.05) and YC (p<0.05).

28 These data suggest that long-term running does not counteract the previously reported age-
29 related increase in tendon CSA, but instead, may have an additive effect. The greatest Achilles
30 tendon CSA was observed in sprinters followed by endurance runners and older controls,
31 suggesting that adaptation to running exercise is loading intensity dependent. Achilles tendon
32 stiffness was maintained in older groups even though all older groups displayed larger tendon
33 CSA and lower tendon Young's modulus. Shorter soleus muscle fascicles in older endurance
34 runners may be an adaptation to life-long endurance running.

35 **Keywords:** Achilles tendon, mechanical properties, muscle architecture, aging, exercise

36 **Introduction**

37 Loss of muscle function with aging is associated with physical limitations and disability (40).
38 Decline in muscle mass is undoubtedly an important contributor to the deterioration in muscle
39 function with aging (16). However, longitudinal studies have shown a clear dissociation in loss
40 of muscle function and cross-sectional area or mass with aging (9, 17), suggesting that other
41 factors may also contribute to the age-related loss of muscle function. Muscle architecture and
42 tendon mechanical properties greatly affect muscle performance (28, 51) and have been found to
43 differ between young and old sedentary adults (36, 37, 44). Thus, age-related alterations in
44 muscle architecture and tendon mechanical properties may partially explain the loss of muscle
45 performance with age that occurs at a disproportionately faster rate than the decline in muscle
46 mass.

47 Regular exercise is a key aspect supporting healthy aging. Indeed, it has been suggested that
48 older athletes provide a model of exceptionally successful biological aging (46). For example,
49 previous studies have shown that aged athletes with systematic exercise training habits exhibit
50 much better cardiorespiratory, metabolic and bone health than their less active counterparts (22,
51 49). Regular exercise training, especially strength and sprint exercise, also helps to maintain
52 muscle mass, function (21, 52) and composition (43), thus counteracting the age-related decline
53 in functional performance typically observed in normal populations (9, 39).

54 In spite of several known beneficial effects of regular exercise on the musculoskeletal system in
55 old age, little is known about the effects of regular participation in planned exercise on muscle
56 architecture and tendon properties in older adults. Two previous studies have compared
57 untrained older adults to older endurance runners. Firstly, Karamanidis and Arampatzis (19)

58 found that muscle architecture and tendon stiffness in medial gastrocnemius and vastus lateralis
59 were largely similar in older endurance runners compared to untrained older adults. The only
60 significant difference was greater medial gastrocnemius pennation angle in endurance runners.
61 Secondly, Couppé et al. (7) recently found that older endurance runners had a greater patella
62 tendon cross-sectional area but similar tendon stiffness compared to untrained peers. These
63 previous studies were conducted on endurance runners and thus knowledge of the long-term
64 effects of different types of exercise loading on muscle architecture and tendon properties in
65 older adults is missing.

66 Therefore, the aim of this study was to examine the association between different types of life-
67 long exercise and muscle-tendon properties by comparing muscle architecture and tendon
68 properties in older sprint and endurance runners to both age-matched and young untrained adults.
69 Triceps surae muscles were studied because of their important role in locomotion and because
70 they exhibit the greatest functional limitation of all lower limb muscle groups in older adults
71 during locomotion (24). Endurance running provides a model of high volume and moderate
72 intensity loading while sprint running represents a model of low volume but high intensity
73 loading of triceps surae muscles. The hypothesis was that older athletes with a life-long regular
74 running background would exhibit muscle fascicle length, pennation angle, muscle size, muscle
75 strength and tendon mechanical properties in the triceps surae muscle group that are more similar
76 to those of young adults than untrained older adults. In addition, based on previous cross-
77 sectional studies conducted in young adults (1, 2), it was hypothesized that sprint-trained older
78 athletes would be stronger, have stiffer Achilles tendons, lower pennation angle, and longer
79 muscle fascicles compared to endurance-trained older athletes.

80 **Materials and methods**

81 **Subjects**

82 Male subjects were recruited in two age categories, one from 18 to 30 years old (untrained young
83 controls, YC, n=18) and the other from 70 to 80 years old. The older cohort was recruited in
84 three groups: untrained older controls (OC, n=33), older athletes competing in endurance running
85 events (OE, n=10) and older athletes competing in sprint running events (OS, n=10).

86 Untrained young and older control groups were part of a Europe-wide collaborative study called
87 MyoAge (34) and included in the current study to represent general populations of healthy young
88 and older adults. We defined untrained as a person who may be recreationally active but is not
89 training for, or participating in, competitive sport. YC were recruited from among university
90 students using study advertisements via e-mail and bulletin boards. We excluded those who
91 studied sport sciences as well as competitive athletes. OC were recruited from the University of
92 the Third Age or from weekly community meetings of retired people. The aim was to recruit
93 healthy older people who were socially active and free from comorbidity. Using telephone
94 interviews, an equal number of sedentary and physically active (competitive athletes excluded)
95 older subjects were recruited to obtain a representative sample of older people with varying
96 physical activity levels. Sedentariness was defined as exercising for fitness and health one or
97 fewer times per week. Physically active was defined as exercise three or more times per week
98 (30 min or more with intensity sufficient to cause sweating or breathlessness). Results (44) and
99 more detailed description of the recruitment (34) of YC and OC have been presented earlier.

100 Older athletes were recruited among the participants of the World Master Athletics Indoor
101 Championships held in Jyväskylä, Finland in 2012. Twenty male athletes were recruited based

102 on the events they participated in during the championships. Ten subjects were recruited from
103 sprint running events (60 m, 60 m hurdles, 200 m, 400 m) and 10 were recruited from endurance
104 running events (3 km, ½ marathon and 8 km cross country running). Some subjects in the OS
105 and OE groups participated in several sprint or endurance running events, respectively. Mean
106 results of the subjects competing in the championships were: 60 m 9.13 ± 0.48 (sec, n=8), 60 m
107 hurdles 10.15 (sec, n=1), 200 m 30.64 ± 1.97 (sec, n=7), 400 m $1:13 \pm 8$ (min:sec, n=3), 3000 m
108 $13:48 \pm 60$ (min:sec, n=4), ½ marathon $1:43:37 \pm 11:34$ (hr:min:sec, n=5) and 8 km cross
109 country running $44:19 \pm 6:25$ (min:sec, n=7). These results correspond to 8, 1, 11, 16, 22 and 16
110 % slower than the world record times for 75 year old men in 60 m, 60 m hurdles, 200 m, 400 m,
111 3000 m and ½ marathon respectively. Thus, the participating subjects can be considered to be
112 highly competitive athletes.

113 Subject exclusion criteria were Achilles tendon pain, history of Achilles tendon rupture or
114 surgery, pain in calf muscles during measurements, neurologic and progressive severe illnesses,
115 insulin treated diabetes, fracture within the previous year, immobilization for one week during
116 the last three months, daily use of painkillers, use of immunosuppressive drugs or anticoagulants,
117 or severe visual or hearing impairment.

118 The ethics committee of the Central Finland Health Care District approved the study. All
119 participants signed an informed consent prior to participating in the study and measurements
120 were conducted according to the standards set by the latest revision of the Declaration of
121 Helsinki.

122 **Measurements**

123 Training characteristics of OE and OS groups were assessed with self-reported questionnaire.
124 The athletes were asked about their training history (yr.), overall training volume (h/wk.) and
125 amount of endurance (km/wk.), sprint (sessions/wk.) and strength training (sessions/wk.) in their
126 current normal training routines.

127 Laboratory measurements included assessment of triceps surae muscle architecture and size and
128 Achilles tendon cross-sectional area and mechanical properties. The measurement procedures
129 have been previously described in detail (44) but are briefly described below.

130 For the measurements of Achilles tendon and both gastrocnemius and soleus muscle architecture
131 and size at rest, the subjects were lying prone facing down with ankle angle at 90°. Tendon
132 cross-sectional area (mm^2) was measured from a B-mode ultrasound image taken 4 cm proximal
133 from the proximal border of the calcaneal tubercle where the free Achilles tendon typically
134 reaches its smallest cross-sectional area (38). Body mass normalized tendon cross-sectional area
135 was calculated by dividing cross-sectional area by body mass^{2/3} (20). Muscle architecture from
136 medial gastrocnemius and soleus muscles was assessed from ultrasound images taken at 50% of
137 medial gastrocnemius length and mid-muscle belly in the medial-lateral direction. Fascicle
138 length (mm), pennation angle (°) and muscle thickness (mm) were measured from the images. In
139 order to take into account between-subject differences in stature, fascicle length was normalized
140 to tibia length. The combined anatomical cross-sectional area (cm^2) of medial and lateral
141 gastrocnemius was measured from a panoramic B-mode ultrasound image taken at 50% of
142 medial gastrocnemius length as a measure of the size of the gastrocnemius muscles. All
143 measurements from ultrasound images were taken twice using an open source computer program

144 (ImageJ 1.44b, National Institutes of Health, US) and the mean was used for subsequent data
145 analysis.

146 For the measurement of Achilles tendon mechanical properties, the subjects were seated in a
147 custom built dynamometer with ankle angle at 90° , knee fully extended and hip at 60° of flexion
148 (full extension 0°). After a standardized warm-up, three maximal voluntary contractions (MVC)
149 lasting approximately 3 seconds were performed with strong verbal encouragement to measure
150 plantar flexion strength (Nm). The highest value obtained during MVC trials was used for
151 subsequent analysis. Warm-up and plantar flexion MVCs served to precondition the tendon
152 before the measurement of tendon mechanical properties (30). Achilles tendon mechanical
153 properties were measured from several isometric plantar flexion contractions up to a force level
154 of 80% of MVC. Tendon force was calculated by multiplying measured reaction force by the
155 ratio between Achilles tendon moment arm length and moment arm of the reaction force.
156 Achilles tendon moment arm was defined as the distance from the center of the Achilles tendon
157 to the outermost tip of the medial malleolus in the sagittal plane measured using a ruler. The
158 moment arm of the reaction force around the ankle joint was defined as the sagittal plane
159 distance between the outermost tip of the medial malleolus and the head of the first metatarsal.
160 Achilles tendon elongation (mm) was defined as the change in the distance between the proximal
161 border of the calcaneal tubercle and the medial gastrocnemius muscle-tendon junction. Changes
162 in the location of the calcaneal tubercle in the laboratory coordinate system were measured using
163 a potentiometer that measures heel lift from the dynamometer footplate. Medial gastrocnemius
164 muscle-tendon junction location in the laboratory coordinate system was measured with a
165 combination of B-mode ultrasonography and motion analysis. Ultrasound images of the muscle-
166 tendon junction were collected at 70 Hz and the location of the muscle tendon junction within the

167 image was defined by automatic tracking software (32). The location of the muscle-tendon
168 junction was converted to the laboratory coordinate system using video based motion capture of
169 the ultrasound probe. Two parameters that describe tendon mechanical properties were
170 calculated, tendon stiffness (N/mm) and Young's modulus (GPa). Tendon stiffness characterizes
171 mechanical properties of the tendon and is defined as the slope of the linear portion of the tendon
172 force-elongation curve. We calculated tendon stiffness as a linear fit to force-elongation data
173 from 10 to 80% MVC force since the curves were almost perfectly linear in this region (Fig. 1,
174 $r^2=0.999$ from linear fits to average force-elongation curves). Tendon Young's modulus is the
175 slope of the linear portion of the tendon stress-strain curve and represents tendon stiffness
176 normalized to tendon dimensions. Young's modulus describes the mechanical properties of the
177 material from which a tendon is composed. To derive Young's modulus, Achilles tendon stress
178 (Pa) was calculated by dividing Achilles tendon force (N) by tendon cross-sectional area (m^2)
179 and strain (%) was calculated by dividing elongation (mm) by initial tendon length (mm).
180 Young's modulus was calculated as a linear fit to force-elongation data from 10 to 80% MVC
181 force.

182 **Statistical analyses**

183 Due to inadequate image quality, soleus muscle architecture data were excluded for two subjects
184 from the OS group and three from the OC group, while medial gastrocnemius muscle
185 architecture data were excluded for one subject from OE, and gastrocnemius cross-sectional area
186 data from one subject from OC. Data were first checked for normality with Shapiro-Wilk test
187 and for homogeneity of variance with Levene's test. Differences in muscle and tendon properties
188 between the groups were tested using single factor analysis of variance and Tukey-Kramer post
189 hoc test. Games-Howell post hoc test was used when inhomogeneous variances between the

190 groups were observed and Kruskal-Wallis test with Bonferroni correction for non-normally
191 distributed variables. Differences in training characteristics between OE and OS were tested
192 using Mann-Whitney U-test. The level of statistical significance was set at $\alpha = 0.05$ for all tests.
193 Statistical analyses were performed using IBM SPSS Statistics (version 20.0.0.2). Standardized
194 mean differences between YC and groups of older adults were calculated for main results of the
195 study (Table 2 and 3) as a measure of effect sizes using Hedges' g including a correction for
196 small sample bias (12).

197 **Results**

198 Subject characteristics and training status for the older athletes are reported in Table 1. Older
199 adults in the three different groups were matched for age, height and body mass. YC were
200 significantly taller than OC ($p < 0.001$). OC had significantly greater BMI compared to YC
201 ($p = 0.006$) and OE ($p = 0.009$). Significantly lower plantar flexion strength was found in OC (34%,
202 $p = 0.001$) and OE (42%, $p < 0.001$) compared to YC but not in OS compared to YC ($p = 0.077$). OE
203 and OS groups did not differ in years of training, hours of training per week, or number of
204 strength training sessions per week. Endurance training measured in distance was 8 times greater
205 in OE in comparison with OS ($p < 0.001$), and OS did 3 times more sprint training sessions per
206 week than OE ($p = 0.006$).

207 Achilles tendon cross-sectional area was 22, 45 and 71% larger in OC ($p = 0.022$), OE ($p = 0.001$)
208 and OS ($p < 0.001$) compared to YC, respectively (Table 2). Tendon cross-sectional area in OS
209 was significantly larger than in OC ($p = 0.033$). Body mass normalized tendon cross-sectional area
210 yielded similar results to the unnormalized values. No statistically significant differences were

211 observed between the groups in Achilles tendon stiffness (Figure 1) but Young's modulus was
212 31, 35, and 44% smaller in OC ($p<0.001$), OE ($p=0.001$) and OS ($p<0.001$) in comparison to YC,
213 respectively. Maximal tendon force during MVC was significantly lower in OC (35%, $p<0.001$)
214 and OE (38%, $p<0.001$) but not in OS ($p=0.156$) compared to YC. Average tendon stress during
215 MVC was greater in YC than the older groups ($p<0.001$). Tendon elongation at 80% MVC was
216 significantly greater in YC compared to OC ($p=0.014$) but the difference did not reach statistical
217 significance in OE ($p=0.114$) or OS ($p=0.352$). However, effect sizes between YC and OE and
218 OS were greater than the effect size between YC and OC. The groups did not differ significantly
219 in tendon strain at 80% MVC.

220 Results of soleus and gastrocnemius muscle architecture and size, as well as plantar flexion
221 muscle strength, are presented in Table 3. Soleus fascicle length was significantly shorter in OE
222 compared to YC (absolute $p=0.014$, normalized $p=0.002$) and also compared to OC (absolute
223 $p=0.047$, normalized $p<0.001$). No significant differences were found in soleus pennation angle
224 or muscle thickness. Medial gastrocnemius fascicle length and pennation angle did not differ
225 between the groups. In OC, medial gastrocnemius muscle thickness was significantly smaller in
226 contrast to YC ($p=0.043$) and gastrocnemius cross-sectional area was significantly smaller in
227 contrast to YC ($p=0.011$) and OS ($p=0.011$).

228 **Discussion**

229 We examined selected triceps surae muscle-tendon properties of two differently trained groups
230 of older athletes with an average of 42 years of regular running training, and compared them to
231 untrained age-matched older and young adults. The main findings of the study were that Achilles

232 tendon cross-sectional area was significantly larger in all older adult groups than young adults,
233 and in older sprinters compared to age-matched untrained older adults, while there were no
234 statistically significant group differences in Achilles tendon stiffness. The greater tendon cross-
235 sectional area was also reflected in tendon Young's modulus and tendon average tensile stress
236 during maximal isometric force production, both of which were significantly lower in all older
237 groups compared to young untrained adults. Only minor differences were observed in triceps
238 surae muscle architecture, the most important being significantly shorter fascicle length in soleus
239 muscle in older endurance runners. The current study adds new insight into possible effects of
240 exercise loading on muscle and tendon structure and function in older age. The novelty of the
241 current study is that measurements of triceps surae muscle architecture and Achilles tendon
242 properties were made from top-level older athletes that included both endurance and sprint
243 runners.

244 *Achilles tendon properties*

245 To the best of our knowledge, this is the first study to show greater Achilles tendon cross-
246 sectional area in older adults with a regular exercise training background. Contradicting our
247 hypothesis, the results suggest that long-term exercise did not counteract the age-related increase
248 in Achilles tendon cross-sectional area. Previous cross-sectional studies suggest that Achilles
249 tendon cross-sectional area increases in response to both long-term exercise loading (20, 33) and
250 normal aging (31, 44). The present results suggest that the Achilles tendon responds to regular
251 loading by increasing cross-sectional area in an intensity-dependent manner. Moreover, the
252 increase in cross-sectional area appears to be additive to the increase due to normal aging. This
253 finding supports recent findings by Couppé et al. (7), who showed that regular endurance

254 running was associated with larger patella tendon cross-sectional area in both young and older
255 adults.

256 A possible explanation for ageing and exercise training to be associated with larger tendon cross-
257 sectional area is that tendon hypertrophy is needed to compensate for an age-related decrease in
258 mechanical properties of the tendon collagen structure. Another possible explanation is that
259 greater tendon cross-sectional area in older adults is observed as a consequence of intratendinous
260 accumulation of lipids or water. These two possible mechanisms are not exclusive and could
261 together explain the observed results. The following paragraphs introduce these proposed
262 explanations in more detail.

263 In animal models, aging has been linked with an increase in type V collagen and a greater
264 proportion of small collagen fibrils, which probably contribute to concurrently observed reduced
265 ultimate tensile stress (10, 48). Greater tendon cross-sectional area in older adults could be due to
266 a necessary adaptation to reduce maximal tendon stress to safe levels for older tendons that
267 possibly have reduced ultimate tensile stress. To reduce the stress to a safe level, cross-sectional
268 area must be proportional to maximal force acting on the tendon, thus explaining the greater
269 cross-sectional area in older sprint runners compared to older untrained adults observed in the
270 current study.

271 Greater tendon cross-sectional area in older adults could also serve to maintain sufficient
272 stiffness, which could be important both for protecting the tendon from strain-induced damage
273 and for muscle function. A possible age-related reduction in stiffness of tendon collagen
274 structure may be partly compensated by an age-related increase in collagen cross-links,
275 especially in advanced glycation end product cross-links (6), which stabilize collagen structure

276 and may increase tendon stiffness. Life-long endurance running has been shown to be associated
277 with lower advanced glycation end product cross-link density (7). If older athletes in the current
278 study had a lower density of collagen cross-links, this could explain the requirement for older
279 athletes to have even greater tendon cross-sectional area compared to untrained older adults, in
280 order to maintain tendon stiffness with aging.

281 Based on current knowledge of tendon adaptation, loading intensity is the main factor
282 determining adaptations in tendon mechanical properties (5). Thus, it seems unlikely that sprint-
283 trained older athletes would have the lowest Achilles tendon Young's modulus among the groups
284 in the current study. A possible explanation could be that larger tendon cross-sectional area in
285 older adults is not an adaptation to lowered tendon Young's modulus. Instead it could be due to
286 accumulation of tendon subcomponents that do not markedly affect tendon mechanical behavior.
287 These could include extracellular lipid deposits and proteoglycans and glycosaminoglycans that
288 attract water. Extracellular lipid deposits within tendon have been associated with aging (14) and
289 this could be common to all older adults irrespective of exercise training. On the other hand,
290 production of proteoglycans and glycosaminoglycans could be increased with exercise training
291 induced tendon loading (15). This would explain the observed lower Young's modulus and stress
292 of the tendon in older adults in the present study, and also explains why greater tendon cross-
293 sectional area was not related to greater tendon stiffness.

294 Within- and between-operator reliability of Achilles tendon cross-sectional area measurement
295 using ultrasound imaging has been reported to be good (11, 50). In the current study, duplicate
296 analysis of tendon cross-sectional area images produced intraclass correlation 0.989 and typical
297 error 2.1%. However, validity of tendon cross-sectional area measurement using ultrasound
298 imaging is not known, thus the results should be interpreted with some caution. Future studies

299 should try to replicate the findings of the current study, preferably using magnetic resonance
300 imaging, which allows measurements of tendon cross-sectional area along the whole tendon.
301 More research examining tendon composition and collagen structure in older adults is also
302 warranted to explain the mechanisms behind changes in tendon cross-sectional area.

303 In contrast to our hypothesis that life-long running would mitigate age-related changes in tendon
304 mechanical properties, we found that Young's modulus was significantly lower in older
305 compared to young adults irrespective of training status, with no significant differences between
306 the older groups. There were also no significant between-group differences in initial tendon
307 length or tendon stiffness. Thus, the lower Young's modulus in older compared to young adults
308 can be attributed mainly to the larger tendon cross-sectional area in older adults.

309 It should be noted that a toe-region with a lower slope of the tendon force-elongation curve at
310 low forces or stresses was not observed (Figure 1). We think that the reason for highly linear
311 force-elongation/stress-strain curves is initial force acting on the Achilles tendon at a 90° ankle
312 angle, and the fact that we calculated the curve starting from 10% MVC force. Lack of toe-
313 region has also been previously observed for Achilles tendon in vivo when elongation is
314 measured from the medial gastrocnemius muscle-tendon junction (27), as done in the present
315 study.

316 To summarize the findings regarding tendon mechanical properties, Young's modulus of the
317 Achilles tendon was significantly lower in older compared to young adults, irrespective of
318 training status. Despite this, Achilles tendon stiffness was conserved in all groups of older adults.
319 Thus, the lower muscle strength, greater tendon cross-sectional area and conserved tendon
320 stiffness resulted in reduced maximal tendon stress and strain in older adults. Reduced tendon

321 stress and strain could be a necessary mechanism to decrease the probability of tendon injury, as
322 aging may decrease tendon fascicle sliding that possibly leads to greater loading of the fascicles
323 themselves (47). A functional consequence of similar Achilles tendon stiffness but lower muscle
324 strength in older compared to young adults is a limited maximal capacity for elastic energy
325 storage and subsequent utilization during locomotion. This may contribute to the reported greater
326 metabolic cost of transport in older compared to young adults (35).

327 *Triceps surae muscle architecture, size and strength*

328 The present data also suggest that, in general, muscle architecture is not greatly different in older
329 habitual runners in contrast to both untrained older or young adults. Soleus fascicle length was
330 found to be significantly shorter in endurance-trained older adults than young and older
331 untrained adults. Although somewhat speculative, it may be that shorter fascicles observed in
332 long-term endurance runners in the current study are due to adaptation that improves the
333 efficiency of force production in locomotion. Soleus has short muscle fascicles compared to
334 tendon length (51). Consequently, soleus muscle operates mainly as a force rather than a power
335 producer in locomotion (4). Thus, as this muscle does not need to produce large amounts of
336 work, short fascicles may decrease the energy cost of force production due to lower activated
337 muscle volume per unit of force output compared to longer fascicles (29). We recently observed
338 that shorter fascicle length in soleus and gastrocnemius was associated with better mobility in
339 older adults (45), further supporting the suggestion that shorter soleus fascicle length in older
340 endurance runners may be an adaptive response to life-long exercise training.

341 Another finding of the present study is that long-term endurance running was not associated with
342 greater strength or size of triceps surae muscles compared to untrained older controls. Plantar

343 flexion strength and maximal tendon force in endurance-trained older adults was significantly
344 lower in comparison to young adults. Moreover, the effect sizes for the difference in
345 gastrocnemius thickness and cross-sectional area were comparable to those between young and
346 older controls, which were also statistically significant. Taken together, these results suggest that
347 endurance running is not a sufficient stimulus for maintenance of muscle mass and size with
348 aging.

349 In contrast, the current data suggest that high intensity loading due to sprint training may be an
350 effective stimulus to counteract the age-related decline in both muscle mass and strength in
351 triceps surae muscles. We observed that gastrocnemius muscle cross-sectional area was
352 significantly larger in sprint-trained older adults compared to untrained older controls. In
353 addition, plantar flexion strength and maximal tendon force were not significantly different from
354 those of young controls, with about half the effect size as in endurance-trained older adults
355 compared to young controls. These findings are supported by previous studies in young adults in
356 which sprint running but not endurance running was associated with greater muscle strength and
357 size in triceps surae muscles (18, 23). It may be that the beneficial effects of sprint training
358 preferentially target gastrocnemius muscle, which contains more fast twitch muscle fibers than
359 soleus (13).

360 *Methodological considerations*

361 The strengths of the current study are that the world-class older athletes measured in the present
362 study had a life-long physical activity background and had performed many decades of regular
363 exercise training. In addition, both the trained and untrained older adults were over 70 years old
364 and thus can be assumed to be affected by primary biological aging.

365 Limitations of the present study include the cross-sectional study design which does not allow
366 conclusions about cause-effect relationships that a longitudinal study design may allow. Cross-
367 sectional studies can be affected by selection bias. It is possible that subjects with favorable
368 muscle-tendon properties for endurance or sprint running were more likely to participate in such
369 activities. However, we did not observe differences between older trained and untrained subjects
370 in genetically determined variables such as Achilles tendon moment arm, forefoot length or
371 Achilles tendon length, all of which are related to running performance (3, 25, 26, 42). This
372 suggests that selection bias caused by genetic predisposition towards favorable musculoskeletal
373 properties for running did not considerably affect our data, although the possibility of selection
374 bias cannot be completely excluded. Another limitation of the current study is the small sample
375 size. However, it was not possible to obtain a larger sample of older athletes from the highest
376 performance level.

377 *Conclusions*

378 The current findings suggest that triceps surae muscle size, architecture, strength and tendon
379 stiffness are relatively unaffected by long-term running training in older adults. The reason for
380 this finding may be that the triceps surae muscle group is highly loaded in daily activities and
381 thus training produces only a small relative overload to this muscle group. Considering the
382 unparalleled physical performance of the older athletes in the present study, it appears that the
383 measured triceps surae muscle-tendon properties are not the key determining factors in their
384 physical performance. However, relatively high individual variation in these properties suggests
385 that a well-functioning muscle-tendon unit may be achieved via different combinations of muscle
386 and tendon properties. In addition, it is likely that in the current study there were differences
387 between the groups in factors that were not measured but that affect physical performance. These

388 include muscle fiber type, composition, molecular level modifications in contractile proteins and
389 neural activation (8, 21, 41). To further elucidate the importance of muscle architecture and
390 tendon mechanical properties for physical performance, future studies should investigate how
391 aging and physical loading affect muscle-tendon interaction during locomotion.

392 In conclusion, our data suggest that long-term physical loading induced by either endurance or
393 sprint running does not have a significant effect on Achilles tendon stiffness in older adults.
394 However, the loading patterns associated with sprint and endurance training in older age both
395 appear to increase Achilles tendon cross-sectional area in an intensity dependent manner.
396 Furthermore, the present results suggest that sprint running but not endurance running may
397 mitigate age-related loss of muscle mass and strength in triceps surae muscles. On the other
398 hand, endurance training in older age may alter muscle architecture in a way that is beneficial for
399 movement economy.

400 **Grants**

401 This work was supported by grants from the Finnish Cultural Foundation to L.S. and by the EC
402 FP7 Collaborative Project MYOAGE (GA-223576).

403 **References**

- 404 1. **Abe T, Kumagai K, Brechue WF.** Fascicle length of leg muscles is greater in sprinters
405 than distance runners. *Med Sci Sports Exerc* 32: 1125–9, 2000.
- 406 2. **Arampatzis A, Karamanidis K, Morey-Klapsing G, De Monte G, Stafilidis S.**
407 Mechanical properties of the triceps surae tendon and aponeurosis in relation to intensity
408 of sport activity. *J Biomech* 40: 1946–1952, 2007.

- 409 3. **Baxter JR, Novack T a., Van Werkhoven H, Pennell DR, Piazza SJ.** Ankle joint
410 mechanics and foot proportions differ between human sprinters and non-sprinters. *Proc R*
411 *Soc B Biol Sci* 279: 2018–2024, 2012.
- 412 4. **Biewener AA, Roberts TJ.** Muscle and tendon contributions to force, work, and elastic
413 energy savings: a comparative perspective. *Exerc Sport Sci Rev* 28: 99–107, 2000.
- 414 5. **Bohm S, Mersmann F, Arampatzis A.** Human tendon adaptation in response to
415 mechanical loading: a systematic review and meta-analysis of exercise intervention
416 studies on healthy adults. *Sport Med - Open* 1: 7, 2015.
- 417 6. **Couppé C, Hansen P, Kongsgaard M, Kovanen V, Suetta C, Aagaard P, Kjaer M,**
418 **Magnusson SP.** Mechanical properties and collagen cross-linking of the patellar tendon in
419 old and young men. *J Appl Physiol* 107: 880–886, 2009.
- 420 7. **Couppé C, Svensson RB, Grosset JF, Kovanen V, Nielsen RH, Olsen MR, Larsen JO,**
421 **Praet SFE, Skovgaard D, Hansen M, Aagaard P, Kjaer M, Magnusson SP.** Life-long
422 endurance running is associated with reduced glycation and mechanical stress in
423 connective tissue. *Age* 36: 9665, 2014.
- 424 8. **D’Antona G, Pellegrino MA, Carlizzi CN, Bottinelli R.** Deterioration of contractile
425 properties of muscle fibres in elderly subjects is modulated by the level of physical
426 activity. *Eur J Appl Physiol* 100: 603–611, 2007.
- 427 9. **Delmonico MJ, Harris TB, Visser M, Park SW, Conroy MB, Velasquez-Mieyer P,**
428 **Boudreau R, Manini TM, Nevitt M, Newman AB, Goodpaster BH, Health and Body**
429 **A.** Longitudinal study of muscle strength, quality, and adipose tissue infiltration. *Am J*
430 *Clin Nutr* 90: 1579–1585, 2009.
- 431 10. **Dressler MR, Butler DL, Wenstrup R, Awad HA, Smith F, Boivin GP.** A potential
432 mechanism for age-related declines in patellar tendon biomechanics. *J Orthop Res* 20:
433 1315–1322, 2002.
- 434 11. **Dudley-Javoroski S, McMullen T, Borgwardt MR, Peranich LM, Shields RK.**
435 Reliability and Responsiveness of Musculoskeletal Ultrasound in Subjects with and
436 without Spinal Cord Injury. *Ultrasound Med Biol* 36: 1594–1607, 2010.
- 437 12. **Durlak JA.** How to select, calculate, and interpret effect sizes. *J. Pediatr. Psychol.* 34:
438 917–928, 2009.
- 439 13. **Edgerton VR, Smith JL, Simpson DR.** Muscle fibre type populations of human leg
440 muscles. *Histochem J* 7: 259–66, 1975.
- 441 14. **Finlayson R, Woods SJ.** Lipid in the Achilles tendon. A comparative study.
442 *Atherosclerosis* 21: 371–89, 1975.

- 443 15. **Franchi M, Torricelli P, Giavaresi G, Fini M.** Role of moderate exercising on Achilles
444 tendon collagen crimping patterns and proteoglycans. *Connect Tissue Res* 54: 267–74,
445 2013.
- 446 16. **Frontera WR, Hughes VA, Lutz KJ, Evans WJ.** A cross-sectional study of muscle
447 strength and mass in 45- to 78-yr-old men and women. *J Appl Physiol* 71: 644–650, 1991.
- 448 17. **Goodpaster B, Park S, Harris T, Kritchevsky S, Nevitt M, Schwartz A, Simonsick E,
449 Tylavsky F, Visser M, Newman A.** The loss of skeletal muscle strength, mass, and
450 quality in older adults: the health, aging and body composition study. *Journals Gerontol
451 Ser A Biol Sci Med Sci* 61: 1059–1064, 2006.
- 452 18. **Hansen P, Aagaard P, Kjaer M, Larsson B, Magnusson SP.** Effect of habitual running
453 on human Achilles tendon load-deformation properties and cross-sectional area. *J Appl
454 Physiol* 95: 2375–2380, 2003.
- 455 19. **Karamanidis K, Arampatzis A.** Mechanical and morphological properties of human
456 quadriceps femoris and triceps surae muscle-tendon unit in relation to aging and running.
457 *J Biomech* 39: 406–417, 2006.
- 458 20. **Kongsgaard M, Aagaard P, Kjaer M, Magnusson SP.** Structural Achilles tendon
459 properties in athletes subjected to different exercise modes and in Achilles tendon rupture
460 patients. *J Appl Physiol* 99: 1965–1971, 2005.
- 461 21. **Korhonen MT, Cristea A, Alén M, Häkkinen K, Sipilä S, Mero A, Viitasalo JT,
462 Larsson L, Suominen H.** Aging, muscle fiber type, and contractile function in sprint-
463 trained athletes. *J Appl Physiol* 101: 906–917, 2006.
- 464 22. **Korhonen MT, Heinonen A, Siekkinen J, Isolehto J, Alén M, Kiviranta I, Suominen
465 H.** Bone density, structure and strength, and their determinants in aging sprint athletes.
466 *Med Sci Sports Exerc* 44: 2340–2349, 2012.
- 467 23. **Kubo K, Ikebukuro T, Yata H, Tomita M, Okada M.** Morphological and mechanical
468 properties of muscle and tendon in highly trained sprinters. *J Appl Biomech* 27: 336–344,
469 2011.
- 470 24. **Kulmala JP, Korhonen MT, Kuitunen S, Suominen H, Heinonen A, Mikkola A,
471 Avela J.** Which muscles compromise human locomotor performance with age? *J R Soc
472 Interface* 11, 2014.
- 473 25. **Kunimasa Y, Sano K, Oda T, Nicol C, Komi P V., Locatelli E, Ito a., Ishikawa M.**
474 Specific muscle-tendon architecture in elite Kenyan distance runners. *Scand J Med Sci
475 Sport* 24: 1–6, 2014.
- 476 26. **Lee SSM, Piazza SJ.** Built for speed: musculoskeletal structure and sprinting ability. *J
477 Exp Biol* 212: 3700–3707, 2009.

- 478 27. **Lichtwark GA, Wilson AM.** In vivo mechanical properties of the human Achilles tendon
479 during one-legged hopping. *J Exp Biol* 208: 4715–4725, 2005.
- 480 28. **Lieber RL, Friden J.** Clinical significance of skeletal muscle architecture. *Clin Orthop*
481 *Relat Res* : 140–151, 2001.
- 482 29. **Mademli L, Arampatzis A.** Mechanical and morphological properties of the triceps surae
483 muscle-tendon unit in old and young adults and their interaction with a submaximal
484 fatiguing contraction. *J Electromyogr Kinesiol* 18: 89–98, 2008.
- 485 30. **Maganaris CN.** Tendon conditioning: artefact or property? *Proc R Soc B Biol Sci* 270
486 Suppl: S39–42, 2003.
- 487 31. **Magnusson SP, Beyer N, Abrahamsen H, Aagaard P, Neergaard K, Kjaer M.**
488 Increased cross-sectional area and reduced tensile stress of the Achilles tendon in elderly
489 compared with young women. *J Gerontol A Biol Sci Med Sci* 58: 123–127, 2003.
- 490 32. **Magnusson SP, Hansen P, Aagaard P, Brond J, Dyhre-Poulsen P, Bojsen-Moller J,**
491 **Kjaer M.** Differential strain patterns of the human gastrocnemius aponeurosis and free
492 tendon, in vivo. *Acta Physiol Scand* 177: 185–195, 2003.
- 493 33. **Magnusson SP, Kjaer M.** Region-specific differences in Achilles tendon cross-sectional
494 area in runners and non-runners. *Eur J Appl Physiol* 90: 549–553, 2003.
- 495 34. **McPhee JS, Hogrel JY, Maier AB, Seppet E, Seynnes OR, Sipila S, Bottinelli R,**
496 **Barnouin Y, Bijlsma AY, Gapeyeva H, Maden-Wilkinson TM, Meskers CG, Paasuke**
497 **M, Sillanpaa E, Stenroth L, Butler-Browne G, Narici M V, Jones DA.** Physiological
498 and functional evaluation of healthy young and older men and women: design of the
499 European MyoAge study. *Biogerontology* 14: 325–337, 2013.
- 500 35. **Mian OS, Thom JM, Ardigò LP, Narici M V., Minetti a. E.** Metabolic cost, mechanical
501 work, and efficiency during walking in young and older men. *Acta Physiol* 186: 127–139,
502 2006.
- 503 36. **Morse CI, Thom JM, Birch KM, Narici M V.** Changes in triceps surae muscle
504 architecture with sarcopenia. *Acta Physiol Scand* 183: 291–298, 2005.
- 505 37. **Narici M V, Maganaris CN, Reeves ND, Capodaglio P.** Effect of aging on human
506 muscle architecture. *J Appl Physiol* 95: 2229–2234, 2003.
- 507 38. **Peltonen J, Cronin NJ, Avela J, Finni T.** In vivo mechanical response of human
508 Achilles tendon to a single bout of hopping exercise. *J Exp Biol* 213: 1259–1265, 2010.
- 509 39. **Rantanen T, Masaki K, Foley D, Izmirlian G, White L, Guralnik JM.** Grip strength
510 changes over 27 yr in Japanese-American men. *J Appl Physiol* 85: 2047–2053, 1998.

- 511 40. **Rantanen T.** Midlife hand grip strength as a predictor of old age disability. *JAMA* 281:
512 558, 1999.
- 513 41. **Reid KF, Doros G, Clark DJ, Patten C, Carabello RJ, Cloutier GJ, Phillips EM,**
514 **Krivickas LS, Frontera WR, Fielding RA.** Muscle power failure in mobility-limited
515 older adults: preserved single fiber function despite lower whole muscle size, quality and
516 rate of neuromuscular activation. *Eur J Appl Physiol* 112: 2289–2301, 2012.
- 517 42. **Scholz MN, Bobbert MF, van Soest AJ, Clark JR, van Heerden J.** Running
518 biomechanics: shorter heels, better economy. *J Exp Biol* 211: 3266–3271, 2008.
- 519 43. **Sipilä S, Suominen H.** Quantitative ultrasonography of muscle: detection of adaptations
520 to training in elderly women. *Arch Phys Med Rehabil* 77: 1173–8, 1996.
- 521 44. **Stenroth L, Peltonen J, Cronin NJ, Sipila S, Finni T.** Age-related differences in
522 Achilles tendon properties and triceps surae muscle architecture in vivo. *J Appl Physiol*
523 113: 1537–1544, 2012.
- 524 45. **Stenroth L, Sillanpää E, McPhee JS, Narici M V., Gapeyeva H, Pääsuke M,**
525 **Barnouin Y, Hogrel J-YJ-Y, Butler-Browne G, Bijlsma A, Meskers CGM, Maier AB,**
526 **Finni T, Sipilä S, Sillanpää E, Paasuke M, Sipila S.** Plantarflexor muscle-tendon
527 properties are associated with mobility in healthy older adults. *Journals Gerontol. Ser. A*
528 *Biol. Sci. Med. Sci.* (2015). doi: 10.1093/gerona/glv011.
- 529 46. **Tanaka H, Seals DR.** Endurance exercise performance in Masters athletes: age-
530 associated changes and underlying physiological mechanisms. *J Physiol* 586: 55–63,
531 2008.
- 532 47. **Thorpe CT, Udeze CP, Birch HL, Clegg PD, Screen HRC.** Capacity for sliding
533 between tendon fascicles decreases with ageing in injury prone equine tendons: A possible
534 mechanism for age-related tendinopathy? *Eur Cells Mater* 25: 48–60, 2012.
- 535 48. **Wood LK, Arruda EM, Brooks S V.** Regional stiffening with aging in tibialis anterior
536 tendons of mice occurs independent of changes in collagen fibril morphology. *J. Appl.*
537 *Physiol.* 111: 999–1006, 2011.
- 538 49. **Yataco AR, Busby-Whitehead J, Drinkwater DT, Katzel LI.** Relationship of body
539 composition and cardiovascular fitness to lipoprotein lipid profiles in master athletes and
540 sedentary men. *Aging* 9: 88–94.
- 541 50. **Ying M, Yeung E, Li B, Li W, Lui M, Tsoi CW.** Sonographic evaluation of the size of
542 Achilles tendon: The effect of exercise and dominance of the ankle. *Ultrasound Med Biol*
543 29: 637–642, 2003.
- 544 51. **Zajac FE.** Muscle and tendon: properties, models, scaling, and application to
545 biomechanics and motor control. *Crit Rev Biomed Eng* 17: 359–411, 1989.

546 52. **Zampieri S, Pietrangelo L, Loeffler S, Fruhmann H, Vogelauer M, Burggraf S, Pond**
547 **A, Grim-Stieger M, Cvecka J, Sedliak M, Tirpakova V, Mayr W, Sarabon N, Rossini**
548 **K, Barberi L, De Rossi M, Romanello V, Boncompagni S, Musaro A, Sandri M,**
549 **Protasi F, Carraro U, Kern H, Tírpáková V, Mayr W, Sarabon N, Rossini K,**
550 **Barberi L, De Rossi M, Romanello V, Boncompagni S, Musarò A, Sandri M, Protasi**
551 **F, Carraro U, Kern H.** Lifelong physical exercise delays age-associated skeletal muscle
552 decline. *J. Gerontol. A. Biol. Sci. Med. Sci.* (2014). doi: 10.1093/gerona/glu006.

553

554 **Figure legend**

555 FIGURE 1. Mean Achilles tendon force-elongation (upper) and stress-strain (lower)
556 relationships for young controls (YC), older controls (OC), older endurance runners (OE), and
557 older sprint runners (OS). Lines are linear fits and represent Achilles tendon stiffness and
558 Young's modulus, respectively. Values are calculated at 10% MVC increments from 10 to 80%
559 MVC. Standard deviations are omitted for clarity.





TABLE 1. Subject characteristics and training status of older athletes

	YC	OC	OE	OS
Number of subject	18	33	10	10
Age (yr.)	23.7 ± 2.0	74.8 ± 3.6**	74.0 ± 2.8**	74.4 ± 2.8**
Height (cm)	181 ± 6	173 ± 5**	175 ± 7	176 ± 7
Body mass (kg)	75.4 ± 9.0	76.1 ± 7.7	69.9 ± 6.9	74.3 ± 7.1
BMI (kgm ⁻²)	23.1 ± 2.6	25.4 ± 2.4**††	22.7 ± 1.6	24.1 ± 1.9
Plantar flexion strength (Nm)	199 ± 56	132 ± 21**	116 ± 25**	153 ± 39
Years of training			39.4 ± 20.9	44.7 ± 19.7
Hours of training per week			6.8 ± 3.3	6.2 ± 2.6
Endurance training per week (km)			55.2 ± 8.8	6.5 ± 2.8††
Sprint training sessions per week			0.8 ± 0.7	2.3 ± 1.2††
Strength training sessions per week			0.5 ± 0.2	0.9 ± 0.2

Values are expressed as mean ± SD. * significantly different from YC, † significantly different from OE, **/†† p<0.01. YC young controls, OC older controls, OE older endurance runners, OS older sprint runners.

TABLE 2. Achilles tendon cross-sectional area and mechanical properties

	YC	OC	OE	OS
Cross-sectional area (mm ²)	56.5 ± 9.6	69.0 ± 12.2 (-1.05)* ‡	82.0 ± 19.8 (-1.69)**	96.5 ± 24.9 (-2.26)**
Stiffness (N/mm)	186 ± 37	164 ± 47 (0.49)	172 ± 39 (0.34)	166 ± 35 (0.51)
Young's modulus (GPa)	0.86 ± 0.20	0.59 ± 0.17 (1.46)**	0.56 ± 0.22 (1.40)**	0.48 ± 0.19 (1.85)**
Max. tendon force (kN)	3.4 ± 0.9	2.2 ± 0.6 (1.58)**	2.1 ± 0.4 (1.64)**	2.6 ± 0.8 (0.80)
Max. tendon stress (MPa)	59.3 ± 14.9	33.1 ± 9.0 (2.22)**	26.5 ± 8.3 (2.36)**	30.1 ± 14.3 (1.86)**
Elongation at 80 % MVC (mm)	14.3 ± 2.5	11.9 ± 6.4 (0.42)*	11.2 ± 4.4 (0.88)	11.9 ± 4.2 (0.70)
Strain at 80 % MVC (%)	5.6 ± 1.5	4.8 ± 2.2 (0.42)	4.5 ± 1.8 (0.66)	4.7 ± 1.7 (0.56)

Values are expressed as mean ± SD (effect size compared to YC). * significantly different from YC, ‡ significantly different from OS, */‡ p<0.05, ** p<0.01. YC young controls, OC older controls, OE older endurance runners, OS older sprint runners, MG medial gastrocnemius.

TABLE 3. Muscle architecture and size

	YC	OC	OE	OS
Soleus fascicle length (mm)	40.6 ± 8.8	38.6 ± 7.6 (0.24) †	31.2 ± 3.9 (1.18)*	35.3 ± 8.3 (0.57)
Normalized soleus fascicle length (mm/mm)	0.102 ± 0.021	0.100 ± 0.021 (0.11) ††	0.073 ± 0.008 (1.55)**	0.083 ± 0.022 (0.83)
Soleus pennation angle (°)	21.0 ± 5.7	21.2 ± 4.0 (-0.05)	23.7 ± 5.3 (-0.46)	21.6 ± 8.3 (-0.08)
Soleus thickness (mm)	14.3 ± 2.6	13.1 ± 2.7 (0.44)	13.4 ± 2.7 (0.33)	12.8 ± 3.7 (0.49)
MG fascicle length (mm)	47.7 ± 6.6	45.0 ± 7.6 (0.35)	45.3 ± 6.5 (0.34)	47.7 ± 7.0 (0.00)
Normalized MG fascicle length (mm/mm)	0.121 ± 0.018	0.117 ± 0.022 (0.17)	0.108 ± 0.015 (0.71)	0.111 ± 0.021 (0.46)
MG pennation angle (°)	24.8 ± 4.0	24.4 ± 4.2 (0.09)	23.3 ± 4.8 (0.34)	24.1 ± 3.5 (0.18)
MG thickness (mm)	20.1 ± 2.5	17.7 ± 3.2 (0.77)*	17.2 ± 3.6 (0.94)	18.6 ± 2.7 (0.55)
Gastrocnemius CSA (cm ²)	24.2 ± 4.5	20.1 ± 4.5 (0.89)* ‡	20.9 ± 3.4 (0.73)	25.1 ± 4.4 (-0.19)

Values are expressed as mean ± SD (effect size compared to YC). * significantly different from YC, † significantly different from OE, ‡ significantly different from OS, */†/‡ p<0.05, **/†† p<0.01. YC young controls, OC older controls, OE older endurance runners, OS older sprint runners.