

**This is an electronic reprint of the original article.  
This reprint *may differ* from the original in pagination and typographic detail.**

**Author(s):** Waller, Benjamin; Munukka, Matti; Rantalainen, Timo; Lammentausta, Eveliina; Nieminen, Miika T.; Kiviranta, Ilkka; Kautiainen, Hannu; Häkkinen, Arja; Kujala, Urho; Heinonen, Ari

**Title:** Effects of high intensity resistance aquatic training on body composition and walking speed in women with mild knee osteoarthritis : a 4-month RCT with 12-month follow-up

**Year:** 2017

**Version:**

**Please cite the original version:**

Waller, B., Munukka, M., Rantalainen, T., Lammentausta, E., Nieminen, M. T., Kiviranta, I., Kautiainen, H., Häkkinen, A., Kujala, U., & Heinonen, A. (2017). Effects of high intensity resistance aquatic training on body composition and walking speed in women with mild knee osteoarthritis : a 4-month RCT with 12-month follow-up. *Osteoarthritis and Cartilage*, 25(8), 1238-1246.  
<https://doi.org/10.1016/j.joca.2017.02.800>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

# Accepted Manuscript

Effects of high intensity resistance aquatic training on body composition and walking speed in women with mild knee osteoarthritis: a 4-month RCT with 12-month follow-up

Benjamin Waller, Matti Munukka, Timo Rantalainen, Eveliina Lammentausta, Miika T. Nieminen, Ilkka Kiviranta, Hannu Kautiainen, Arja Häkkinen, Urho M. Kujala, Ari Heinonen

PII: S1063-4584(17)30869-5

DOI: [10.1016/j.joca.2017.02.800](https://doi.org/10.1016/j.joca.2017.02.800)

Reference: YJOCA 3978

To appear in: *Osteoarthritis and Cartilage*


Received Date: 3 August 2016

Revised Date: 31 January 2017

Accepted Date: 21 February 2017

Please cite this article as: Waller B, Munukka M, Rantalainen T, Lammentausta E, Nieminen MT, Kiviranta I, Kautiainen H, Häkkinen A, Kujala UM, Heinonen A, Effects of high intensity resistance aquatic training on body composition and walking speed in women with mild knee osteoarthritis: a 4-month RCT with 12-month follow-up, *Osteoarthritis and Cartilage* (2017), doi: 10.1016/j.joca.2017.02.800.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Effects of high intensity resistance aquatic training on body composition and walking**  
2 **speed in women with mild knee osteoarthritis: a 4-month RCT with 12-month follow-up**

3  
4 Benjamin Waller<sup>1</sup>, Matti Munukka<sup>1</sup>, Timo Rantalainen<sup>2</sup>, Eveliina Lammentausta<sup>3</sup>, Miika T  
5 Nieminen<sup>3,4,5</sup>, Ilkka Kiviranta<sup>6</sup>, Hannu Kautiainen<sup>7,8</sup>, Arja Häkkinen<sup>1</sup>, Urho M Kujala<sup>1</sup>, Ari  
6 Heinonen<sup>1</sup>

7  
8 Email addresses:

9 [ben.waller@jyu.fi](mailto:ben.waller@jyu.fi)

10 [matti.munukka@jyu.fi](mailto:matti.munukka@jyu.fi)

11 [t.rantalainen@deakin.edu.au](mailto:t.rantalainen@deakin.edu.au)

12 [eveliina.lammentausta@oulu.fi](mailto:eveliina.lammentausta@oulu.fi)

13 [miika.nieminen@oulu.fi](mailto:miika.nieminen@oulu.fi)

14 [ilkka.kiviranta@helsinki.fi](mailto:ilkka.kiviranta@helsinki.fi)

15 [hannu.kautiainen@medcare.fi](mailto:hannu.kautiainen@medcare.fi)

16 [arja.h.hakkinen@jyu.fi](mailto:arja.h.hakkinen@jyu.fi)

17 [urho.m.kujala@jyu.fi](mailto:urho.m.kujala@jyu.fi)

18 [ari.o.heinonen@jyu.fi](mailto:ari.o.heinonen@jyu.fi)

19  
20 1) Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland.

21 2) Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition  
22 Sciences, Deakin University, Melbourne, Australia

23 3) Department of Diagnostic Radiology, Oulu University Hospital, Oulu, Finland.

24 4) Research Unit of Medical Imaging, Physics and Technology, University of Oulu, Oulu,  
25 Finland

26 5) Medical Research Center, University of Oulu and Oulu University Hospital

27 6) Department of Orthopedics and Traumatology, University of Helsinki and Helsinki  
28 University Hospital, Helsinki, Finland.

29 7) Department of General Practice and Primary Health Care, University of Helsinki,  
30 Helsinki, Finland

31 8) Unit of Primary Health Care, Kuopio University Hospital, Kuopio, Finland

32  
33 *Corresponding author: MSc Benjamin Waller. University of Jyväskylä, Department of Health*  
34 *Sciences, P.O. Box 35, FI-40014 University of Jyväskylä, Finland. +358408053569. E-mail:*  
35 *[ben.waller@jyu.fi](mailto:ben.waller@jyu.fi)*

36  
37 **Running title:** Aquatic training improves walking speed in knee OA

38 **ABSTRACT**

39 **Objective:** To investigate the effects of 4-months' intensive aquatic resistance training on  
40 body composition and walking speed in post-menopausal women with mild knee  
41 osteoarthritis, immediately after intervention and after 12-months follow-up. Additionally,  
42 influence of leisure time physical activity (LTPA) will be investigated.

43 **Design:** This randomised clinical trial assigned 87 volunteer postmenopausal women into  
44 two study arms. The intervention group (n=43) participated in 48 supervised intensive aquatic  
45 resistance training sessions over 4-months while the control group (n=44) maintained normal  
46 physical activity. 84 participants continued into the 12-months' follow-up period. Body  
47 composition was measured with dual-energy X-ray absorptiometry. Walking speed over 2km  
48 and the knee injury and osteoarthritis outcome score (KOOS) were measured. LTPA was  
49 recorded with self-reported diaries.

50 **Results:** After the 4-month intervention there was a significant decrease ( $p=0.002$ ) in fat  
51 mass (mean change:  $-1.17\text{kg}$ ; 95%CI:  $-2.00$  to  $-0.43$ ) and increase ( $p=0.002$ ) in walking speed  
52 ( $0.052\text{m/sec}$ ; 95%CI:  $0.018$  to  $0.086$ ) in favour of the intervention group. Body composition  
53 returned to baseline after 12-months': In contrast, increased walking speed was maintained  
54 ( $0.046\text{m/sec}$  (95%CI  $0.006$  to  $0.086$ ,  $p=0.032$ ). No change was seen in lean mass or KOOS.  
55 Daily LTPA over the 16-months had a significant effect ( $p=0.007$ ) on fat mass loss ( $f^2=0.05$ )  
56 but no effect on walking speed.

57 **Conclusions:** Our findings show that high intensity aquatic resistance training decreases fat  
58 mass and improves walking speed in post-menopausal women with mild knee OA. Only  
59 improvements in walking speed were maintained at 12-months' follow-up. Higher levels of  
60 LTPA were associated with fat mass loss.

61 **Keywords:** Osteoarthritis; Aquatic Exercise; Body Composition; Walking speed

62 **Trial registration number:** ISRCTN6534659

## 1 INTRODUCTION

2 Knee osteoarthritis (OA) is a common cause of pain and activity limitations causing  
3 significant burden on healthcare services<sup>1</sup>. While there is no known treatment that prevents or  
4 reverses OA, traditional management of OA focuses on reducing the symptoms, i.e. pain and  
5 activity limitations, associated with the disease. Recently focus has shifted from treatment of  
6 end-stage OA to preventing progression of the disease, especially in early knee OA<sup>2</sup>. One  
7 possible approach could be to use interventions that address known risk factors for  
8 progression of OA. Risk factors predicting worsening in symptoms and activity limitations  
9 include slow walking speeds, obesity, older age and decreased leisure time physical activity  
10 (LTPA)<sup>3,4</sup>. Obesity is associated with knee OA progression through sub-optimal  
11 biomechanical loading and low-grade systematic inflammation related to high body fat-mass<sup>2</sup>,  
12 <sup>5</sup>. Further, people with knee OA have been shown to walk slower and adapt their gait patterns  
13 in order to avoid pain and to redistribute joint loading<sup>6,7</sup>.

14  
15 Exercise has been shown to evoke positive changes on symptoms and functional capacity as  
16 well as facilitate weight loss<sup>8,9</sup> and is therefore strongly recommended in the management of  
17 knee OA<sup>1,10</sup>. However, pain is a major modulator for activity avoidance in patients with OA  
18 and may limit compliance with land-based exercise<sup>11</sup>. The aquatic environment allows the  
19 individual to exercise with reduced weight bearing and impact on the affected joints<sup>12</sup>. Recent  
20 studies have shown that individuals with lower-limb OA experience significantly less pain  
21 during aquatic compared to land-based exercise of equivalent intensity<sup>13,14</sup>. Our recent  
22 systematic reviews revealed that aquatic exercise evokes both a small and a moderate effect  
23 on physical functioning in people with lower limb OA<sup>15</sup> and healthy older people<sup>16</sup>,  
24 respectively. The difference in effect size is thought, in part, to be due to the higher intensity  
25 of training implemented with the healthy older adults<sup>17</sup>. Further, lack of reporting of actual

26 training intensities achieved in all the included aquatic exercise studies, limits interpretation  
27 of the results. Moreover, higher levels LTPA can have a positive impact on body composition  
28 and also predict a slower progression of OA related symptoms and activity limitations<sup>3, 4</sup>.  
29 LTPA levels have not been reported in any previous aquatic exercise studies and therefore the  
30 effect of this important cofounding factor has not been previously investigated.

31

32 In order to prevent knee OA progression, the exercise intervention should be prescribed early  
33 in the disease progression<sup>2</sup>. To the authors knowledge only one previous study has  
34 investigated the effect evoked by aquatic exercise in the early stage of knee OA  
35 development<sup>18</sup>. Our study, a randomised controlled trial (RCT), indicated that 4-months of  
36 aquatic resistance training improved estimated cardiovascular fitness and had a small  
37 significant impact on tibiofemoral cartilage as measured with quantitative magnetic  
38 resonance imaging (qMRI)<sup>18</sup>. Therefore, aims of this study are to report the effect of 4-  
39 months intensive aquatic resistance training program on body composition and functional  
40 capacity in postmenopausal women with mild knee OA, and whether possible changes are  
41 maintained after 12-months' follow-up. The effect of leisure time physical activity on the  
42 results and the training intensities achieved during the aquatic resistance training will also be  
43 investigated.

## 44 MATERIALS AND METHODS

### 45 Study design

46 This study uses previously unreported outcome data collected from the registered AquaRehab  
47 research project (ISRCTN65346593), a RCT consisting of a 4-month aquatic intervention  
48 with a 12-month follow-up period. Data was collected from January 2012 until April 2014.  
49 The full description of the protocol can be found on open access<sup>19</sup>, which was followed  
50 without changes and a full report of participant recruitment can be found from our previous  
51 study<sup>18</sup>. This study has two experimental arms: 1) aquatic resistance training and 2) control.  
52 Included participants were women aged 60-68 years old with mild knee OA. In this study we  
53 classify mild knee OA as experiencing knee pain on most days, not exceeding 5/10 VAS,  
54 with radiographic changes in tibiofemoral joint grades I (possible osteophytes) or II (definite  
55 osteophytes, possible joint space narrowing) according to the Kellgren Lawrence (K/L)  
56 classification<sup>20</sup>. Pre- and post-intervention results for the qMRI outcomes and patient reported  
57 symptoms have been previously reported<sup>18</sup>. This current study, in addition to patient reported  
58 symptoms, will report the outcomes for body composition and walking speed taken pre- and  
59 post-intervention as well as after the 12-months follow-up<sup>18</sup>. The study design and reporting  
60 follows the CONSORT recommendations for the conducting and reporting of randomized  
61 controlled trials<sup>21</sup>. The study protocol (Dnro 19U/2011) was approved by the Ethics  
62 Committee of the Central Finland Health Care District and conforms to the Declaration of  
63 Helsinki. Written informed consent was obtained from all participants prior to enrolment.

### 65 Subject recruitment

66  
67 Participants were recruited from the county of Central Finland using newspaper  
68 advertisements and telephone recruitment methods. Eligibility criteria was female aged 60-68

69 years old, body mass index (BMI) <35, experiences knee pain almost daily, K/L grades I or II  
70 and no medical reason preventing full participation in intensive exercise. Full eligibility  
71 criteria are described elsewhere<sup>19</sup>.

72

### 73 **Randomisation and blinding**

74

75 The subjects were randomly allocated into one of the two arms of the study by a blinded  
76 external statistician, provided only with randomisation number and OA severity, using a  
77 computer generated block randomization of size of ten, stratified according to K/L grading.

78 The first author performed the DXA imaging but analysis was performed using the  
79 manufactures' in-built software without modification. Physical therapists providing the  
80 intervention also performed the physical performance measures. Principal investigators were  
81 blinded to group allocation.

82

### 83 **Interventions**

84

85 Those participants in the intervention group participated in an aquatic resistance training  
86 sessions lasting 1 hour, 3 times a week for 16 weeks (48 sessions in total). Variable resistance  
87 equipment was used to progress training intensity with three resistance levels; barefoot, small  
88 resistance fins (Theraband products, The Hygienic Corporation, Akron, OH 44310 USA) and  
89 large resistance boots (Hydro-boots, Hydro-Tone Fitness Systems, Inc. Orange, CA 92865-  
90 2760, USA). Training intensity was set at as "hard and fast as possible": A full description of  
91 the training program, its progression and daily training program can be found elsewhere<sup>18</sup>.

92 The control group maintained usual care and were asked to continue their usual leisure time  
93 activities. They were offered the possibility of participating in two sessions consisting of 1



94 hour of light stretching, relaxation and social interaction during the 4-month intervention  
95 period.

96

### 97 **Measures of exercise intensity and perceived exertion**

98

99 Maximum training intensity was ensured by measuring the maximum and average heartrates  
100 and rating of perceived exertion (RPE) for every training session using heart rate monitors  
101 (Polar Oy, Kemble. Finland). Maximum heartrate was estimated using the Karvonen formula  
102 ( $220 - \text{age} = \text{maxHR}$ ) with no adjustments made for the possible effects of immersion. During  
103 the twelfth week capillary blood lactates and repetitions completed for all three training  
104 situations were measured. Self-reported emotional state felt during training was measured  
105 with a 1-5 Likert scale (1-Poor, 2-Tolerable, 3-Satisfactory, 4-Good, 5-Excellent).

106

### 107 **Outcome measures**

108

109 Outcomes for this study are body composition, walking speed and self-reported symptoms.  
110 Body composition (total body fat and lean body mass (kg)) was measured with dual-energy  
111 X-ray absorptiometry (DXA, Lunar Prodigy; GE Lunar Healthcare, Madison, WI, USA). All  
112 full body and regional images were analysed as per manufacturers' protocols using enCORE  
113 software (enCORE 2011, version 13.60.033). In vivo precision of these measurements has  
114 been reported to be high (CV 1.3-2.2%)<sup>22</sup>. Walking speed was calculated from the UKK 2km  
115 walking test. This test requires the subject to walk 2km around a 200m flat track as quickly as  
116 possible without running<sup>23</sup>. Walking speed was calculated in metres per second (m/sec) and  
117 describes walking ability and is a surrogate for aerobic fitness. Self-report pain (Pain),  
118 symptoms (Sym), activities of daily living (ADL), sports and recreation (Sport&Rec) and

119 quality of life (QoL) were measured using be the five domains of the Finnish version of the  
120 Knee Injury and Osteoarthritis Outcome Score (KOOS)<sup>24</sup>. Scores are transformed into a score  
121 0-100 with a score of 0 indicating extreme knee problems and 100 no knee problems<sup>25</sup>.

122

### 123 **Leisure time physical activity (LTPA)**

124

125 LTPA for each participant was calculated for the whole 4-month intervention and 12-month  
126 follow-up period using a daily physical activity diary. Participants recorded type of activity  
127 and self-perceived intensity of each activity, i.e. low, moderate or high, from which metabolic  
128 equivalent task hours (MET/h) per month was calculated<sup>26</sup>. The LTPA for the intervention  
129 group was calculated by combining the MET/h calculated from the aquatic resistance training  
130 and the physical activity diary.

131

### 132 **Statistical Methods**

133

134 Results are displayed as mean and standard deviation (SD) unless otherwise stated. Between  
135 group baseline comparisons were performed using a bootstrap type t-test and Chi-squared.  
136 Repeated measures for walking speed, body composition and all domains of the KOOS were  
137 analysed using generalised linear mixed-models with unstructured correlation structure. Fixed  
138 effects were group, time and group-time interaction. Effect size, standardised beta coefficient  
139 (Beta ( $\beta$ )) adjusted for baseline values, was calculated for post intervention and at 12-month  
140 follow-up. Cohen's standard for Beta values above 0.10, 0.30 and 0.50 represent small,  
141 moderate and large effects respectively<sup>27</sup>. Between group differences in average monthly  
142 LTPA during the 4 month intervention and 12-months' period was tested using a Fisher-  
143 Pitman permutation test for two independent samples. The relationship between average

144 monthly LTPA and body composition and walking speed, after removal of group allocation,  
145 was calculated using a mixed-effects regression model and represented as Cohen's  $f^2$ , where  
146 0.02, 0.15 and 0.35 indicate a small, moderate and large effect respectively<sup>28</sup>. Repeated  
147 ANOVA was used to compare the differences between the three training intensities (barefoot,  
148 small fins and large boots) and measures of training response i.e. RPE, heart rates, blood  
149 lactates, number of repetition performed per session and emotional state. Statistical analyses  
150 were performed using statistical software (Stata, release 13.1, StataCorp, College Station,  
151 Texas).

152

153 This study is a post hoc analysis of original data thus the target sample size (n=70, 35 per  
154 research arm) was calculated based on expect change in the qMRI outcome<sup>18</sup>.

155 **RESULTS**

156

157 In total, 87 participants fulfilled the eligibility criteria and after attending baseline  
158 measurement were randomised into the two treatment arms of the study. There were no  
159 significant differences between the groups in any descriptive variables at baseline (Table 1).  
160 85 participants completed the intervention and 84 agreed to participate in the 12-month  
161 follow-up. In total 76 participants attended measurement at 12-months' follow-up. Participant  
162 recruitment and reasons for loss to follow-up are shown in Figure 1.

163

164

*Table 1 here.*

165

166

*Figure 1 here.*

167

168 **Training intensities achieved during aquatic resistance training**

169

170 Adherence to the aquatic training program was high (88%), with only three subjects attending  
171 less than 70%. Pain during aquatic resistance training in the affected knee was reported more  
172 frequently during the first month (37 times), followed by a gradual decrease in frequency as  
173 the training progressed, with a three-fold reduction in the frequency (12 times) by the 4<sup>th</sup>  
174 month. Pain experienced in affected knee during the intervention was mild 14(16) (VAS 0-  
175 100mm). Training intensity recorded from each complete training session is shown in Table  
176 2. There was a gradual increase in RPE when progressing from barefoot to large resistance  
177 boots while no significant differences in heart rates were measured. A full description of the  
178 daily training intensities measured and pain experienced during training can be found from

179 the supplemental material Appendix A. The attendance for the control group sessions was  
180 68%.

181

182 *Table 2 here.*

183

#### 184 **Treatment effects and maintenance at 12-months**

185

186 Summaries of the treatment effects after 4-months and their maintenance at 12-months  
187 follow-up are presented in Figure 2 and Table 3. After 4-months aquatic resistance training  
188 there was a significant ( $p=0.002$ ) moderate (Beta ( $\beta$ ): 0.32; 95% CI: 0.14 to 0.51) decrease in  
189 fat mass (-4.7% and 0.25% in training and control group respectively) and over-all moderate  
190 ( $\beta$ : 0.34; 0.15 to 0.52) decrease (-1.4% and 0.21%) in body weight ( $p=0.004$ ), both in favour  
191 of the intervention group. There was a significant ( $p<0.001$ ) decrease in fat mass in both legs  
192 -0.47kg (-0.74 to -0.20) or a loss of -4.5% in the training group compared to 1.1% increase in  
193 the control group. There was a similar significant change ( $p=0.007$ ) in the trunk -0.63kg (-1.1  
194 to -0.17) or a loss of -3.1% compared to a 1.0% increase in the control group. Both  
195 significant findings were lost a 12-months follow-up. No localised change in lean mass was  
196 seen at any time point. After the intervention, a significant increase in walking speed  
197 ( $p=0.002$ ) was observed in favour of the intervention group ( $\beta$ : 0.3; 0.12 to 0.50). At 12-  
198 month follow-up walking speed ( $p=0.032$ ) in the intervention group remained significantly  
199 faster compared to the control group ( $\beta$ :0.2; 0.01 to 0.44). No other significant between group  
200 differences could be seen in any domain of the KOOS questionnaire.

201

*Figure 2 here*

202

*Table 3 here*

203

## 204 **Effects of physical activity**

205

206 There was a significant ( $p < 0.001$ ) between group difference in average monthly LTPA during  
207 the intervention period 160 (53) versus 104 (63) MET/h for intervention and control groups  
208 respectively. This difference was immediately lost following cessation of the aquatic training  
209 (Table 3), monthly group averages are depicted in Figure 3. After removal of group allocation  
210 there was a small (Cohen's  $f^2 = 0.05$ ) statistically significant ( $p = 0.007$ ) relationship between  
211 higher average monthly LTPA (MET/h) and greater loss of fat mass. There was no  
212 relationship between LTPA ( $p = 0.52$ ) and lean mass ( $f^2 = 0.002$ ) and a small ( $f^2 = 0.02$ ) but  
213 non-significant ( $p = 0.25$ ) relationship with walking speed. While walking was the most  
214 popular form of LTPA (40.1%) there was no difference seen in activity type or intensities  
215 between the control and intervention group at any time point.

216

217 *Figure 3 here.*

218

## 219 **Harms**

220

221 As previously reported<sup>18</sup>, one subject stopped the intervention following pain experienced  
222 after the first use of the large resistance boot (session 16). One subject complained of  
223 dyspnoea. After education training the participant was able to complete the intervention and  
224 attend follow-up measurements. The results of both participants are included as per intention-  
225 to-treat analysis. No subjects at pre- or post-intervention measurements were unable to walk  
226 the 2km. At 12 months' follow-up two subjects, one from each group could not complete the  
227 2km; both due to a lower-limb injury unrelated to their knee OA.

228 **DISCUSSION**

229

230 Our study indicates that an intensive aquatic resistance training program is effective at  
231 decreasing fat mass as well as improving walking speed in post-menopausal women with  
232 mild knee OA. This is the first randomised controlled study investigating the effects of  
233 aquatic resistance training on individuals with mild knee OA with a 12-month follow-up  
234 period. While, our results show that the improvements in body composition are lost at 12-  
235 months' follow-up, the improvements in walking speed were maintained. Importantly, higher  
236 average monthly LTPA was related with greater loss of fat mass over the 16-month study  
237 period. Further, this is the first study to report the actual training intensities achieved by the  
238 subjects during an aquatic exercise intervention and the effect of LTPA during the  
239 intervention and follow-up period.

240

241 Increased fat mass is linked to knee OA through biomechanical<sup>29</sup> and low-grade  
242 inflammatory mechanisms<sup>30</sup>, and is associated with an increased risk of suffering from knee  
243 OA as well as a more rapid progression of the disease<sup>31-33</sup>. A change of -1% in body weight  
244 has been shown to have a significant association with slower loss of tibial cartilage volume  
245 and improvement in symptoms suggesting our 1.4% weight change could have a meaningful  
246 impact on both cartilage health and OA related symptoms<sup>34</sup>. Our findings indicate a superior  
247 improvement in body composition compared with the two previous studies investigating the  
248 effects of aquatic exercise on body and fat mass in persons with OA<sup>17,35</sup>. The respective -  
249 1.17kg and -1.1kg decreases in fat and body mass evoked in our study are larger than the  
250 non-significant decrease in fat mass (-0.7kg) reported by Lim et al.<sup>17</sup> and significant  
251 reduction in body mass (-0.76kg) reported by Kim et al.<sup>35</sup>. While this could be due our  
252 slightly longer duration i.e. 4 weeks longer, we also utilised a much higher training

253 intensities. Lim et al.<sup>17</sup> set intensity at 65% Max HR, while Kim et al.<sup>35</sup> set intensity at RPE  
254 12-13 (Borg 6-20), approximately 60% Max HR<sup>36</sup>. In our study, average maximum heartrates  
255 during the main set were close to 85% with measured maximum HR up to 105%. Further, our  
256 finding demonstrate that high intensity aquatic exercise, appears to have similar effects as  
257 land-based exercise programs on body and fat mass. Messier et al.<sup>9</sup>, for example, reported  
258 that an 18 month, 3 times a week land-based exercise-only program produced a (-1.8kg) loss  
259 in total body weight . However, the decrease in fat mass was only 1% (-0.4 kg) and loss of  
260 lean mass was 1% (-2.6kg) in the exercise-only group. A loss of lean-mass and therefore  
261 reduction in muscle strength, is a common negative side effect of weight loss. Reduction in  
262 muscle mass and strength is associated with the development and faster progression of knee  
263 OA<sup>37,38</sup>, therefore preserving muscle mass during periods of weight loss is vital in this  
264 population<sup>33</sup>. Our study showed no change in lean mass and previously reported muscle  
265 strength<sup>18</sup>, indicating that while the training was intensive enough to evoke a decrease in fat  
266 mass it was also sufficient to preserve strength and lean mass.

267

268 Improvements in walking speed after 4-months aquatic resistance training, and its  
269 maintenance, after a 12-month follow-up period, are in contrast to the results for body  
270 composition changes. Slower walking speeds are associated with faster progression of OA  
271 related symptoms and activity limitations and the small but sustained improvement of  
272 0.05m/sec achieved in our study indicate a meaningful and lasting improvement in functional  
273 capacity<sup>39</sup>. Given that the weight lost during the intervention was regained during follow-up,  
274 our findings suggest that the effect on walking speed may not have been weight related. The  
275 results could indicate an improvement in cardiovascular fitness; however, the lack of between  
276 group differences in LTPA over the 12-months' follow-up period and at a level that had no  
277 effect on cardiovascular fitness in controls during the intervention period, suggests that this



278 alone cannot explain the maintenance of walking speed. While there were no improvements  
279 in muscle strength of the knee extensors and flexors, (previously reported<sup>18</sup>) we cannot rule  
280 out improvements in the un-measured ankle plantar flexors or hip abductors which could  
281 improve gait biomechanics and efficiency<sup>40</sup>. Further, strength alone is not a marker of  
282 improved gait biomechanics with efficient gait requiring co-ordination between agonist and  
283 antagonist muscles<sup>7, 41, 42</sup>. Immersion results in a decrease in nociceptor stimulation and  
284 afferent feedback<sup>43, 44</sup>, and reduces the sensation of pain<sup>13, 14, 45, 46</sup>. These conditions may  
285 create a suitable training condition for improving gait biomechanics<sup>14</sup>. Alternatively, the high  
286 intensity intervention exposed the subjects to the sensation of high physical exertion. This  
287 could have taught the participants that it was safe for them to exert themselves at a higher  
288 intensity than previously thought. It is feasible to speculate that this exercise pedagogy was  
289 retained 12-months after intervention cessation. However, walking frequency and intensity as  
290 part of the monthly LTPA did not differ between groups suggesting the improvement in  
291 walking speed was not utilised. Ultimately, the mechanisms behind the effect of aquatic  
292 resistance training on walking speed, deserves further investigation.

293

294 Education on life-style changes has been suggested as a vital part of management of both  
295 early and late-stage OA, in order to sustain improved levels of physical activity following an  
296 intervention study<sup>47</sup>. Participants in the training group did not have higher leisure time  
297 physical activity after the intervention than the control group therefore it is plausible to  
298 conclude that they returned back to pre-intervention level. Therefore, the increased walking  
299 speed may only describe improved functional capacity and may not be associated with  
300 increased walking speeds utilised in daily life. In combination with the possible exercise  
301 pedagogical effect of the high intensity exercise and implementation of a life-style education  
302 program, including dietary and advice, may have maintained or even continued the

303 improvements body composition and walking speed. Importantly, our results showed an  
304 association between higher levels LTPA and loss of fat mass irrespective of group allocation.  
305 Our results suggest that involvement in the intervention did not increase LTPA during the 12-  
306 months' follow-up period. However, LTPA was measured using self-reported questionnaires  
307 and it is plausible to hypothesise that after the intervention the participants in the training  
308 group may have changed their perception of activity intensities. Further, inclusion in a study  
309 may have caused a general increase in LTPA explaining the results. No acute worsening of  
310 clinical symptoms, as measured with the KOOS, was seen, possibility a result of the low  
311 impairment at baseline, the fluctuating nature of OA symptoms and the relatively short  
312 follow-up period<sup>2</sup>.

313

314 The strengths of this study included the randomised control design. The high adherence to the  
315 intervention and small number of drop-outs optimised the treatment response and shows  
316 motivation to participate in such an aquatic resistance exercise intervention. This is the first  
317 study to monitor leisure time physical activity, in addition to the exercise intervention, during  
318 an aquatic exercise intervention in participants with knee OA<sup>15</sup>, controlling an important  
319 confounding factor. The main limitation of this study was the use of strict inclusion criteria,  
320 essential for the original primary qMRI outcomes, which resulted in a homogeneous sample  
321 limiting direct application of our results to persons with more severe knee OA. However, it is  
322 conceivable to assume that, adapted, this program, would be suitable to improve functional  
323 capacity and decrease weight in subjects with more severe knee OA. Further studies are  
324 needed to confirm its efficacy in subjects with hip OA. The use of un-equal interventions i.e.  
325 only 2 sessions in the control group, introduces at least some degree of bias in favour of the  
326 intervention. Therefore, these results only indicate that aquatic resistance training is effective  
327 compared to no intervention and not more effective than another intervention. The lack of

328 assessor blinding to the intervention may have resulted in bias, however, assessors had no  
329 vested interest in the results of this study and primary investigator was blinded throughout.  
330 Dietary intake was not measured or controlled for. Inclusion in a study has been shown to  
331 affect participants' dietary habits as well as physical activity and therefore we cannot directly  
332 attribute all the changes as a pure effect of the intervention. Diet alone, however, would not  
333 have accounted for the maintenance of lean body mass<sup>9</sup>. Further, greater increases in lean  
334 mass and decreases in fat mass may have occurred with appropriate diet<sup>48</sup>. Intensity of the  
335 self-reported LTPA may have been affected after the intervention therefore use of objective  
336 measure after the intervention, e.g. accelerometers, would have given more accurate  
337 information<sup>49</sup>. It is not known if the mechanisms improving walking ability occurred earlier  
338 during the intervention therefore, future studies could look at the effectiveness of a shorter  
339 intensive aquatic exercise intervention.

340

### 341 **Conclusion**

342

343 To conclude, our findings show that a relatively short high intensity aquatic resistance  
344 training program decreases fat mass and improves walking speed in post-menopausal women  
345 with mild knee OA. Only improvements in walking speed were maintained at 12-months'  
346 follow-up. Further, LTPA appeared more important for controlling body composition than  
347 walking speed. Therefore, future research should investigate if lifestyle education following  
348 an intensive aquatic resistance training intervention optimises long term benefits for people  
349 with knee OA. Additionally, research is needed, to discover through which mechanism  
350 aquatic resistance training improves walking speed.

351 **Author contributions**

352 Waller, Benjamin: Analysis and interpretation of the data, drafting of the article, critical revision of  
353 the article for important intellectual content, final approval of the article, obtaining of funding,  
354 collection and assembly of data.

355 Munukka, Matti: Analysis and interpretation of the data, drafting of the article, critical revision of the  
356 article for important intellectual content, final approval of the article, obtaining of funding, collection  
357 and assembly of data.

358 Rantalainen, Timo: Analysis and interpretation of the data, critical revision of the article for important  
359 intellectual content, final approval of the article, administrative, technical, or logistic support,  
360 collection and assembly of data.

361 Lammentausta, Eveliina: Conception and design, analysis and interpretation of the data, critical  
362 revision of the article for important intellectual content, final approval of the article, administrative,  
363 technical, or logistic support.

364 Nieminen, Miika : Conception and design, analysis and interpretation of the data, critical revision of  
365 the article for important intellectual content, final approval of the article, administrative, technical, or  
366 logistic support.

367 Kiviranta, Ilkka: Conception and design, analysis and interpretation of the data, critical revision of the  
368 article for important intellectual content, final approval of the article.

369 Kautiainen, Hannu: Conception and design, analysis and interpretation of the data, drafting of the  
370 article, critical revision of the article for important intellectual content, final approval of the article,  
371 statistical expertise, collection and assembly of data.

372 Häkkinen, Arja: Conception and design, analysis and interpretation of the data, drafting of the article,  
373 critical revision of the article for important intellectual content, final approval of the article.

374 Kujala, Urho: Conception and design, analysis and interpretation of the data, drafting of the article,  
375 critical revision of the article for important intellectual content, final approval of the article.

376 Heinonen, Ari: Conception and design, analysis and interpretation of the data, drafting of the article,  
377 critical revision of the article for important intellectual content, final approval of the article, obtaining  
378 of funding.

379

### 380 **Role of funding sources**

381 Financial supporters were: Academy of Finland (ref: 253198), The Social Insurance  
382 Institution of Finland (KELA) (ref: 34/26/2011), Finnish Cultural Foundation and Yrjö  
383 Jahnsson foundation. None of these had a role in study design, collection, analysis and  
384 interpretation of data; in the writing of the manuscript; and in the decision to submit the  
385 manuscript for publication.

386

### 387 **Conflict of interest**

388 There is no conflict of interest for any authors.

389

## 390 References

- 391 1. McAlindon T, Bannuru R, Sullivan M, Arden N, Berenbaum F, Bierma-Zeinstra S, *et al.*  
392 OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis*  
393 *Cartilage* 2014;22:363-88.
- 394 2. Roos EM and Arden NK. Strategies for the prevention of knee osteoarthritis. *Nat Rev*  
395 *Rheumatol* 2016;12:92-101. doi: 10.1038/nrrheum.2015.135.
- 396 3. van Dijk GM, Veenhof C, Spreeuwenberg P, Coene N, Burger BJ, van Schaardenburg D,  
397 *et al.* Prognosis of limitations in activities in osteoarthritis of the hip or knee: a 3-year cohort  
398 study. *Arch Phys Med Rehabil* 2010;91:58-66. doi: 10.1016/j.apmr.2009.08.147.
- 399 4. Pisters MF, Veenhof C, van Dijk GM, Heymans MW, Twisk JW, Dekker J. The course of  
400 limitations in activities over 5 years in patients with knee and hip osteoarthritis with moderate  
401 functional limitations: risk factors for future functional decline. *Osteoarthritis Cartilage*  
402 2012;20:503-10. doi: 10.1016/j.joca.2012.02.002.
- 403 5. Murray IR, Benke MT, Mandelbaum BR. Management of knee articular cartilage injuries  
404 in athletes: chondroprotection, chondrofacilitation, and resurfacing. *Knee Surg Sports*  
405 *Traumatol Arthrosc* 2016;24:1617-26. doi: 10.1007/s00167-015-3509-8.
- 406 6. Mundermann A, Dyrby CO, Hurwitz DE, Sharma L, Andriacchi TP. Potential strategies to  
407 reduce medial compartment loading in patients with knee osteoarthritis of varying severity:  
408 reduced walking speed. *Arthritis Rheum.* 2004;50:1172-8. doi: 10.1002/art.20132.
- 409 7. Zeni JA, Jr and Higginson JS. Differences in gait parameters between healthy subjects and  
410 persons with moderate and severe knee osteoarthritis: a result of altered walking speed? *Clin*  
411 *Biomech* 2009;24:372-8. doi: 10.1016/j.clinbiomech.2009.02.001.

- 412 8. Fransen M, McConnell S, Harmer AR, van der Esch M, Simic M, Bennell KL. Exercise  
413 for osteoarthritis of the knee. *Br J Sports Med* 2015;10.1136/bjsports-2015-095424: doi:  
414 10.1002/14651858.CD004376.pub3.
- 415 9. Messier SP, Mihalko SL, Legault C, Miller GD, Nicklas BJ, DeVita P, *et al.* Effects of  
416 intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among  
417 overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial.  
418 *JAMA* 2013;310:1263-73. doi: 10.1001/jama.2013.277669; 10.1001/jama.2013.277669.
- 419 10. Hochberg MC, Altman RD, April KT, Benkhalti M, Guyatt G, McGowan J, *et al.*  
420 American College of Rheumatology 2012 recommendations for the use of nonpharmacologic  
421 and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res*  
422 2012;64:465-74.
- 423 11. Holla JF, Sanchez-Ramirez DC, van der Leeden M, Ket JC, Roorda LD, Lems WF, *et al.*  
424 The avoidance model in knee and hip osteoarthritis: a systematic review of the evidence. *J*  
425 *Behav Med* 2014;37:1226-41. doi: 10.1007/s10865-014-9571-8.
- 426 12. Harrison RA, Hillman M, Bulstrode S. Loading of the lower limb when walking partially  
427 immersed: implications for clinical practice. *Physiotherapy* 1992;78:164-6.
- 428 13. Denning WM, Bressel E, Dolny DG. Underwater treadmill exercise as a potential  
429 treatment for adults with osteoarthritis. *International Journal of Aquatic Research and*  
430 *Education* 2010;4:70-80.
- 431 14. Roper JA, Bressel E, Tillman MD. Acute aquatic treadmill exercise improves gait and  
432 pain in people with knee osteoarthritis. *Arch Phys Med Rehabil* 2013;94:419-25. doi:  
433 10.1016/j.apmr.2012.10.027.

- 434 15. Waller B, Ogonowska-Słodownik A, Vitor M, Lambeck J, Daly D, Kujala UM, *et al.*  
435 Effect of therapeutic aquatic exercise on symptoms and function associated with lower limb  
436 osteoarthritis: systematic review with meta-analysis. *Phys Ther* 2014;94:1383-95. doi:  
437 10.2522/ptj.20130417.
- 438 16. Waller B, Ogonowska-Słodownik A, Vitor M, Rodionova K, Lambeck J, Heinonen A, *et*  
439 *al.* The effect of aquatic exercise on physical functioning in the older adult: a systematic  
440 review with meta-analysis. *Age Ageing* 2016;doi: 10.1093/ageing/afw102.
- 441 17. Lim JY, Tchai E, Jang SN. Effectiveness of aquatic exercise for obese patients with knee  
442 osteoarthritis: a randomized controlled trial. *PM R* 2010;2:723-31.
- 443 18. Munukka M, Waller B, Rantalainen T, Hakkinen A, Nieminen MT, Lammentausta E, *et*  
444 *al.* Efficacy of progressive aquatic resistance training for tibiofemoral cartilage in  
445 postmenopausal women with mild knee osteoarthritis: a randomised controlled trial.  
446 *Osteoarthritis Cartilage* 2016;24(10):1708-17. doi: 10.1016/j.joca.2016.05.007.
- 447 19. Waller B, Munukka M, Multanen J, Rantalainen T, Poyhonen T, Nieminen MT, *et al.*  
448 Effects of a progressive aquatic resistance exercise program on the biochemical composition  
449 and morphology of cartilage in women with mild knee osteoarthritis: protocol for a  
450 randomised controlled trial. *BMC Musculoskelet.Disord.* 2013;14:82,2474-14-82. doi:  
451 10.1186/1471-2474-14-82.
- 452 20. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, *et al.* Development of  
453 criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of  
454 the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism  
455 Association. *Arthritis Rheum* 1986;29:1039-49.



- 456 21. Moher D, Hopewell S, Schulz KF, Montori V, Gotzsche PC, Devereaux PJ, *et al.*  
457 CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group  
458 randomised trials. *Int J Surg* 2012;10:28-55. doi: 10.1016/j.ijisu.2011.10.001;  
459 10.1016/j.ijisu.2011.10.001.
- 460 22. Uusi-Rasi K, Rauhio A, Kannus P, Pasanen M, Kukkonen-Harjula K, Fogelholm M, *et al.*  
461 Three-month weight reduction does not compromise bone strength in obese premenopausal  
462 women. *Bone* 2010;46:1286-93. doi: 10.1016/j.bone.2009.10.013.
- 463 23. Laukkanen R, Oja R, Pasanen M, Vuori I. Criterion validity of a two-kilometer walking  
464 test for predicting the maximal oxygen uptake of moderately to highly active middle-aged  
465 adults. *Scand J Med Sci Sports* 1993;3:267-72.
- 466 24. Koli J, Multanen J, Häkkinen A, Kiviranta I, Kujala U, Heinonen A. Reliability of the  
467 Finnish versions of WOMAC and KOOS forms for knee osteoarthritis. *Physiotherapy*  
468 2011;97:D203-4.
- 469 25. Bekkers JE, de Windt TS, Raijmakers NJ, Dhert WJ, Saris DB. Validation of the Knee  
470 Injury and Osteoarthritis Outcome Score (KOOS) for the treatment of focal cartilage lesions.  
471 *Osteoarthritis Cartilage* 2009;17:1434-9.
- 472 26. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Jr DRB, Tudor-Locke C, *et al.*  
473 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med*  
474 *Sci Sports Exerc* 2011;43:1575-81.
- 475 27. Cohen J. A power primer. *Psychol Bull* 1992;112:155-9.
- 476 28. Cohen J. *Statistical power analysis for the behavioral sciences*. Hillsdale, N.J. : Lawrence  
477 Erlbaum, 1988.

- 478 29. Browning RC and Kram R. Effects of obesity on the biomechanics of walking at different  
479 speeds. *Med Sci Sports Exerc* 2007;39:1632-41. doi: 10.1249/mss.0b013e318076b54b [doi].
- 480 30. Vincent HK, Heywood K, Connelley J, Hurley RW. Weight loss and obesity in the  
481 treatment and prevention of osteoarthritis. *PM R*. 2012;4:S59-67. doi:  
482 10.1016/j.pmrj.2012.01.005.
- 483 31. Houard X, Goldring MB, Berenbaum F. Homeostatic mechanisms in articular cartilage  
484 and role of inflammation in osteoarthritis. *Curr Rheumatol Rep* 2013;15:375,013-0375-6. doi:  
485 10.1007/s11926-013-0375-6.
- 486 32. Vuolteenaho K, Koskinen A, Moilanen E. Leptin - a link between obesity and  
487 osteoarthritis. applications for prevention and treatment. *Basic Clin Pharmacol Toxicol*  
488 2014;114:103-8. doi: 10.1111/bcpt.12160.
- 489 33. Ding C, Stannus O, Cicuttini F, Antony B, Jones G. Body fat is associated with increased  
490 and lean mass with decreased knee cartilage loss in older adults: a prospective cohort study.  
491 *Int J Obes* 2013;37:822-7.
- 492 34. Teichtahl AJ, Wluka AE, Tanamas SK, Wang Y, Strauss BJ, Proietto J, *et al*. Weight  
493 change and change in tibial cartilage volume and symptoms in obese adults. *Ann Rheum Dis*  
494 2015;74:1024-9. doi: 10.1136/annrheumdis-2013-204488.
- 495 35. Kim IS, Chung SH, Park YJ, Kang HY. The effectiveness of an aquarobic exercise  
496 program for patients with osteoarthritis. *Appl Nurs Res* 2012;25:181-9. doi:  
497 10.1016/j.apnr.2010.10.001.
- 498 36. Whaley MH, Brubaker PH, Otto RM, Eds. *ACSM's Guidelines for Exercise Testing and*  
499 *Prescription*. Philadelphia: Lippin cott Williams and Wilkins, 2006.

- 500 37. van der Esch M, Holla JF, van der Leeden M, Knol DL, Lems WF, Roorda LD, *et al.*  
501 Decrease of muscle strength is associated with increase of activity limitations in early knee  
502 osteoarthritis: 3-year results from the cohort hip and cohort knee study. *Arch Phys Med*  
503 *Rehabil* 2014;95:1962-8. doi: 10.1016/j.apmr.2014.06.007.
- 504 38. Oiestad BE, Juhl CB, Eitzen I, Thorlund JB. Knee extensor muscle weakness is a risk  
505 factor for development of knee osteoarthritis. A systematic review and meta-analysis.  
506 *Osteoarthritis Cartilage* 2015;23:171-7. doi: 10.1016/j.joca.2014.10.008.
- 507 39. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and  
508 responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc*  
509 2006;54:743-9.
- 510 40. Kulmala JP, Korhonen MT, Kuitunen S, Suominen H, Heinonen A, Mikkola A, *et al.*  
511 Which muscles compromise human locomotor performance with age? *J R Soc Interface*  
512 2014;11:20140858. doi: 10.1098/rsif.2014.0858.
- 513 41. Mills K, Hettinga BA, Pohl MB, Ferber R. Between-limb kinematic asymmetry during  
514 gait in unilateral and bilateral mild to moderate knee osteoarthritis. *Arch Phys Med Rehabil.*  
515 2013;94:2241-7. doi: 10.1016/j.apmr.2013.05.010.
- 516 42. Mills K, Hunt MA, Leigh R, Ferber R. A systematic review and meta-analysis of lower  
517 limb neuromuscular alterations associated with knee osteoarthritis during level walking. *Clin*  
518 *Biomech* 2013;28:713-24. doi: 10.1016/j.clinbiomech.2013.07.008.
- 519 43. Cronin NJ, Valtonen AM, Waller B, Poyhonen T, Avela J. Effects of short term water  
520 immersion on peripheral reflex excitability in hemiplegic and healthy individuals: A  
521 preliminary study. *J Musculoskelet Neuronal Interact* 2016;16:58-62.

- 522 44. Pöyhönen T and Avela J. Effect of head-out water immersion on neuromuscular function  
523 of the plantarflexor muscles. *Aviat Space Environ Med* 2002;73:1215-8.
- 524 45. Hall J, Swinkels A, Briddon J, McCabe CS. Does aquatic exercise relieve pain in adults  
525 with neurologic or musculoskeletal disease? A systematic review and meta-analysis of  
526 randomized controlled trials. *Arch Phys Med Rehabil* 2008;89:873-83. doi:  
527 10.1016/j.apmr.2007.09.054.
- 528 46. Fiskén A, Waters DL, Hing WA, Steele M, Keogh JW. Perception and Responses to  
529 Different Forms of Aqua-Based Exercise Among Older Adults With Osteoarthritis.  
530 *International Journal of Aquatic Research & Education* 2014;8:32-52.
- 531 47. Filardo G, Kon E, Longo UG, Madry H, Marchettini P, Marmotti A, *et al.* Non-surgical  
532 treatments for the management of early osteoarthritis. *Knee Surg.Sports Traumatol Arthrosc.*  
533 2016;26:1775-85. doi: 10.1007/s00167-016-4089-y.
- 534 48. Thomas DM, Bouchard C, Church T, Slentz C, Kraus WE, Redman LM, *et al.* Why do  
535 individuals not lose more weight from an exercise intervention at a defined dose? An energy  
536 balance analysis. *Obes Rev* 2012;13:835-47. doi: 10.1111/j.1467-789X.2012.01012.x.
- 537 49. Ainsworth BE, Caspersen CJ, Matthews CE, Masse LC, Baranowski T, Zhu W.  
538 Recommendations to improve the accuracy of estimates of physical activity derived from self  
539 report. *J Phys Act Health* 2012;9 Suppl 1:S76-84.

540

541 **TABLES****Table 1** Baseline demographic and clinical characteristics

	Exercise group (n=43)	Control group (n=44)
Age (years)	63.8 (2.4)	63.9 (2.4)
Height (cm)	161.7 (5)	161.6 (5)
Body mass (kg)	69.6 (10.3)	71.0 (11.2)
Body mass index (kg/m <sup>2</sup> )	26.6 (3.8)	27.1 (3.5)
Affected knee (right/left)	36/7	34/10
K/L grade, n (%)		
Grade 1	23 (53.5)	24 (54.5)
Grade 2	20 (46.5)	20 (45.5)
Analgesia, n (%)*	11 (26)	9 (20)
LTPA (METh/week)	29 (31)	36 (33)
Smoker n (%)		
Never	17	13
Current	3	3
Previous	23	28
Blood pressure, n (%)		
Normal	23	14
Elevated	9	11
Medical management	11	19

542 Values are means (SD) unless otherwise noted.

543 \*Number of participants using analgesia for knee pain on inclusion to study

544 LTPA = leisure time physical activity, METh = metabolic equivalent task hour.

545

546 **Table 2** Description of training intensities achieved during the aquatic resistance training and  
547 psychological feelings experienced per progression

	Barefoot	Small Fins	Large boots
No. Sessions	8	14	26
RPE*	13.7 (1.0)	14.9 (1.3)	15.0 (1.5) <sup>†</sup>
Average HR (%)	61 (5.9)	61 (5.3)	61 (6.3)
Max HR (%)	85 (7.8)	84 (8.9)	84 (8.0)
Blood lactates (mmol/L) <sup>‡</sup>	4.9 (2.1)	4.5 (1.9)	4.0 (1.8)
Repetitions per session	481 (66)	408 (71)	376 (65) <sup>†</sup>
Self-reported emotional state	4.2 (0.33)	4.2 (0.36)	4.3 (0.40)

548 Mean and (SD) unless otherwise stated. \*RPE = Rating of perceived exertion (BORG 6-20)

549 <sup>‡</sup> Measured directly after sessions 35-37, <sup>†</sup>Bare vs Large (p<0.001)

550

551 **Table 3** Effect of aquatic resistance training on walking speed, body composition and clinical symptoms.

552

Variable	n	Aquatic training (AT)		Control		Mean Difference (95% CI)	p-value*	n	AT	Control	Mean Difference (95% CI)	p-value**
		BL mean (SD)	FU mean (SD)	BL mean (SD)	FU mean (SD)				12-FU mean (SD)	12-FU mean (SD)		
Walking speed (m/sec)	87	1.74 (0.15)	1.83 (0.16)	1.73 (0.17)	1.76 (0.17)	0.052 (0.018 to 0.086)	0.002	73	1.82 (0.14)	1.77 (0.13)	0.046 (0.006 to 0.90)	0.032
<b>Body Composition</b>												
Body Mass (Kg)	87	69.2 (10.3)	68.2 (10.4)	70.8 (11.2)	70.9 (11.3)	-1.11 (-1.85 to -0.42)	0.004	76	68.6 (10.6)	70.8 (11.5)	-0.39 (-1.51 to 0.64)	0.543
BMI	87	26.6 (3.8)	26.2 (3.9)	27.1 (3.5)	27.1 (3.6)	-0.46 (-0.74 to -0.19)	0.001	76	26.4 (4.0)	26.9 (3.7)	0.001 (-0.47 to 0.47)	0.892
Lean Mass (kg)	87	40.3 (3.9)	40.6 (3.9)	41.4 (4.4)	41.7 (4.4)	0.083 (-0.29 to 0.45)	0.590	76	40.1 (4.0)	41.9 (4.2)	-0.30 (-0.79 to 0.12)	0.410
Fat Mass (Kg)	87	26.0 (8.6)	24.8 (8.8)	26.5 (8.0)	26.4 (8.1)	-1.17 (-2.00 to -0.43)	0.002	76	25.7 (8.8)	26.1 (8.5)	-0.14 (-1.24 to 0.90)	0.700
<b>KOOS (0-100)</b>												
Pain	87	80.6 (10.4)	84.3 (10.5)	82.1 (11.8)	83.3 (11.7)	2.3 (-1.93 to 6.31)	0.184	76	86.8 (10.5)	85.1 (12.4)	1.45 (-2.72 to 5.66)	0.187
Symptoms	87	74.4 (12.9)	80.9 (12.1)	74.8 (14.1)	77.5 (14.9)	4.07 (-0.43 to 8.54)	0.091	76	81.4 (11.4)	77.9 (14.5)	3.31 (-1.19 to 7.30)	0.119
ADL	87	84.5 (10.4)	87.7 (9.7)	85.2 (11.0)	86.0 (14.6)	3.36 (-0.38 to 7.118)	0.105	74	89.2 (11.2)	88.3 (11.0)	0.97 (-2.64 to 4.32)	0.397
Sport&Rec	87	63.6 (20.5)	70.6 (21.7)	64.8 (22.2)	67.6 (26.5)	4.81 (-3.00 to 12.61)	0.223	76	71.0 (20.7)	68.7 (24.6)	2.45 (-4.76 to 8.96)	0.396
QoL	87	66.0 (17.5)	72.6 (18.1)	70.6 (20.1)	74.1 (23.1)	2.76 (-3.51 to 8.66)	0.248	75	75.0 (18.2)	76.4 (24.4)	1.21 (-5.97 to 7.98)	0.308
LTPA (MET/h)	85	-	160 (53)	-	104 (63)	56 (-81.4 to -31.1)	<0.001†	76	100 (57)	107 (56)	-7.3 (-31.5 to 16.9)	0.56†

553 BL = baseline, FU = post-intervention (4-months), 12-FU = 12-months follow-up, \*post-intervention follow-up compared to baseline (mixed

554 model), \*\*12-month follow-up compared to baseline (mixed-model),

555 LTPA = average monthly total Leisure Time Physical Activity, MET/h = metabolic equivalent task hour. †Fisher-Pitman permutation test

556

557 **FIGURE LEGENDS**558 **Figure 1** Flow chart showing participant recruitment, randomisation and retention559 **Figure 2** Changes in A) fat and lean mass (kg) and B) walking speed (m/sec) following a 4-  
560 months aquatic resistance training and 12-month follow-up period.561 **Figure 3** Monthly leisure time physical activity (METH)

