

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Ng, Kwok; Tynjälä, Jorma; Sigmundová, Dagmar; Augustine, Lilly; Sentenac, Mariane; Rintala, Pauli; Inchley, Jo

Title: Physical activity among adolescents with long-term illnesses or disabilities in 15 European countries

Year: 2017

Version:

Please cite the original version:

Ng, K., Tynjälä, J., Sigmundová, D., Augustine, L., Sentenac, M., Rintala, P., & Inchley, J. (2017). Physical activity among adolescents with long-term illnesses or disabilities in 15 European countries. *Adapted Physical Activity Quarterly*, 34(4), 456-465.
<https://doi.org/10.1123/apaq.2016-0138>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Physical activity among adolescents with long term illnesses or disabilities in 15

European countries

Kwok Ng, PhD. Faculty of Sport and Health Sciences, University of Jyväskylä, Finland.

Kwok.ng@jyu.fi

Jorma Tynjälä, PhD. Faculty of Sport and Health Sciences, University of Jyväskylä, Finland.

Jorma.tynjala@jyu.fi

Dagmar Sigmundova, PhD. Faculty of Physical Culture, Palacky University, Czech Republic.

Dagmar.sigmundova@upol.cz

Lilly Augustine, PhD. Kristianstad university, Dept. Human sciences. Sweden.

lilly.augustine@hkr.se

Mariane Sentenac, PhD. INSERM, U.1027, Toulouse, F-31000 France ; Paul-Sabatier

University, Toulouse, F-31400 France. marianesentenac@yahoo.fr

Pauli Rintala, PhD. Faculty of Sport and Health Sciences, University of Jyväskylä, Finland.

Jo Inchley, PhD. University of St Andrews, Scotland. jci2@st-andrews.ac.uk

Corresponding author: Kwok Ng, kwok.ng@jyu.fi

1 Physical activity among adolescents with long-term illnesses or disabilities in 15
2 European countries

3 **Abstract:**

4 Physical activity (PA) is an important health-promoting behaviour that adolescents
5 with long-term illnesses or disabilities (LTID) can benefit from. It is important to monitor
6 differences across countries in adherence with PA recommendations for health (PARH). The
7 aim of this study was to compare PA levels among 15 European countries after
8 disaggregating data by disability. Data from pupils (Mean age = 13.6y, SD=1.64)
9 participating in the 2013/14 Health Behaviour in School-aged Children (HBSC) study were
10 analysed to compare adolescents without LTID, with LTID and with LTID that affects their
11 participation (affected LTID). Logistic regression models adjusted for age and family
12 affluence, stratified by gender and country group with PARH as the outcome variable. With
13 the data pooled, 15% (n=9,372) of adolescents reported to have LTID and 4% (n=2566) with
14 affected LTID. Overall, fewer boys with LTID met PARH than boys without LTID, although
15 it was not statistically significant at the national levels nor for girls.

16 Keywords: inclusion; mainstream schools; health; chronic conditions, disability

1 **Introduction**

2 A mass of evidence exists confirming the health benefits from regular physical
3 activity (PA) among adolescents (Poitras et al., 2016). PA is defined as, “any bodily
4 movement produced by skeletal muscles that requires energy expenditure” (Caspersen,
5 Powell, & Christenson, 1985, p. 126) and moderate intensity PA (usually above 3METs) is
6 considered as health-enhancing PA (Biddle, Sallis, & Cavill, 1998). International PA
7 recommendations for health (PARH) for children and adolescents aged 5-17 years old is set
8 to participation of at least 60 minutes of moderate to vigorous-intensity physical activity daily
9 (WHO, 2010). It is also important for youth with disabilities to engage in PA, and meet the
10 PARH since regular participation in PA distinctly reduces health complications secondary to
11 disability conditions (Rimmer, Schiller, Chen, Schiller, & Chen, 2012).

12 According to the International Classification of Functioning, Disability and Health
13 (ICF; WHO, 2001), people may experience disability because of the interaction between
14 health conditions and contextual factors, thus influencing the child’s ability to participate in
15 various activities including PA. Common barriers to participation in PA reported by people
16 with disabilities include attitudes of others, lack of friends, high costs, low self-determination,
17 fatigue, as well as accessibility (Ross et al., 2016). The degree to which these barriers exist to
18 influence levels of PA in adolescents with different impairments may vary, as with
19 adolescents with other chronic health conditions diagnosed by the doctor (Ng, Rintala,
20 Tynjälä, Villberg, & Kannas, 2014). Yet, it is difficult to draw conclusions about these
21 differences when studies use different protocols, instruments, and questions to measure PA.
22 In addition, only six percent of studies published in *Adapted Physical Activity Quarterly*
23 between 2004 and 2013 came from multiple samples, rather than subtypes of disabilities
24 (Haegele, Lee, & Porretta, 2015). The high possibility of conditions and impairments co-

25 existing in adolescents, makes increasing PA among young people with a disability an
26 important public health concern (Ding et al., 2016).

27 Public health and health promotion experts often use monitoring and surveillance
28 studies to assess progress over time between countries and for global evaluations (Brown,
29 Cueto, & Fee, 2006). Global coverage of PA surveillance has increased from 64% of the
30 European adolescent population in 2012 to 68% in 2016 (Sallis et al., 2016). However, it is
31 unknown what proportion, if any, of these surveys involve people with disabilities. The
32 Active Healthy Kids Global Alliance has recognised this in their most recent report card
33 (Tremblay et al., 2016) which presented population-level data from 38 countries but
34 highlighted the need for evidence concerning PA among children with disabilities. These
35 suggestions are in line with the statements in the 2030 sustainable developmental goals that
36 support reporting disaggregation by disability (Tardi & Njelesani, 2015).

37 In the most recent PA progress report by Sallis and colleagues (Sallis et al., 2016),
38 adolescents that failed to meet the PARH (WHO, 2010) were considered as “inactive”. In
39 other words, for the purpose of surveillance, only adolescents who achieved at least 60
40 minutes of moderate to vigorous physical activity on a daily basis were considered as active
41 adolescents. Although this benchmark does not allow for consideration of overall differences
42 in PA levels, it serves the purpose for analysis against the PARH which is of primary interest
43 to policy makers for national and international monitoring (Kalman et al., 2015). Due to the
44 absence of data on adolescents with disabilities in existing reports, it is of great importance to
45 report PA levels after disaggregation for disability. Furthermore, reporting of PA against
46 benchmarks used in global surveys is necessary for making comparisons of young people
47 with and without disabilities. Therefore, the purpose of this study is to use the PARH as a
48 reference point to compare prevalence of physically active adolescents across Europe after
49 disaggregating for disability and adjusting for age and family affluence.

50 **Method**

51 Data were analysed from the 2013/14 WHO Collaborative Cross-national Health
52 Behaviour in School-aged Children (HBSC) study. The 15 countries in the current analysis
53 included the same questions on disabilities. All countries participating in HBSC comply with
54 a standardised international protocol, including back translation of items. The samples were
55 at the class level to form nationally representative estimates of 11-, 13- and 15-year-olds.
56 Each team obtained approval to conduct the study through an ethics review board or
57 equivalent regulatory body. The school administrators, parents and pupils granted consent
58 (explicit or implicit, varied by country) for pupil completion of a self-report survey in the
59 classroom without further assistance. Responses from the participants were anonymous, and
60 participation was voluntary. Response rates varied by country, although was >70% at the
61 international student level, with a proportion of the non-responses from absentees.

62 The pupils were asked to report their sex, month and year of birth as well as family
63 affluence, measured using the Family Affluence Scale (FAS III). FAS III is a six-item
64 assessment of material assets or activities and regarded as a child friendly indicator of social
65 economic status (see Torsheim et al., 2016). FAS III was divided into three groups, low FAS
66 (as represented by the lowest 20 percentile), medium FAS (the middle 60 percentile), and
67 high FAS (highest 20 percentile). A separate analysis was performed for Armenia, since four
68 (FAS II) of the six FAS III items reported.

69 **LTID status**

70 Two items were used to group adolescents according to their long-term illness or
71 disability (LTID) status. Two yes or no questions were asked: 1) “Do you have a long-term
72 illness, disability, or medical condition (like diabetes, arthritis, allergy, or cerebral palsy) that
73 has been diagnosed by a doctor?” and 2) “Does your long-term illness or disability affect

74 your attendance and participation at school?” Students were classified into three mutually
75 exclusive categories, i) adolescents without LTID, ii) adolescents for whom the participation
76 is not affected by their LTID – grouped as LTID, and iii) adolescents with participation
77 affected by the LTID – grouped as affected LTID. Guided by the International Classification
78 of Functioning, Disability and Health (ICF) definition of disability, the affected LTID group
79 was classified as a group with greater severity of disability to the group of LTID. Data from
80 Bulgaria and Czech Republic consisted of only the first question. The Finnish team collected
81 data on this item only from 13 and 15 year-olds.

82 **Physical activity measures**

83 A single item assessed the number of days the pupil participated in moderate to
84 vigorous physical activity (MVPA) frequency of at least 60 minutes during the last 7 days
85 and was used to measure adherence to the PARH. Text to define PA was included and the
86 question was as follows

87 *Physical activity is any activity that increases your heart rate and makes you get out*
88 *of breath some of the time. Physical activity can be done in sports, school activities, playing*
89 *with friends, or walking to school. Some examples of physical activity are running, brisk*
90 *walking, rollerblading, biking, dancing, skateboarding, swimming, soccer, basketball,*
91 *football, & surfing [country specific examples can be given].*

92 *Over the past 7 days, on how many days were you physically active for a total of at*
93 *least 60 minutes per day?*

94 *Please **add up** all the time you spent in physical activity each day.*

95 Response categories ranged from 0 day to 7 days. Results from validity studies
96 include correlation coefficient of 0.40 when compared with accelerometer data in clinical
97 settings (Murphy, Rowe, Belton, & Woods, 2015; Prochaska, Sallis, & Long, 2001) and the

98 item has been in use in the HBSC study since 2001/2002 survey. The ICC value from a test-
99 retest was 0.82 (Liu et al., 2010). In accordance with the PARH (WHO, 2010), and the
100 definitions by the PA progress report (Sallis et al., 2016), responses were dichotomised into
101 0-6 days as “inactive”, in other words, not meeting the PARH, and 7 days as “active”, as
102 meeting the PARH.

103 **Statistics**

104 Descriptive statistics were conducted on the data stratified by gender and country.
105 Missing cases from the LTID status were grouped and tested independently against PARH.
106 Since differences were $p < 0.05$, the cases were removed. Differences in prevalence of LTID
107 between boys and girls were tested using a chi-square test of independence (Table 1). Binary
108 logistic regression analyses were performed separately by country, with PARH as the
109 outcome variable and LTID status as the main independent variable. Reporting of odds ratios
110 (OR), with the 95% confidence intervals, first by the overall sample, then for each country,
111 were performed. The reference group was those without LTID. All analyses were conducted
112 after controlling for age categories and family affluence. Analyses were done with SPSS for
113 Windows (version 24.0)

114 **Results**

115 **LTID prevalence**

116 The pooled sample size was 61,329 participants (48.7% boys, mean age=13.6yr
117 SD=1.7) from the 15 countries (see Table 1 for the list of countries). Less than one in six
118 (15.3%, n=9,372) of adolescents reported to have LTID. One in twenty (4.2%, n=2566)
119 reported that their disability affects their participation at school and were therefore classified
120 as affected LTID. The proportions of affected LTID increased from 11-y (3.4%; 95% CI:
121 3.1%-3.6%), 13-y (4.2%; 95% CI: 3.9%-4.5%), to 15-y olds (4.9%; 95% CI: 4.6%-5.2%).

HEADER: PA among adolescents with LTID in Europe

122 Gender differences were only significant in some countries, whereby there were more boys
123 than girls with LTID or affected LTID (Armenia, Scotland and Ireland), or the prevalence
124 was greater amongst girls than boys (Bulgaria, Romania). There were no significant
125 differences in the prevalence across the different FAS groups.

126 **PA recommendations for health**

127 After pooling data from all countries, more boys (24.4%; 95% CI: 23.9%-24.9%) met
128 the PARH than girls (15.5%; 95% CI:15.1%-15.9%), more 11y- (25.2%; CI:24.5%-25.8%),
129 and 13y – (19.8%; CI:19.3%-20.3%) met the PARH than 15y –olds (14.8%; CI: 14.3%-
130 15.3%). In addition, more adolescents from the high FAS group met the PARH (24.3%;
131 CI=23.5%-25.1%) than medium FAS (18.7%; CI:18.3%-19.1%) and low FAS (18.0%;
132 CI:17.3%-18.8%). The patterns between age, gender and FAS III were similar across
133 disability groups.

134 Overall, boys with LTID were less likely to meet the recommendations than boys
135 without LTID, although differences amongst girls were not significant. At country level, the
136 proportions of boys with and without LTID meeting PARH were not significantly different.
137 In one country (Slovakia), a significant association was observed between affected LTID and
138 meeting the PARH amongst both boys and girls with OR 2.1 (CI: 1.1-4.2) and OR=1.9
139 (CI:=1.0-3.6) respectively. In Romania, the likelihood of meeting PARH increased in girls
140 with LTID with OR=1.8 (CI=1.1-3.1) and with affected LTID with OR=3.6 (CI=1.7-7.9)
141 when compared with girls without LTID (Table 2).

142 **Discussion**

143 Through the same measures reported in the 2016 PA progress report, data in this
144 study were disaggregated by disability across 15 countries to report proportions of
145 adolescents who meet PARH. Consistent with the previous reports, meeting the PARH was

HEADER: PA among adolescents with LTID in Europe

146 more common among boys, younger adolescents, and those from higher affluence families
147 (Kalman et al., 2015). According to the results from this study, there were no significant
148 differences in meeting PARH between boys or girls with LTID and those without LTID at a
149 national level, with two exceptions (Romania and Slovakia). After the data were pooled
150 together, boys with LTID were less likely to meet the PARH than their peers without LTID,
151 but this effect was not observed for girls.

152 The results of this study add to the current literature on adolescents that met PARH by
153 providing data disaggregated by disability on adolescents in mainstream schools, and using a
154 comparable measure of PA across countries. Previous studies generally report lower levels of
155 PA in adolescents with disabilities compared to their peers without disabilities (Ross et al.,
156 2016). In this study, this finding was only observed among boys and after pooling the data
157 across countries. This may be attributed to the fact that previous studies have often focused
158 on specific disability subcategories (Haegele et al., 2015) whereas the current study included
159 data from general schools with multiple samples of various (self-reported) impairments and
160 health conditions. Although overall, 20% of adolescents reported disabilities or chronic
161 conditions, less than one in twenty said that their disability affected their daily functioning at
162 school. As such, the majority of adolescents with LTID reported their conditions were
163 independent of their ability to participate in school. In addition, this study indicates that the
164 restrictions felt by adolescents in the affected LTID did not influence meeting PARH when
165 compared with their peers. Since overall proportions that met the PARH were low in all
166 countries, it is clear that there needs to be vast improvements in PA levels in all children,
167 regardless of disability. Practitioners with access to adolescents in mainstream schools may
168 need to consider the increasing rate of inclusion in schools. Furthermore, practitioners are
169 encouraged to promote PA to girls, and take care in addressing the divide in PA levels among
170 boys with and without LTID.

171 Europe is a culturally diverse area with inclusive practices in Central and Eastern
172 European countries different from Western European countries (UNICEF, 2012). Despite
173 such differences, significant differences at the national level were found in only two
174 countries. In Slovakia (boys and girls) and Romania (girls only), those with LTID or affected
175 LTID were more likely to meet the PARH. Cultural differences in educational policies,
176 access to support services and availability of physical activity opportunities may play a role.
177 For example, both are Central and Eastern European countries and segregated school systems
178 are still commonplace for children with specific needs. This may mean adolescents with
179 disabilities who attend mainstream schools are less likely to have severe disabilities, and
180 those that do, can benefit the most with the existing support services (UNICEF, 2012). These
181 differences may explain the unexpected increased proportion of affected LTID that met the
182 PARH over adolescents without LTID

183 The findings from this study were dependent on the way adolescents reported LTID
184 and PA as well as responded to the questions unaided. As such, there are some study
185 limitations to consider. PA was measured by self-report and there could be inaccuracies to the
186 response. However, self-report surveys are considered an appropriate way to collect data
187 from large population samples on PA for the purposes of meeting PARH (Haskell, 2012) and
188 all young people participating in this study were asked the same questions in the same way.
189 In relation to reporting on levels of physical activity, the practice used in this study was
190 replicated from Sallis and colleagues (2016) which considered young people as active only if
191 they achieved at least 60 minutes of MPVA daily in line with current international policy
192 recommendations. However, this may inadvertently mask differences between people (e.g. 0
193 days vs 2 days vs 6 days) that could be illuminating and should be investigated further in
194 future research.

195 In relation to the question used to identify those with a long-term illness or disability,
196 it should be noted that disability is not synonymous with being ill, although they often co-
197 exist. Disabilities were measured at two levels and not by impairment types. The first level
198 was the type of disability that the individual reports. This could be a manageable health
199 condition, such as asthma or speaking impairments, which might not have a major effect on
200 physical activity participation. The second level is the type of disability whereby the
201 disability or health condition affects daily participation. It remains unknown what context,
202 other than of school participation this was, but in reference to the ICF (WHO, 2001), these
203 pupils would be expected to have more difficulties in daily functioning. In addition, to
204 improve the global PA matrix, disability statistics are needed to confirm a systematic method
205 to compare data (Tremblay et al., 2016). Despite the study limitations, this paper has results
206 that can be used for international comparisons for PARH and contribute to data sets like the
207 global matrix.

208 **References**

- 209 Biddle, S. J. H., Sallis, J. F., & Cavill, N. (1998). *Youth and Active? Young people and*
210 *health-enhancing physical activity - evidence and implications*. London: Health
211 Education Authority. Retrieved from
212 <http://www.nice.org.uk/niceMedia/documents/youngandactive.pdf>
- 213 Brown, T. M., Cueto, M., & Fee, E. (2006). The World Health Organization and the
214 transition from international to global public health. *American Journal of Public Health,*
215 *96*, 62–72. <http://doi.org/10.2105/AJPH.2004.050831>
- 216 Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and
217 physical fitness: definitions and distinctions for health-related research. *Public Health*
218 *Reports, 100*, 126–31. doi:10.2307/20056429
- 219 Ding, D., Lawson, K. D., Kolbe-Alexander, T. L., Finkelstein, E. A., Katzmarzyk, P. T., van

- 220 Mechelen, W., & Pratt, M. (2016). The economic burden of physical inactivity: A global
221 analysis of major non-communicable diseases. *The Lancet*, *6736*, 1–14.
222 doi:10.1016/S0140-6736(16)30383-X
- 223 Haegele, J. A., Lee, J., & Porretta, D. L. (2015). Research trends in adapted physical activity
224 quarterly from 2004 to 2013. *Adapted Physical Activity Quarterly*, *32*(3), 187–205.
225 doi:10.1123/APAQ.2014-0211
- 226 Haskell, W. L. (2012). Physical activity by self-report: a brief history and future issues.
227 *Journal of Physical Activity & Health*, *9 Suppl 1*, S5-10.
- 228 Kalman, M., Inchley, J., Sigmundova, D., Iannotti, R. J., Tynjälä, J. A., Hamrik, Z., ...
229 Bucksch, J. (2015). Secular trends in moderate-to-vigorous physical activity in 32
230 countries from 2002 to 2010: A cross-national perspective. *European Journal of Public*
231 *Health*, *25*, 37–40. doi:10.1093/eurpub/ckv024
- 232 Liu, Y., Wang, M., Tynjälä, J., Lv, Y., Villberg, J., Zhang, Z., & Kannas, L. (2010). Test-
233 retest reliability of selected items of Health Behaviour in School-aged Children (HBSC)
234 survey questionnaire in Beijing, China. *BMC Medical Research Methodology*, *10*, 73.
235 doi:10.1186/1471-2288-10-73
- 236 Murphy, M. H., Rowe, D., Belton, S., & Woods, C. (2015). Validity of a two-item physical
237 activity questionnaire for assessing attainment of physical activity guidelines in youth.
238 *BMC Public Health*, *15*, 1–8. doi:/10.1186/s12889-015-2418-6
- 239 Ng, K., Rintala, P., Tynjälä, J., Villberg, J., & Kannas, L. (2014). Physical activity patterns of
240 adolescents with long term illnesses or disabilities in Finnish general education.
241 *European Journal of Adapted Physical Activity*2014, *7*, 58–72.
- 242 Poitras, V. J., Gray, C. E., Borghese, M. M., Carson, V., Chaput, J., Janssen, I., ... Tremblay,
243 M. S. (2016). Systematic review of the relationships between objectively measured
244 physical activity and health indicators in school-aged children and youth. *Applied*

- 245 *Physiology, Nutrition, and Metabolism*, 41, S197–S239. doi:10.1139/apnm-2015-0663
- 246 Prochaska, J. J., Sallis, J. F., & Long, B. (2001). A physical activity screening measure for
247 use with adolescents in primary care. *Archives of Pediatrics & Adolescent Medicine*,
248 155, 554–559. doi:10.1001/archpedi.155.5.554
- 249 Rimmer, J. H., Schiller, W., Chen, M.-D., Schiller, W., & Chen, M. (2012). Effects of
250 Disability-Associated Low Energy Expenditure Deconditioning Syndrome Interactive
251 Exercise Technologies and Exercise Physiology for Persons with Disabilities, Chicago,
252 IL. *Exercise Sport Science Review*, 40, 22–29. doi:10.1097/JES.0b013e31823b8b82
- 253 Ross, S. M., Bogart, K. R., Logan, S. W., Case, L., Fine, J., & Thompson, H. (2016).
254 Physical Activity Participation of Disabled Children: A Systematic Review of
255 Conceptual and Methodological Approaches in Health Research. *Frontiers in Public*
256 *Health*, 4, 187. doi:10.3389/fpubh.2016.00187
- 257 Sallis, J. F., Bull, F., Guthold, R., Heath, G. W., Inoue, S., Kelly, P., ... Hallal, P. C. (2016).
258 Progress in physical activity over the Olympic quadrennium. *The Lancet*, 6736, 1–12.
259 doi:10.1016/S0140-6736(16)30581-5
- 260 Tardi, R., & Njelesani, J. (2015). Disability and the post-2015 development agenda.
261 *Disability and Rehabilitation*, 37, 1496–1500. doi:10.3109/09638288.2014.972589
- 262 Torsheim, T., Cavallo, F., Levin, K. A., Schnohr, C., Mazur, J., Niclasen, B., & Currie, C.
263 (2016). Psychometric Validation of the Revised Family Affluence Scale: a Latent
264 Variable Approach. *Child Indicators Research*, 9, 771–784. doi:10.1007/s12187-015-
265 9339-x
- 266 Tremblay, M. S., Barnes, J. D., González, S. A., Katzmarzyk, P. T., Onywera, V. O., Reilly,
267 J. J., ... Team, G. M. 2. . R. (2016). Global Matrix 2.0: Report Card Grades on the
268 Physical Activity of Children and Youth Comparing 38 Countries and the Global Matrix
269 2.0 Research Team. *Journal of Physical Activity and Health*, 13, 343–366.

270 doi:10.1123/jpah.2016-0594

271 UNICEF. (2012). *The right of children with disabilities to education: A rights-based*
272 *approach to inclusive education*. Geneva. Retrieved from

273 https://www.unicef.org/ceecis/IEPositionPaper_ENGLISH.pdf

274 WHO. (2001). *International Classification of Functioning, Disability and Health*. Geneva:

275 WHO

276 WHO. (2010). *Global recommendations on physical activity for health*. Geneva: WHO

277 doi:10.1080/110264804100343

Table 1. Prevalence of LTID (%) in different countries in Europe by boys and girls

	Boys				Girls				χ^2*	p-value
	N	No LTID %	LTID %	Affected LTID %	N	No LTID %	LTID %	Affected LTID %		
Armenia	946	92.0	4.1	3.9	1315	94.8	3.6	1.7	11.45	.003
Bulgaria	2396	90.7	9.3	nd	2225	88.3	11.7	nd	7.80	.006
Czech Republic	2367	73.6	26.4	nd	2619	71.7	28.3	nd	2.33	.127
England	2311	75.6	18.6	5.8	2193	76.3	17.4	6.3	1.64	.441
Finland	1454	70.4	25.3	4.3	1671	68.3	26.0	5.6	3.53	.171
France	2523	79.0	16.1	5.0	2553	79.4	16.3	4.3	1.21	.547
Hungary	1464	76.2	20.4	3.5	1632	74.7	22.3	3.0	2.14	.342
Ireland	1320	76.0	17.8	6.2	2172	80.9	14.0	5.0	12.20	.002
MKD	1417	92.3	5.3	2.4	1656	93.6	4.6	1.8	2.17	.337
Poland	2182	82.8	10.6	6.6	2228	81.0	11.6	7.5	2.62	.271
Romania	1248	93.3	5.5	1.2	1502	90.7	6.9	2.5	8.16	.017
Scotland	2376	80.0	15.0	5.1	2493	83.2	11.4	5.4	13.48	.001
Sweden	3113	75.8	16.8	7.4	3266	74.8	16.4	8.8	4.22	.121
Slovakia	2299	81.7	16.3	2.0	2476	79.6	17.6	2.7	4.42	.110
Wales	1912	80.6	13.2	6.2	1987	82.9	11.0	6.1	4.65	.098
Total	29341	80.5	15.4	4.1	31988	80.5	15.2	4.3	1.53	.465

nd – no data

LTID – Long term illnesses or disabilities

χ^2* Chi-square test of independence of adolescents without LTID between boys and girls

Table 2. Proportion of no LTID, LTID, and affected LTID (%), adolescents who meet PA recommendations for health with adjusted odds ratio and 95% Confidence Intervals

	Boy										Girl							
	No LTID					Affected LTID					No LTID					Affected LTID		
	%	%	OR	LCI	UCI	%	OR	LCI	UCI	%	%	OR	LCI	UCI	%	OR	LCI	UCI
Armenia ^a	26.7	17.9	0.55	0.23	1.36	24.3	0.59	0.24	1.45	17.3	8.5	0.45	0.16	1.27	27.3	2.03	0.76	5.39
Bulgaria ^b	32.9	37.8	1.28	0.94	1.74	nd				22.0	21.8	1.01	0.73	1.40	nd			
Czech ^b	25.4	22.9	0.92	0.74	1.14	nd				17.8	19.3	1.16	0.93	1.45	nd			
England	20.8	21.1	0.95	0.71	1.27	22.4	1.01	0.62	1.62	13.0	16.3	1.30	0.94	1.81	12.2	1.08	0.62	1.87
Finland ^c	23.4	26.1	1.15	0.87	1.52	27.4	1.21	0.67	2.20	17.4	17.7	1.03	0.77	1.39	13.8	0.87	0.47	1.61
France	15.9	16.0	0.98	0.72	1.33	22.4	1.49	0.93	2.37	7.1	10.1	1.44	0.98	2.11	12.7	1.83	0.97	3.44
Hungary	27.9	22.1	0.75	0.55	1.02	25.5	0.86	0.44	1.68	16.3	17.3	1.07	0.77	1.47	14.3	0.91	0.40	2.07
Ireland	32.7	32.8	1.03	0.75	1.42	31.7	0.95	0.57	1.57	15.4	17.7	1.13	0.80	1.59	15.6	1.15	0.65	2.03
MKD	32.6	37.3	1.27	0.76	2.11	26.5	0.65	0.29	1.45	21.2	18.4	0.92	0.50	1.68	13.3	0.65	0.22	1.92
Poland	29.4	31.0	1.09	0.80	1.47	28.0	1.00	0.67	1.47	18.6	17.4	0.99	0.70	1.41	16.9	0.97	0.63	1.50
Romania	26.6	17.4	0.60	0.31	1.20	26.7	0.48	0.10	2.27	13.2	21.4	1.82	1.06	3.12	29.7	3.64	1.69	7.86
Scotland	19.3	20.8	1.00	0.74	1.34	22.5	1.10	0.67	1.82	13.5	15.4	1.11	0.77	1.60	14.8	1.27	0.76	2.12
Sweden	15.9	14.9	0.84	0.63	1.12	13.0	0.82	0.54	1.26	10.5	11.4	1.08	0.79	1.46	10.1	0.95	0.62	1.44
Slovakia	30.4	30.2	0.92	0.70	1.21	40.4	2.09	1.06	4.15	19.2	20.1	1.01	0.76	1.34	25.0	1.91	1.01	3.59
Wales	19.0	18.8	0.94	0.67	1.33	15.1	0.78	0.46	1.33	11.5	14.6	1.32	0.87	1.99	9.1	0.76	0.39	1.49
Total	24.6	23.4	0.89	0.81	0.98	22.5	0.91	0.78	1.06	15.4	16.6	1.09	0.98	1.22	14.2	1.06	0.89	1.26

^a Armenia used FASII

^b Bulgaria and Czech did not include data on severity ; nd – no data

^c Finland did not include data from 11-y

bold text p<.05, LCI = Lower Confidence Interval, UCI = Upper Confidence Interval

LTID – long-term illnesses, disabilities or medical conditions

no LTID is reference group, adjusted for age and FAS