

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Poranen-Clark, Taina; von Bonsdorff, Mikaela; Rantakokko, Merja; Portegijs, Erja; Eronen, Johanna; Pynnönen, Katja; Eriksson, Johan G.; Viljanen, Anne; Rantanen, Taina

**Title:** The temporal association between executive function and life-space mobility in old age

**Year:** 2018

**Version:** Accepted version (Final draft)

**Copyright:** © 2017, Oxford University Press

**Rights:** In Copyright

**Rights url:** <http://rightsstatements.org/page/InC/1.0/?language=en>

**Please cite the original version:**

Poranen-Clark, T., von Bonsdorff, M., Rantakokko, M., Portegijs, E., Eronen, J., Pynnönen, K., Eriksson, J. G., Viljanen, A., & Rantanen, T. (2018). The temporal association between executive function and life-space mobility in old age. *Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 73(6), 835-839. <https://doi.org/10.1093/gerona/glx217>

# The Temporal Association between Executive Function and Life-Space

## Mobility in Old Age

Taina Poranen-Clark, MSc, M.A.(Ed)<sup>1</sup>, Mikaela B. von Bonsdorff, PhD<sup>1,2</sup>, Merja Rantakokko, PhD<sup>1</sup>, Erja Portegijs, PhD<sup>1</sup>, Johanna Eronen<sup>1</sup>, Katja Pynnönen<sup>1</sup>, Johan G. Eriksson, MD, DM Sc<sup>2-4</sup>, Anne Viljanen, PhD<sup>1</sup>, Taina Rantanen, PhD<sup>1</sup>

<sup>1</sup>Gerontology Research Center, Faculty of Sport and Health Sciences, University of

Jyväskylä, Finland

<sup>2</sup>Folkhälsan Research Center, Helsinki, Finland

<sup>3</sup>Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland

<sup>4</sup>Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

### Corresponding author and reprint requests:

Taina Poranen-Clark, MSc, M.A.(Ed)

Gerontology Research Center

Faculty of Sport and Health Sciences

University of Jyväskylä

PO Box 35 (Viveca)

FI-40014 University of Jyväskylä

Tel. +358 50 566 3719, E-mail [taina.poranen-clark@jyu.fi](mailto:taina.poranen-clark@jyu.fi)

© The Author 2017. Published by Oxford University Press on behalf of The Gerontological Society of America. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

## The Temporal Association between Executive Function and Life-Space Mobility in Old Age

### ABSTRACT

**Background:** Life-space mobility, an indicator of community mobility, describes person's movements in terms of the distance from home, the frequency of movement and the need of assistance for movement. Executive function (EF) is a higher order cognitive function that supervises motor control, and plays a key role in a person's ability to function independently. Cognitive impairment often co-occurs with restricted life-space mobility, however, the direction of the longitudinal associations between EF and life-space mobility is unclear. The aim of this study was to investigate the temporal associations between EF and life-space mobility among community-dwelling older people.

**Methods:** 108 community-dwelling persons aged 76 to 91 years participated in the two-year follow-up study. Executive function was measured with the Trail Making Test (TMT). The Life-Space Assessment (range 0-120, higher scores indicate more mobility) was used to assess life-space mobility. Cross-lagged model design was used to examine longitudinal relationship between EF and life-space mobility. The model was adjusted for age and gender.

**Results:** Average age of participants at baseline was 82.2 (SD 4.1) years and 59% were women. Better EF at baseline predicted higher life-space mobility at follow-up (path coefficient = 3.81, 95% Confidential Interval; 0.84, 6.78,  $p = 0.012$ ), while baseline life-space mobility did not predict EF at follow-up.

**Conclusion:** Executive function was a determinant of life-space mobility. Supporting EF may enhance maintaining independence and active participation in old age.

**Keywords:** Cognitive aging, motor control, successful aging

## INTRODUCTION

Life-space mobility reflects a person's physical ability as well as psychosocial capability relative to the environmental requirements<sup>1</sup>. Life-space mobility assessment includes estimates of the distance, the frequency and the need of assistance for mobility<sup>1</sup>. Moving outside more often and travelling further away from home on a regular basis may be viewed as higher environmental complexity of mobility among older people<sup>2,3</sup>. The further away from one's home an individual moves the more complex the environment and the greater cognitive investments might be required<sup>2</sup>. There is evidence that older people with better cognitive function have larger life-space mobility<sup>4</sup>. Executive function (EF) has an important role in performance of control processes, which are required when planning and carrying out complex, goal-oriented tasks<sup>5</sup>. Therefore, EF might be an important higher order cognitive domain for older people enabling them to maintain their life-space mobility<sup>6</sup>.

Coexisting mobility limitations and cognitive impairments are prevalent among community-dwelling older people<sup>7</sup> interfering with their independency and active participation in society. However, according to previous studies the direction of the association between cognition and mobility are inconsistent<sup>8</sup>. There are some evidence from longitudinal studies which used Mini Mental State Examination as a measure of cognitive functioning, that having greater life-space mobility may be a protective factor for cognitive decline<sup>3,9</sup>. Furthermore, constricted life-space mobility has been found to be associated with increased risk of Alzheimer diseases and mild cognitive impairment<sup>10</sup>. Impaired EF is prevalent even among community-dwelling older people without a formal diagnosis of cognitive impairment, and has been found to be associated with longitudinal declines in functional status<sup>11-13</sup>. Further, one previous study suggests that poor EF measured with the Trail Making Test (TMT) is a strong predictor of decline in lower extremity function<sup>14</sup>, which is known to

be associated with constricted life-space mobility<sup>15</sup>. The results of a recent study reported TMT performance being associated with gait adaptation indicating that EF is important for planning and adjusting stepping in situations where rapid gait adjustment may be needed<sup>16</sup>. Thus, importance of EF may increase as task complexity increases in unfamiliar environments farther away from home. The TMT performance requires amongst other things attention, cognitive flexibility and task switching<sup>5</sup> that are cognitive processes needed for everyday challenges for older people when moving around in environment<sup>16</sup>. Consequently, using EF as a measure of cognitive functioning when investigating temporal association between cognition and life-space mobility could add useful information to current literature. The aim of this study was to assess the temporal association between EF and life-space mobility. Establishing the direction of the association between EF and life-space mobility might help to target interventions more accurately to the factor in which declines seem to appear first.

## **METHODS**

### **Study population**

At baseline, altogether 169 community-dwelling people aged 76 to 91 years participated in the Hearing, Cognition and Wellbeing Study. Face-to-face interviews and sensory, physical and cognitive functioning measurements were conducted in the participants' homes. Of the initial 169 persons, 108 participated in the follow-up study two years later (32 were not interested to participate, 19 had deceased, one moved outside study area, and nine were not reached), and formed the analytic sample of this study.

The participants of the current study were part of the Life-Space Mobility in Old Age (LISPE) study, described in detail previously<sup>17</sup>. LISPE is a population-based study among community living older people including 848 participants. Using random number tables a subset of 230 individuals was selected for the Hearing, Cognition and Wellbeing sub-study in year 2014<sup>18</sup>. Those willing to participate and who did not experience severe problems in communicating, underwent the examinations and interviews during spring 2014 (n= 169). The Hearing, Cognition and Wellbeing substudy sample (n=169) did not differ from the initial LISPE study sample (n=848) in terms of sex, age, years of education, number of chronic conditions, lower extremity performance measured with the Short Physical Performance Battery (SPPB), or cognitive functioning measured with Mini Mental State Examination (MMSE) (all p-values >0.127).

The LISPE study and the Hearing, Cognition and Wellbeing sub-study both comply with the principles of good scientific conduct and good clinical practice in all aspects of the Declaration of Helsinki, and were approved by the Ethical Committee of the University of Jyväskylä. All participants gave a written informed consent.

### **Executive function**

EF was measured with the Trail Making Test (TMT). TMT is a paper-and-pencil task providing information on visual search, attention, processing speed, cognitive flexibility, and executive function<sup>19,20</sup>. The TMT consists of two parts. In the TMT-A task participants were asked to make a trail with a pencil, sequentially drawing lines to connect randomly arranged encircled numbers (from 1 to 25) spread over a sheet of paper. In the TMT-B task participants were asked to make a trail in numeric and alphabetical order (1-A-2-B-3-C, etc.) by drawing lines to connect randomly arranged encircles containing numbers (from 1 to 13) and letters (from A to L) spread over a sheet. The

participants were asked to perform the task as quickly as possible without lifting the pencil from the paper. If an error occurred the examiner pointed it out immediately. Then the participants were asked to revert to last correct number or letter and continue there to complete the task at the expense of additional time. Time to complete each task was measured in seconds<sup>21,22</sup>. A maximum accepted time to complete each part of the test was 240 seconds and maximum accepted amount of errors was 4<sup>6,21,23</sup>. Delta-TMT was calculated by subtracting time to perform TMT-A from time to perform TMT-B to control for the effects of motor function, visual scanning and processing speed. Delta-TMT has been used in previous studies to indicate executive functioning<sup>24,25</sup>.

EF was categorized into three approximately equal distribution-based groups as follows. At baseline and follow-up, poor EF included those who started to do the TMT test but who could not complete the test an acceptable way; they did not perform TMT-A or TMT-B tasks within the given time (max 240 seconds), or made more than 4 errors, or who did not complete the task so that Delta-TMT could not have been calculated<sup>6</sup>. Intermediate EF included those whose Delta-TMT was between 95 and 179 seconds at the baseline and between 97 and 180 seconds at the follow-up. Good EF included those whose Delta-TMT was 94 seconds or less at the baseline and 97 seconds or less at the follow-up. The cut-off between intermediate and good EF was defined according to the median value of Delta-TMT among those who completed the tests.

### **Life-space mobility**

Life-space mobility was measured with the 15-item University of Alabama at Birmingham Study of Aging Life-Space Assessment (LSA)<sup>1</sup>, which was translated into Finnish<sup>26</sup>. The reliability of the Finnish translation of life-space assessment has been found to be acceptable<sup>26</sup>. Life-space mobility assessment measures person's actual mobility during the four weeks preceding the assessment on

the following life-space levels (=level score): bedroom (score 0), other rooms (1), outside home (2), neighborhood (3), town (4), beyond town (5). Participants were also asked how many times they attained each life-space level (=frequency score); daily (score 4), 4-6 times/ week (3), 1-3 times/ week (2), less than once a week(1), and whether they needed help from another person or used assistive device (=assistance score): no assistance (2), use of device only (1.5), personal assistance (1). A composite score was calculated as follows; level score \* frequency score \* assistance score at respective level, and then summed for all levels. Thus, a composite score reflects distance, frequency, and independence of movement, and was used as an indicator of life-space mobility<sup>26</sup>. The score ranges between 0 and 120, and higher scores indicated higher life-space mobility<sup>17,26</sup>.

### **Covariates and descriptive variables**

Participants' age and gender derived from the national population register were used as covariates in the analysis. Number of self-reported chronic conditions, global cognitive functioning, and lower extremity functioning were used to describe the characteristics of the participants. Number of self-reported chronic conditions was calculated from a list of 22 physician-diagnosed diseases and an additional open-ended question about any other physician-diagnosed chronic diseases.<sup>15,17</sup> Global cognitive functioning was assessed with Mini Mental State Examination (MMSE, range 0-30)<sup>27</sup>. Lower extremity functioning was assessed with Short Physical Performance Battery (SPPB). The test battery comprises three tests assessing standing balance, walking speed over a distance of 2.44 meters and time taken to complete five chair rises. Each test is rated from 0 to 4 points according to established age- and gender-specific cut-off points<sup>28,29</sup>. A sum score (range 1-12) was calculated if at least two of the tests were completed. Higher scores indicate better lower extremity function<sup>6,15</sup>.



## Statistical analyses

The longitudinal cross-lagged associations between EF and life-space mobility were estimated using cross-lagged model design. Cross-lagged analysis enables to study whether EF at baseline is associated with life-space mobility at follow-up after controlling for life-space mobility at baseline, or vice versa. The unstandardized values were used in the analyses. Weighted least square estimator (WLSMV) was used to obtain parameter estimates. The model was adjusted for age and gender. The analysis were performed with MPLUS version 5.21 (Muthén and Muthén, 1998-2009) which uses the Full Information Maximum Likelihood procedure for handling missing values. The proportion of missing data in individual variables varied between 2.7 to 4.6 %. The interaction between gender and EF on life-space mobility was not statistically significant ( $p > 0.185$ ), thus all analyses were pooled by gender. For all tests two-tailed p-values are reported and the level of significance was set at  $p < 0.05$ . The descriptive data analyses were carried out with SPSS IBM version 24.0 (SPSS, Armonk, NY, IBM Corp).

## RESULTS

Characteristics of the study participants at baseline and follow-up are presented in Table 1. The average age of the participants at baseline was 82.2 years (SD 4.1) and 59.3 % of them were women. The proportion of those with poor EF increased during the two year follow-up from 24.3 % to 29.5 %. However, the global cognition measured with Mini Mental State Examination remained nearly stable during the two-year follow-up period. The average composite score of life-space mobility declined during the two-year follow-up from 63.1 (SD 19.7) to 61.5 (SD 21.2) (Table1).

Figure 1 shows the unstandardized coefficients of cross-lagged model with two-year follow-up for EF and life-space mobility. At baseline, better EF was associated with higher life-space mobility. Both EF and life-space mobility showed relatively strong stability between baseline and follow-up assessments. After controlling for age, gender, and taking into account previous measurements of EF and life-space mobility, the analysis revealed that better EF at baseline predicted higher life-space mobility at follow-up (path coefficient = 3.81, 95% confidential interval; 0.84, 6.78,  $p=0.012$ ) while baseline life-space mobility was not associated with EF at follow-up (Figure 1). The cross-lagged model fitted the data well Comparative Fit Index (CFI) was 1.000, Tucker Lewis Index (TLI) was 1.000, and Root Mean Square Error Approximation (RMSEA) less than 0.0005.

Compared to those who did not participate in the follow-up study ( $n=61$ ), those who participated ( $n=108$ ) were younger (82.2 years, SD 4.1 vs. 83.6 years, SD 4.3,  $p=0.038$ ), and they had higher SPPB (9.3, SD 2.0, vs. 7.2, SD 3.4,  $p<0.001$ ), MMSE (26.3, SD 2.7 vs. 25.4, SD 2.5  $p=0.033$ ), and life-space mobility scores (63.1, SD 19.7 vs. 51.2, SD 20.4,  $p<0.001$ ) at baseline. The proportion of those who dropped out was highest among those with poor EF (53.7%), followed by those with intermediate (30.0%) or good (18.9%) EF. Those with poor EF at baseline were five times (OR= 4.99, 95% CI 2.09, 11.92), and those with intermediate EF almost two times (OR =1.84, 95% CI 0.74, 4.61) more likely to drop out from the follow-up compared to those with good EF.

Finally, we conducted additional cross-lagged modelling; first, to examine a stricter cut-point for good TMT, second, to analyze MMSE as an indicator of global cognition, and third, using Delta-TMT as a continuous variable. Using more strict cut-points suggested by an earlier study (good Delta-TMT performance  $<60s$ )<sup>30</sup> to define good EF (at baseline cut-point for good EF was 60s or less and at

follow-up 72s or less) did not change the results. The analysis for MMSE and life-space mobility revealed similar associations as with using EF; coefficients and 95% confidential intervals for baseline MMSE predicting follow-up life-space was 1.22 (0.29, 2.15) and baseline life-space mobility predicting follow-up MMSE 0.01(-0.01, 0.03). The associations for EF as continuous variable and life-space mobility among those who completed both parts of the TMT test (n= 78) were no statistically significant in either of the directions.

## DISCUSSION

To our knowledge, we were the first to examine the temporal association between executive function and life-space mobility among community-dwelling older people. Better EF at baseline predicted higher life-space mobility at follow-up, while baseline life-space mobility did not predict EF at follow-up. Thus, our findings suggest that impaired executive function is a precedent for restricted life-space mobility rather than restriction of life-space would further influence cognitive decline.

It is obvious in the light of previous studies that cognition is interrelated with mobility<sup>7</sup>. In addition to emerging literature on the associations between cognitive function and mobility, we were using specific higher order cognitive domain, EF, and a multidimensional mobility measure, life-space mobility, as measurements. Life-space mobility reflects person's actual mobility through a given area over a specific time period, and incorporates not only physical aspects of mobility but also psychosocial capability for participation in society<sup>1</sup>. Further, we examined which one of these measurements, EF or life-space mobility, is a stronger predictor for one another whereas previous

studies have predicted associations only for one of the possible directions<sup>9,10</sup>. This was done by using analysis that enable simultaneous examination of longitudinal influences of one factor on the other and vice versa, while controlling for contemporary associations between factors, and the stability of each factor over time. The association between EF and mobility has been reported being bidirectional<sup>8,31</sup>. However, our results are in line with some of the previous studies suggesting that EF is a predictor for mobility decline<sup>12-14</sup> and conflicting the previous results that suggest that life-space mobility predicts changes in cognitive functioning<sup>3,9,10</sup>. The latter might be explained by the fact that neither of the studies had EF as a measure of cognitive functioning. After all, a person might have some deficit of EF despite evident cognitive impairment<sup>12</sup>.

According to the environmental complexity theory, a more demanding environment requires more cognitive investment leading to increase in cognitive activity consequently resulting in better cognitive functioning.<sup>2</sup> However, our results suggest that higher order cognitive functioning such as executive functioning is a prerequisite for the capability to move further away from home and enabling to widen one's life-space. Life-space mobility is a mobility measure that takes into account all movement in its' different forms through one's environment. The life-space mobility reaching only to one's close neighborhood may require less cognitive investment than if an individual would be orienteering oneself further away from a familiar environment, and perhaps using a vehicle for transportation. Reduction in life- space mobility may reflect early changes in cognitive functioning. Impairment in EF may affect an individual's willingness to travel further away from home even before any other markers of cognitive decline have appeared. Restricted life-space mobility may be seen as an early sign of cognitive decline resulting from EF impairment.

In recent years, the understanding of the role of the brain in age-related declines in mobility has increased<sup>32,33</sup>. Furthermore, EF has been identified as one of the potential mediators between age-related cognitive decline and functional ability<sup>34</sup>. Cognitive processes such as attention, action-inhibition and task-switching are dependent on EF and needed for motor control<sup>35</sup>. These control processes are primarily supported by frontal areas of the brain that are most vulnerable in the aging process<sup>35</sup>, which might potentially explain the causal relationship between EF and life-space mobility. Decline in life-space mobility has been found to be associated with decline in quality of life<sup>36</sup> and development of ADL disability<sup>37</sup>. Our earlier study suggested that poor EF may predict steeper decline in life-space mobility but the results did not quite reach statistical significance, hence, the causal association between EF and life-space mobility stayed unclear<sup>6</sup>. However, the result of this study confirms our earlier suggestions of EF being a predictor of life-space mobility. Further, poorer lower extremity functioning and difficulties in use of transportation partly explained the association between poor EF and lower life-space mobility<sup>6</sup>. Thus, we did not adjust for SPPB as it being potential mediator between EF and life-space mobility<sup>6</sup>. Future studies should examine if promoting EF by means of cognitive training interventions could possibly delay mobility limitations that further diminish life-space mobility.

Our study had several strengths. By using life-space mobility and EF as measurements in this study, we were able to further existing literature relative to association between cognitive functioning and mobility. Further, we were able to include considerable old people (range 78-93 years at follow-up) from all social strata into this study, and the participant did not have severe cognitive decline at baseline (mean MMSE 26.3, SD 2.7, and 12% < 23). In addition, we used computer-assisted face-to-face interviews by trained interviewers at participants' homes, which likely improve the quality of data. Further, we conducted several further modelling to test our results. Using more strict cut-points for good EF showed that the results were not dependent on the intermediate-good cut-point.

The results of cross-lagged model using MMSE instead of EF as the cognitive measure revealed similar associations. We found no statistically significant associations in either of directions when using EF measure as a continuous variable in cross-lagged modelling. This suggests that the cross-lagged association between baseline EF and follow-up life-space mobility observed in the larger sample where all the participants were included, was driven by the poor values. This also supports our finding that the null life space mobility - EF association was not due to the scale properties of the categorized EF variable but rather due to the absence of a causal relationship. We are aware that the limited sample size in these sensitivity analyses decrease power even further, however, the coefficient is so low that it may not be considered material.

Our study had some limitations. First of all, due to the fact that the participants with poorer functioning were more likely to drop out from the follow-up study, the results can be generalized only to somewhat better functioning community-dwelling older adults, and it is likely that the strength of the associations over time may be underestimated. Second, we had only a two-year follow-up. Although, this is a significant time period among older people to experience noticeable changes in functional capability, a longer follow-up period might reveal stronger effects. Third, we were using a single test to measure executive function instead of multiple tests due to time constraints in the study protocol considering that the participants were very old. Although TMT is a widely used test and simple and easy to perform, further studies should use several test batteries for a more comprehensive evaluation of executive function.

## Conclusions

According to our knowledge, we were the first to explore the temporal association between EF and life-space mobility among community-dwelling older people. The results of the study suggest that better EF is a precedent for higher life-space mobility laying ground to future studies on the topic using larger sample sizes and longer follow-up periods with multiple follow-up points. Supporting EF may help maintain higher life-space mobility enhancing independency and quality of life in old age.

## ACKNOWLEDGEMENTS

### Funding

This study was supported by the Academy of Finland (the Future of Living and Housing Program ASU-LIVE; grant number 255403 to [TR], number 263729 to [AV]; number 285747 to [MR]; number 129369, 129907, 135072, 129255 and 126775 to [JGE] and number 257239 to [MBvB]; Finnish Ministry of Education and Culture to [TR], [MR] and [EP]. TP-C was supported by Yrjö Jahnesson Foundation and Juho Vainio Foundation.

The authors declare that there are no conflicts of interest.

**Legend for the Figure 1** Unstandardized coefficients (95% confidence intervals) of cross-lagged model with two-year follow-up for executive function (EF) and life-space mobility adjusted for age and gender (n=108). Path coefficients are statistically significant if zero is not included in the confidential intervals.

**REFERENCES**

1. Baker PS, Bodner EV, Allman RM. Measuring life- space mobility in community- dwelling older adults. *J Am Geriatr Soc.* 2003;51(11):1610-1614. doi: 10.1046/j.1532-5415.2003.51512.x.
2. Schooler C. Psychological effects of complex environments during the life span: A review and theory. *Intelligence.* 1984;8(4):259-281.
3. Crowe M, Andel R, Wadley VG, Okonkwo OC, Sawyer P, Allman RM. Life-space and cognitive decline in a community-based sample of african american and caucasian older adults. *J Gerontol A Biol Sci Med Sci.* 2008;63(11):1241-1245. doi: 10.1093/gerona/63.11.1241.
4. Sartori AC, Wadley VG, Clay OJ, Parisi JM, Rebok GW, Crowe M. The relationship between cognitive function and life space: The potential role of personal control beliefs. *Psychol Aging.* 2012;27(2):364.
5. Banich MT. Executive function the search for an integrated account. *Current Directions in Psychological Science.* 2009;18(2):89-94.
6. Poranen-Clark T, von Bonsdorff MB, Rantakokko M, et al. Executive function and life-space mobility in old age. *Aging Clinical and Experimental Research.* 2017:1-7.
7. Demnitz N, Esser P, Dawes H, et al. A systematic review and meta-analysis of cross-sectional studies examining the relationship between mobility and cognition in healthy older adults. *Gait Posture.* 2016;50:164-174.
8. Stijntjes M, Aartsen MJ, Taekema DG, et al. Temporal relationship between cognitive and physical performance in middle-aged to oldest old people. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences.* 2016;72(5):662-668.



9. Silberschmidt S, Kumar A, Raji MM, Markides K, Ottenbacher KJ, Al Snih S. Life- Space mobility and cognitive decline among mexican americans aged 75 years and older. *J Am Geriatr Soc*. 2017.
10. James BD, Boyle PA, Buchman AS, Barnes LL, Bennett DA. Life space and risk of alzheimer disease, mild cognitive impairment, and cognitive decline in old age. *The American Journal of Geriatric Psychiatry*. 2011;19(11):961-969.
11. Royall DR, Palmer R, Chiodo LK, Polk MJ. Declining executive control in normal aging predicts change in functional status: The freedom house study. *J Am Geriatr Soc*. 2004;52(3):346-352.
12. Johnson JK, Lui LY, Yaffe K. Executive function, more than global cognition, predicts functional decline and mortality in elderly women. *J Gerontol A Biol Sci Med Sci*. 2007;62(10):1134-1141. doi: 62/10/1134 [pii].
13. Gross AL, Xue QL, Bandeen-Roche K, et al. Declines and impairment in executive function predict onset of physical frailty. *J Gerontol A Biol Sci Med Sci*. 2016;71(12):1624-1630. doi: glw067 [pii].
14. Vazzana R, Bandinelli S, Lauretani F, et al. Trail making test predicts physical impairment and mortality in older persons. *J Am Geriatr Soc*. 2010;58(4):719-723.
15. Portegijs E, Rantakokko M, Mikkola TM, Viljanen A, Rantanen T. Association between physical performance and sense of autonomy in outdoor activities and Life- Space mobility in Community-Dwelling older people. *J Am Geriatr Soc*. 2014;62(4):615-621.
16. Caetano MJD, Menant JC, Schoene D, Pelicioni PH, Sturnieks DL, Lord SR. Sensorimotor and cognitive predictors of impaired gait adaptability in older people. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*. 2016:glw171.

17. Rantanen T, Portegijs E, Viljanen A, et al. Individual and environmental factors underlying life space of older people - study protocol and design of a cohort study on life-space mobility in old age (LISPE). *BMC Public Health*. 2012;12:1018-2458-12-1018. doi: 10.1186/1471-2458-12-1018 [doi].
18. Polku H, Mikkola TM, Rantakokko M, et al. Hearing and quality of life among community-dwelling older adults. *J Gerontol B Psychol Sci Soc Sci*. 2016. doi: gbw045 [pii].
19. Reitan,R., Wolfson,D., ed. *The halstead-reitan neuropsychologic test battery: Theory and clinical interpretation*. Tucson, AZ: Neuropsychology Press; 1993.
20. Strauss E, Sherman E, Spreen O. Trail making test. *A Compendium of Neuropsychological Tests.Administration, Norms, and Commentary*. 2006:655-677.
21. Lezak M, ed. *Neuropsychological assessment*. 3rd edition ed. Oxford: Oxford University Press; 1995.
22. Bowie CR, Harvey PD. Administration and interpretation of the trail making test. *Nature protocols*. 2006;1(5):2277-2281.
23. Lezak MD. *Neuropsychological assessment*. Oxford University Press, USA; 2004.
24. Ble A, Volpato S, Zuliani G, et al. Executive function correlates with walking speed in older persons: The InCHIANTI study. *J Am Geriatr Soc*. 2005;53(3):410-415.
25. Drane DL, Yuspeh RL, Huthwaite JS, Klingler LK. Demographic characteristics and normative observations for derived-trail making test indices. *Neuropsychiatry Neuropsychol Behav Neurol*. 2002;15(1):39-43.
26. Portegijs E, Iwarsson S, Rantakokko M, Viljanen A, Rantanen T. Life-space mobility assessment in older people in finland; measurement properties in winter and spring. *BMC Res Notes*. 2014;7:323-0500-7-323. doi: 10.1186/1756-0500-7-323 [doi].

27. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-198. doi: [http://dx.doi.org/10.1016/0022-3956\(75\)90026-6](http://dx.doi.org/10.1016/0022-3956(75)90026-6).
28. Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: Association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol.* 1994;49(2):M85-94.
29. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med.* 1995;332(9):556-562.
30. Papandonatos GD, Ott BR, Davis JD, Barco PP, Carr DB. Clinical utility of the Trail- Making test as a predictor of driving performance in older adults. *J Am Geriatr Soc.* 2015;63(11):2358-2364.
31. Tian Q, An Y, Resnick SM, Studenski S. The relative temporal sequence of decline in mobility and cognition among initially unimpaired older adults: Results from the baltimore longitudinal study of aging. *Age Ageing.* 2016. doi: afw185 [pii].
32. Holtzer R, Epstein N, Mahoney JR, Izzetoglu M, Blumen HM. Neuroimaging of mobility in aging: A targeted review. *J Gerontol A Biol Sci Med Sci.* 2014;69(11):1375-1388. doi: 10.1093/gerona/glu052 [doi].
33. Rosso AL, Studenski SA, Chen WG, et al. Aging, the central nervous system, and mobility. *J Gerontol A Biol Sci Med Sci.* 2013;68(11):1379-1386. doi: 10.1093/gerona/glt089 [doi].
34. Puente AN, Lindbergh CA, Miller LS. The relationship between cognitive reserve and functional ability is mediated by executive functioning in older adults. *Clin Neuropsychol.* 2015;29(1):67-81.
35. Ren J, Wu YD, Chan JS, Yan JH. Cognitive aging affects motor performance and learning. *Geriatr Gerontol Int.* 2013;13(1):19-27. doi: 10.1111/j.1447-0594.2012.00914.x [doi].

36. Rantakokko M, Portegijs E, Viljanen A, Iwarsson S, Kauppinen M, Rantanen T. Changes in life-space mobility and quality of life among community-dwelling older people: A 2-year follow-up study. *Quality of Life Research*. 2015:1-9.
37. Portegijs E, Rantakokko M, Viljanen A, Sipilä S, Rantanen T. Identification of older people at risk of ADL disability using the life-space assessment: A longitudinal cohort study. *Journal of the American Medical Directors Association*. 2016;17(5):410-414.

Accepted Manuscript


Table 1 Means, standard deviations (SD), and proportions of characteristics of participants at baseline and at two-year follow-up (n=108).

	Baseline	Follow-up
	%	%
<b>Women</b>	59.3	59.2
<b>Executive function (n=103)</b>		
Good	41.7	37.1
Intermediate	34.0	33.0
Poor	24.3	29.5
	<b>Mean (SD)</b>	<b>Mean (SD)</b>
<b>Age</b>	82.2 (4.1)	84.3 (4.1)
<b>Life-space mobility</b>	63.1 (19.7)	61.5 (21.2)
<b>Number of chronic conditions</b>	3.2 (1.6)	3.4 (1.7)
<b>Cognitive functioning (MMSE)</b>	26.3 (2.7)	26.2 (3.0)
<b>Lower extremity function (SPPB)</b>	9.21 (2.1)	9.10 (2.2)*

MMSE = Mini Mental State Examination, SPPB = Short Physical Performance Battery

\*Note, n =106

Figure 1



Accepted Manuscript