

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Ojanen, Tommi; Kyröläinen, Heikki; Igendia, Mikael; Häkkinen, Keijo

Title: Effect of Prolonged Military Field Training on Neuromuscular and Hormonal Responses and Shooting Performance in Warfighters

Year: 2018

Version: Accepted version (Final draft)

Copyright: © Association of Military Surgeons of the United States 2018.

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Ojanen, T., Kyröläinen, H., Igendia, M., & Häkkinen, K. (2018). Effect of Prolonged Military Field Training on Neuromuscular and Hormonal Responses and Shooting Performance in Warfighters. *Military Medicine*, 183(11-12), Article e705. <https://doi.org/10.1093/milmed/usy122>

Pages: 23

Words: 3779

Tables: 2

Figures: 1

References: 40

Contact: Tommi Ojanen

E-mail: tommi.ojanen@mil.fi

Guarantor: Tommi Ojanen

1
2
3
4
5
6
7
8
9
10
11
12 **Effect of prolonged military field training on neuromuscular and hormonal responses and**
13 **shooting performance in warfighters**
14
15
16
17
18

19 Tommi Ojanen, MSc¹

20
21 Heikki Kyröläinen, PhD^{2,3}

22
23 Mikael Igendia, MSc²

24
25
26 Keijo Häkkinen, PhD²
27
28
29
30

31 ¹Finnish Defence Research Agency, Finnish Defence Forces, P.O. Box 5, 04401 Järvenpää,
32
33 Finland;
34
35

36 ²Biology of Physical Activity, Faculty of Sport and Health Sciences, University of Jyväskylä. P.O.
37
38 Box 35, 40014 Jyväskylä, Finland;
39
40

41 ³National Defence University, P.O. Box 7, 00861 Helsinki, Finland
42
43
44

45 **KEYWORDS:** Physiology, Endocrinology, Training, Strength, Performance
46
47
48
49
50
51

52 Conflicts of interest: none
53
54

55 Funding: The study was funded by the Finnish Defence Forces.
56
57
58

59 Acknowledgements: The authors thank Mrs. Elina Vaara, MSc, for statistical support.
60
61
62
63
64
65

STRUCTURED SUMMARY

Introduction

Previous studies have shown that Military Field Training (MFT) has effects on warfighters' hormonal responses, neuromuscular performance and shooting accuracy. The aim of the present study was to investigate the changes in body composition, upper and lower body strength, serum hormone concentrations of testosterone (TES) and cortisol (COR), insulin-like growth factor – 1 (IGF-1) and sex hormone binding globulin (SHBG) and shooting accuracy during prolonged MFT.

Methods

Serum hormone concentrations, isometric strength of the upper and lower extremities and shooting performance were measured four times during the study: before MFT (PRE), after 12 days (MID), at the end of MFT (POST) and after four days recovery (RECO). The study was approved by the Finnish Defence Forces and was granted ethical approval by the Ethical Committee of the University of Jyväskylä.

Results

There was no change in prone shooting score between the measuring points. In the standing position, however, there was a significant ($p \leq 0.001$) decrease from PRE 58.2 ± 12.3 points to MID 45.2 ± 10.4 points. Also POST 61.4 ± 10.8 points and RECO 56.8 ± 13.6 points were significantly ($p \leq 0.001$) higher than MID 45.2 ± 10.4 points. Serum hormone concentrations of TES and IGF-1 decreased significantly during MFT. In COR and SHBG concentrations significant increases were observed during MFT. Individual changes in lower body strength and changes in shooting standing score between the measurement points (PRE - MID / POST / RECO) correlated significantly ($r=0.332$, $p=0.031$; $r=0.335$, $p=0.025$; $r=0.489$, $p=0.001$, respectively). The similar finding was observed with changes in upper body strength and changes in standing shooting between the PRE

1 and RECO measurement points (0.339, $p=0.010$). The changes in COR and the changes in prone
2 shooting showed a positive correlation in all measurement points ($r=0.531$, $p\leq 0.001$; $r=0.337$,
3 $p=0.024$; $r=0.572$, $p\leq 0.001$). The changes in IGF-1 correlated negatively ($r=-0.325$, $p=0.038$) with
4 shooting prone between the PRE and MID measurement points. The changes in shooting standing
5 and the changes in TES between PRE and POST correlated negatively ($r=-0.378$, $p=0.010$).
6
7
8
9

10 **Conclusion**

11
12
13
14
15
16
17
18 In this study we observed a decrease in leg strength from the PRE to MID measurements. When the
19 physical load requirements during the MFT decreased after the MID measurements, leg strength
20 increased. In addition, the shooting score from the standing position decreased from the PRE to MID
21 measurements and improved significantly from the MID to POST measurements. The prone shooting
22 score did not show any significant changes during the study period. Significant positive correlations
23 were found between the changes in standing shooting score and the changes in strength for the legs
24 and upper body. There was a positive correlation between the changes in serum COR concentrations
25 and changes in standing shooting score.
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Altogether, the present study showed that the prolonged MFT has adverse effect on the strength levels
41 and the shooting ability in warfighters. This shows that ensuring warfighters get an appropriate
42 amount of rest while performing their duties is important. Shooting from a prone position was not
43 affected by the changing workloads and this result indicated that soldiers should shoot from a prone
44 position, whenever possible, especially when fatigued.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

1
2
3
4 Warfighters are exposed to physical, environmental and mental stressors when training for military
5 operations. These stressors include factors such as physical and cognitive fatigue caused as a
6 consequence of prolonged physical exertion and/or sleep deprivation and insufficient energy and fluid
7 intake. During military field training (MFT) these factors have shown to cause disruptions in
8 hormonal balance^{1,2,3,4}, leading to reduced physical and cognitive performance^{5,6,7,8,9}, extended
9 recovery times¹⁰ and increased susceptibility to infections¹¹. How this kind of environment effects
10 on warfighters' hormonal responses, neuromuscular performance and shooting accuracy and the
11 consequences of these stressors and their influence on health, physical performance level and work
12 capacity of a warfighter are of vital importance to commanders.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 Previous studies have examined various biomarkers in warfighters in relation to nutrition status, fluid
29 intake, body composition^{5,12}, prolonged physical exertion, neuromuscular performance^{6,13} and sleep
30 deprivation^{14,15}. Decrements in lower and upper body strength have been demonstrated after a
31 prolonged MFT. The type of loading is a key determinant to which part of the body becomes fatigued
32 ^{2,6,7,13}. In order to carry out daily duties, warfighters must have good muscle strength capability¹⁶.
33
34 Even in the face of technological developments in the military field, the modern warfighters tasks
35 are, however, physically very demanding. Both lower and upper body strength must be considered:
36 marching long distances require a warfighter to have a strong lower body, while for lifting and
37 carrying heavy equipment, a warfighter must have good upper body strength^{7,17}. The modern
38 warfighter needs power and strength as well as aerobic capacity and muscular endurance. The optimal
39 level of either one is dependent on occupational tasks of a warfighter¹⁸. Multiple studies have also
40 demonstrated that strength training and higher strength levels coincide with lower injury levels,
41 particularly in the lower extremities^{19,20}.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 The impact of MFT on hormonal changes is well demonstrated. Cortisol (COR) and sex hormone
3
4 binding globulin (SHBG) concentrations have been shown to increase during prolonged
5
6 MFT^{1,3,7,21,22,23}, whereas testosterone (TES) and insulin-like growth factor - 1 (IGF-1) concentrations
7
8 have been shown to decrease^{1,6,21,24,25,26,27,28} Friedl et al. (2000) found that during US Army Ranger
9
10 training a warfighter lost 15 % in body mass, 7 % in fat free mass and 65 % of their fat mass. Nindl
11
12 et al. (2006) studied the effects of short term MFT on hormonal responses. The study showed that
13
14 there was a 3 % loss in body mass and 24 to 30 % loss in free TES levels. Kyröläinen et al. (2008)
15
16 found that during the first couple of days of strenuous MFT there was a significant increase in COR
17
18 levels (32 %) and a decrease in TES levels (27 %). Also, SHBG levels have been found to increase
19
20 during basic military training²³ and during heavy and strenuous MFT²⁶.

21
22
23
24
25
26
27
28
29 Shooting is one of the most important occupational skills for a warfighter. There is limited
30
31 information regarding shooting performance and how it is affected by prolonged MFT. It has been
32
33 shown that shooting performance differs as a result of different loading intensities^{29,30,31,32,33}.
34
35 Warfighters are required to move on the battlefield, and maximizing shot accuracy in a state of
36
37 increased heart rate and fatigue³¹. Several factors affect the accuracy of shooting in warfighters.
38
39 Anxiety has been shown to decrease the accuracy of shooting as well as a negatively impact on
40
41 decision making³⁴. Fatigue is also a significant factor affecting marksmanship, especially after
42
43 anaerobic physical strain^{30,31,32}. In addition to anxiety and fatigue, sleep deprivation has shown a
44
45 detrimental effect on the shooting accuracy. Research on how prolonged MFT in a multistressor
46
47 environment affects the warfighters shooting performance is lacking³³.

48
49
50
51
52
53
54
55
56 The aim of the present study was to investigate the changes in body composition, upper and lower
57
58 body strength, serum hormone (TES, COR, SHBG, IGF-1) concentrations and shooting accuracy

during a prolonged MFT. We hypothesized that we would observe declines in all measured variables and the study would provide better understanding of the associations between warfighters physical performance and shooting capability during MFT.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

METHODS

Subjects

Sixty-one (n=61) male subjects volunteered for the present study. Each subject (conscript) was a Finnish Army member conducting infantry training, performed during their six-month mandatory service in the Finnish Defence Forces. Based on exit interviews, twelve individuals dropped out due to discomfort of having to give blood samples and unsustained motivation to participate in the study. Forty-nine subjects between 19 to 22 years of age completed the study. The mean (\pm standard deviation) age was 20 ± 1 years, height of 179 ± 6 cm, body mass 73.5 ± 8.7 kg and body fat 12.6 ± 5.0 %.

All subjects were fully informed of the experimental design and possible risks, and every subject signed an informed consent before the study commenced. The subjects were informed that they could cancel their participation in the present study at any stage if they so wished without any consequences. This study was conducted according to the provisions of the Declaration of Helsinki and was granted ethical approval by the Ethical Committee of the University of Jyväskylä. The study was also approved by the Finnish Defence Forces.

Experimental Design

A week before MFT, all the subjects were tested for their baseline values. The same tests were also performed on the day 12 of MFT, at the end of MFT and after a recovery period of four days. Blood samples, isometric strength of the upper and lower extremities and shooting performance were recorded during the measurement days.

1 The PRE measurement week was a normal training week for the study participants. Duties included
2 lectures in the classroom, rifle maintenance and preparation for MFT. The entire 21-day MFT period,
3
4 which was divided into three phases, was performed in field conditions. During the first phase (ST)
5
6 the subjects performed combat drills and live-fire shooting exercises. The goal for each conscript was
7
8 to improve their combat and shooting skills and advance their weapon handling abilities. In this phase,
9
10 the aim was not to physically exhaust the conscripts but rather to ensure that each of them maintained
11
12 a high level of performance by ensuring that they had an appropriate amount of rest and sleep. The
13
14 normal training day started at 07:00 hours and ended not later than 19:00 hours.
15
16
17
18
19
20

21 In the second phase (MFT), they practiced moving from their base to their attacking positions. The
22
23 tasks performed included reconnaissance, combat maneuvers, patrolling and tactical road marches.
24
25

26 In the last phase (MFT) they executed combat mission as a part of larger military exercise. The tasks
27
28 were the same as in the second phase. After the prolonged MFT, the subjects had four days of
29
30 recovery, two at home and two at the garrison before the final study measurements were taken.
31
32
33
34
35

36 **Measurements**

37
38
39
40

41 *Neuromuscular performance*

42

43 Maximal isometric strength of the upper and lower extremities was measured with dynamometers.
44
45 The measurements were conducted using a leg and bench press dynamometer manufactured by the
46
47 University of Jyväskylä, Department of Biology of Physical Activity, Finland. The knee angle was
48
49 set to 107 degrees³⁵. During the maximal strength test for upper extremities the equipment was
50
51 adjusted for each subject so that when in sitting position with their feet flat on the floor their arms
52
53 were parallel to the floor and the elbow angle was 90 degrees. The test was performed by pushing the
54
55 bar horizontally. The subjects were given one trial attempt before the two actual test trials were
56
57
58
59
60
61
62
63
64
65

1 conducted on both leg extension and bench press. The subjects were instructed to produce maximal
2 force as fast as possible in both leg extension and bench press. On all trials, the testing personnel
3 encouraged them vocally during the maximal effort. The best performance was selected for analysis.
4
5
6

7 8 9 *Serum Hormone concentrations*

10 Venous blood samples were drawn four times from the antecubital vein after an overnight fast
11 between 0630 - 0730 to analyze TES, COR, SHBG and IGF-1 (Siemens Immulite 2000 XPI, Siemens
12 Healthcare, USA). Interassay coefficients of variance were 9.4% for TES, 7,6% for COR, 5.6% for
13 SHBG and 6.7% for IGF-1. The samples were centrifuged (Megafire 1.0 R Heraeus, DJB Lab Care,
14 Germany) at 3500 rpm for 10 minutes and frozen and transported to the laboratory for later analysis.
15
16
17
18
19
20
21
22
23
24
25

26 *Shooting*

27 The shooting test was performed from both prone and standing positions. First, the conscripts fired
28 ten shots at the target from a prone position and then ten shots from a standing position. The best
29 possible score was 100 points, and the result for each person was given at an accuracy of 0.1 points.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66 *Statistical Analysis*

67 The data was analyzed using IBM SPSS Statistics 22.0 (IBM Corporation, Armonk, NY). Statistical
68 analysis included descriptive statistics, Pearson correlation, and multivariate analysis of variance with
69 repeated measures. Probability adjusted *t* tests were used for pairwise comparisons. A general linear
70
71
72
73
74
75

1 model, with repeated measures ANOVA was used to analyze the differences between the different
2 measuring points. Bivariate correlation was used for correlation analysis where the changes in the
3 variables between the different time points were tested. The $p < 0.05$ criterion was used for
4 establishing the statistical significance.
5
6
7
8
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

RESULTS

Neuromuscular performance

There was a significant increase ($p \leq 0.05$) in isometric leg press force from MID of 3406 ± 923 N to POST 3532 ± 1011 N measurement. No other significant changes were found in leg press. Force of upper body isometric bench press increased significantly ($p \leq 0.001$) from PRE of 890 ± 181 N to MID 929 ± 179 N. There was a decline in POST of 900 ± 179 N ($p \leq 0.05$) and RECO 873 ± 178 N ($p \leq 0.001$) measurements when compared to MID 929 ± 179 N. RECO 873 ± 178 N ($p \leq 0.001$) was also significantly lower than POST of 900 ± 179 N measurement. (Table 1)

Shooting

There was no change in prone shooting score between the measuring points. In the standing position however there was significant ($p \leq 0.001$) decrease from PRE 58.2 ± 12.3 points to MID 45.2 ± 10.4 points. Also POST of 61.4 ± 10.8 points and RECO of 56.8 ± 13.6 points were significantly ($p \leq 0.001$) higher than MID of 45.2 ± 10.4 points. (Table 1)

Serum hormone concentrations

Testosterone

Serum TES concentrations decreased significantly ($p \leq 0.001$) from PRE 18.5 ± 4.5 nmol/l to MID 13.8 ± 4.9 nmol/l (-25.0%) and POST 16.0 ± 4.2 nmol/l (-13.0%). The POST value was also significantly ($p \leq 0.01$) higher (15.9%) than the MID. RECO 19.9 ± 3.7 nmol/l was significantly ($p \leq 0.001$) higher (44.2 %) than MID and POST (24.4 %) and also higher than PRE ($p \leq 0.05$) (8.2 %). (Table 2)

Cortisol

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

There was a significant increase in serum COR concentrations between PRE 301 ± 86 nmol/l and MID 355 ± 76 nmol/l ($p \leq 0.05$) (17.8 %), POST 396 ± 69 nmol/l ($p \leq 0.001$) (31.3 %) and RECO 385 ± 85 nmol/l ($p \leq 0.001$) (27.8 %). Also POST cortisol was significantly ($p \leq 0.001$) higher than that of MID (15.9%). (Table 2)

Insulin-like Growth Factor-1

IGF-1 showed a significant ($p \leq 0.001$) reduction from PRE of 40.6 ± 7.7 nmol/l to MID 32.5 ± 8.9 nmol/l (-20.0 %) and POST 32.5 ± 7.7 nmol/l (-20.0 %). Also a significant increase of the same magnitude ($p \leq 0.001$) was observed between RECO of 39.4 ± 7.8 nmol/l and MID (21.2 %) and POST (21.2 %). (Table 2)

Sex Hormone Binding Globulin

There was a significant increase in SHBG from PRE of 30.1 ± 7.6 nmol/l to MID 32.8 ± 7.9 nmol/l ($p \leq 0.01$) (9.0 %) and POST of 34.3 ± 9.1 nmol/l ($p \leq 0.001$) (14.0 %). A significant ($p \leq 0.001$) decline was observed between POST and RECO 31.5 ± 8.1 nmol/l (-8.2 %). (Table 2)

Associations between strength, shooting and serum concentrations

In the present study individual changes in lower body strength and changes in shooting standing score between the measurement points (PRE - MID / POST / RECO) correlated significantly ($r = 0.332$, $p = 0.031$; $r = 0.335$, $p = 0.025$; $r = 0.489$, $p = 0.001$). The same effect was found with changes in upper body strength and changes in shooting standing between PRE and RECO measurement points ($r = 0.339$, $p = 0.010$). With regard to serum concentrations a negative correlation was observed between the changes in SHBG and changes in IGF-1 in all measurement points ($r = -0.310$, $p = 0.043$; $r = -0.482$, $p = 0.001$; $r = -0.382$, $p = 0.010$). There was also a positive correlation between the changes in SHBG and the change in TES ($r = -0.330$, $p = 0.027$) when comparing the PRE and RECO measurement points.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The changes in COR correlated negatively with the changes in TES ($r=-0.341$, $p=0.025$) and the changes in IGF-1 ($r=-0.346$, $p=0.023$) between the PRE and MID measurement points. The changes in COR and the changes in shooting prone showed a positive correlation in all measurement points ($r=0.531$, $p\leq 0.001$; $r=0.337$, $p=0.024$; $r=0.572$, $p\leq 0.001$). The changes in IGF-1 correlated negatively ($r=-0.325$, $p=0.038$) with shooting prone between the PRE and MID measurement points. The changes in shooting standing and the changes in TES between PRE and POST correlated negatively ($r=-0.378$, $p=0.010$).

DISCUSSION

1
2
3
4
5 Our study supports previous investigations and adds to the existing literature on stressors and key
6 military performance measures. The present study showed the decrease in leg strength from the PRE
7 to MID measurements. When the physical load decreased after the MID measurements, the leg
8 strength increased. In addition, the shooting score from the standing position decreased from the PRE
9 to MID measurements and improved significantly from the MID to POST measurements. The
10 shooting prone score did not show any significant changes during the study period. Serum COR and
11 SHBG concentrations increased from PRE to MID and POST. SHBG increased in RECO
12 measurements, but COR did not change. Serum TES and IGF-1 concentrations decreased during MFT
13 and increased in RECO measurements. Significant positive correlations were found between the
14 changes in standing shooting score and the changes in strength for the legs and upper body. There
15 was a positive correlation between the changes in serum cortisol concentrations and changes in
16 standing shooting score. All the measured variables except cortisol returned close to the baseline level
17 after the recovery period.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Previous military field exercise studies^{2,6,7,13} have shown that prolonged physical activity and
40 negative energy balance combined with sleep deprivation have negative impact on neuromuscular
41 performance and hormonal balance of a warfighter. In the present study, maximal upper body strength
42 increased significantly from the PRE to MID measurements, but decreased back to PRE levels in the
43 POST and RECO measurements. This was probably partly due to learning, but indicated that upper
44 body strength was not affected by prolonged MFT.
45
46
47
48
49
50
51
52
53
54
55

56 It has been shown that lower body strength is an important factor for warfighters to be successful in
57 their duties^{16,17}. In the previous studies the lower body power has been reported both to either decrease
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

or increase² after MFT. In the present study, a slight, but not significant, decrease in leg extension values in the MID measurements. This was probably due to the higher loading of the first part of MFT as the four days before the MID measurements were physically the most demanding part of MFT. These findings suggest that lower body strength levels are related to the loads of MFT. There were no further declines after the loading was reduced. This indicates that it is possible to recover physically during the present type of MFT. Kyröläinen et al. (2008) had similar findings with reduced load during prolonged MFT. They showed that a lighter period between two intensive MFT parts can allow strength and hormone levels to recover before the commencing of the second intensive training period.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Serum TES concentration decreased significantly from the PRE to MID (-25.0%) and PRE to POST (-13.0%) measurement. Similar findings have been reported earlier^{1,3,21}. A significant increase was observed between MID to POST (15.9%) and POST to RECO (24.4%) measurements. This was probably due to the amount of physical load during MFT, as it was physically harder in the beginning of MFT.

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Serum COR concentration on the other hand, increased during MFT (31.3%) and did not return back to resting levels during the recovery period (27.8%). Friedl et al. (2000) and Nindl et al. (2007) found similar effect with US Army Rangers. Cortisol concentration increased significantly during the study period, but no recovery period was included. Kyröläinen et al. (2008) found that during a 20 day MFT cortisol levels increased by 32 %. COR, unlike TES and strength values did not begin to increase after the MID measurements. The present results showed that despite the loading was reduced in the latter part of MFT, the stress levels were at highest at the end of MFT, when the warfighters' training reached its capstone exercise.

57
58
59
60
61
62
63
64
65

Previous studies have shown that SHBG concentration has increased during strenuous physical training²². In a study with Finnish conscripts, Tanskanen et al. (2011) showed that SHBG values were

1 elevated during eight weeks of military training. Accordingly, Alemany et al. (2008) have shown an
2 increase in SHBG during MFT. In the present study, SHBG concentrations increased significantly
3 from the PRE values in the MID (9.0%) and POST (14.0%) measurement points, but returned back
4 to the PRE values in the RECO measurements.
5
6
7
8
9

10 IGF-I is one particular biomarker that has demonstrated a positive relationship between circulating
11 levels and aerobic fitness³⁶. Additional biomarkers (e.g. COR, TES, SHBG) have also shown to
12 predict excessive rates of physical performance change. In addition to the ability of biomarkers to
13 predict fitness measures, certain biomarkers that are sensitive to changes in homeostasis may serve
14 as early indicators of stress or overreaching^{3,23,26}. Rosendahl et al. (2002) reported a significant
15 decline in total IGF-1 values after 11 weeks of garrison training. There have also been decreases in
16 IGF-1 values after MFT³⁷ and longer special force training⁶.
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 In the present study IGF-1 concentration decreased from the PRE to MID measurements by 20.0%.
32 The values stayed the same from MID to POST and returned back to resting levels in the RECO
33 measurements. Similar findings have been reported in earlier studies^{7,28,38}. Even more drastic changes
34 (-62.0%) have been found during special force training⁶. The decrease in IGF-1 concentration is
35 mainly due to physical strain and energy deficit, which has been shown to decrease IGF-1
36 concentration²¹. This decrease in IGF-1 concentration indicates that warfighters' ability to recovery
37 from physical strain and to repair muscle damage is weakened and can lead to decrease in physical
38 performance. IGF-1 concentration seems to be a good marker to evaluate warfighters' recovery and
39 physical performance.
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 In previous studies soldiers' shooting performance has mainly been measured after different
55 simulations of military work. These studies have shown that drastic anaerobic work periods can
56 decrease shooting accuracy, but it has been shown to recover quickly³³. Evans et al. (2003) observed
57
58
59
60
61
62
63
64
65

1 the fatiguing upper body obstacle course decreased shooting accuracy significantly, but it recovered
2 quickly. Load carriage has also been shown to decrease shooting accuracy⁴⁰, although it seems that
3 there is no change before the carried loads of 45% of body weight³⁴. In a study of Tenan et al. (2017)
4 shooting performance was studied during a loaded march with live-fire shooting in the field. They
5 found that load carriage and marching did not affect shooting accuracy, actually the shooting
6 improved with lighter load. In the present study we found that shooting from the prone position did
7 not alter during MFT. Standing shooting was more sensitive to fatigue, but it seemed to decrease only
8 after the first and most physically demanding part of MFT. Declines in lower body strength and
9 hormonal concentrations have been shown to occur during MFT^{6,7}. This might explain the decline in
10 standing shooting score as physical exhaustion causes the heart rate to elevate and influence the
11 shooting mechanisms³³. More research should be done in the future to investigate the interaction
12 between mechanical and physiological effects and differences in the standing and prone shooting
13 positions in soldiers.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 The change in standing shooting score correlated positively with the change in lower body strength.
32 The similar correlation was observed between the change in upper body strength and the change in
33 standing shooting only between the PRE and RECO measurements. It seems that muscular strength,
34 especially in the lower body seems to be important for the standing shooting accuracy of a warfighter
35 during MFT. Both lower and upper body strength seems to be important for enhancing the recovery
36 process and thus the standing shooting accuracy in the recovery phase.
37
38
39
40
41
42
43
44
45
46

47 In the RECO measurements almost all the measured values were recovered to the PRE measurement
48 values. Serum COR concentration was significantly higher than in the PRE measurements. Also, body
49 weight (BW) and skeletal muscle mass (SMM) were significantly lower than in the PRE
50 measurements, but had almost recovered to the PRE values. It seems like four days is adequate for
51 the warfighter to recover from the physical strain of the present prolonged MFT in all measured
52 variables except for COR concentration.
53
54
55
56
57
58
59
60
61

1 The present study had some limitations. Participants were representatives of the larger population
2 doing their military service. A number of the subjects in the study represents only a small fraction of
3
4 the total number of service members. Thus, some caution should be used when extrapolating the
5
6 physiological data to the general population. Also, the number of dropped subjects in the study might
7
8 have had an influence on the results, because most of the dropped subjects were among the poorest
9
10 half with regard to physical fitness. On the other hand, this might also decrease the motivational effect
11
12 of the results, when the least motivated subjects were dropped out of the study. The present subjects
13
14 were familiarized with most of the physical tests, but some new tests were implemented (maximal
15
16 isometric tests), which could have had some influence in the results due to learning effect.
17
18 Nevertheless, all the measurements were carefully controlled by the trained researchers, and the tests
19
20 were completed in the same order and same time of day in all the measurement points.
21
22
23
24
25
26

27 The strength of the present study was that the conscripts performed similar duties during the study
28
29 period. In addition, there was a daily control of the subjects and the researchers were able to monitor
30
31 the loading during the whole study period. The MFT lasted for 21 days, which made it more realistic
32
33 when compared to short (three to five days) MFTs. The physical loading in the latter part of MFT
34
35 was lower than expected. This may have influenced the observed measures as they (shooting and
36
37 isometric maximal strength) did not decline after the MID measurements as expected based on prior
38
39 studies.
40
41
42
43
44

45 The present study showed that the prolonged MFT has adverse effect on the strength levels and the
46
47 shooting ability in warfighters. Therefore, to ensure that warfighters get an appropriate amount of
48
49 rest, while performing their duties is important. Shooting from a prone position was not affected by
50
51 the changing loads and this result indicated that soldiers should shoot from a prone position, whenever
52
53 possible, especially when fatigued.
54
55
56
57
58
59
60
61
62
63
64
65

References:

1. Alemany JA, Nindl BC, Kellogg MD, Tharion WJ, Young AJ and Montain SJ. Effects of dietary protein content on IGF-I, testosterone, and body composition during 8 days of severe energy deficit and arduous physical activity. *J Appl Physiol.* 2008 Jul;105(1):58-64.
2. Chester AL, Edwards AM, Crowe M, Quirk F. Physiological, biochemical, and psychological responses to environmental survival training in the royal Australian air force. *Mil Med.* 2013 Jul;178(7):829-35.
3. Kyröläinen H, Karinkanta J, Santtila M, Koski H, Mäntysaari M, Pullinen T. Hormonal responses during a prolonged military field exercise with variable exercise intensity. *Eur J Appl Physiol.* 2008 Mar;102(5):539-46.
4. Tyyskä J, Kokko J, Salonen M, Koivu M, Kyröläinen H. Association with physical fitness, serum hormones and sleep during a 15-day military field training. *J Sci Med Sport.* 2010 May;13(3):356-9.
5. Margolis LM, Rood J, Champagne C et al. Energy balance and body composition during US Army special forces training. *Applied Physiology, Nutrition, and Metabolism.* April 2013, Vol. 38 Issue 4, p396.
6. Nindl BC, Barnes BR, Alemany JA, Frykman PN, Shippee RL, Friedl KE. Physiological consequences of U.S. Army Ranger training. *Med Sci Sports Exerc.* 2007 Aug;39(8):1380-7.
7. Nindl BC, Leone CD, Tharion WJ et al. Physical performance responses during 72 h of military operational stress. *Med Sci Sports Exerc.* 2002 Nov;34(11):1814-22.
8. Richmond VL, Horner FE, Wilkinson DM, Rayson MP, Wright A, Izard R. Energy balance and physical demands during an 8-week arduous military training course. *Mil Med.* 2014 Apr;179(4):421-7.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
9. Tharion WJ, Lieberman HR, Montain SJ, Young AJ, Baker-Fulco, CJ, DeLany JP, Hoyt RW. Energy requirements of military personnel. *Appetite* 44:47-65, 2005.
10. Henning PC, Scofield DE, Spiering BA et al. Recovery of Endocrine and Inflammatory Mediators Following an Extended Energy Deficit. *J Clin Endocrinol Metab*, March 2014, 99(3);956-964.
11. Dimend BC, Fortes MB, Greeves JP et al. Effect of daily mixed nutritional supplementation on immune indices in soldiers undertaking an 8-week arduous training programme. *Eur J Appl Physiol* 2012 Apr;112(4):1411-18.
12. Friedl KE, Body composition and military performance - many things to many people. *J Strength Cond. Res.* 2012 Jul; 26(2): S87-100.
13. Sporiš G, Harasin D, Bok D, Matika D, Vuleta D. Effects of a training program for special operations battalion on soldiers' fitness characteristics. *J Strength Cond Res.* 2012 Oct;26(10):2872-82.
14. Williams SG, Collen J, Wickwire E, Lettieri CJ, Mysliwiec V. The Impact of sleep on soldier's performance. *Current Psychiatry Reports*, Aug 2014, Vol. 16 Issue 8.
15. Lentino CV, Purvis DL, Murphy KJ, Deuster PA. Sleep as a component of the performance triad: the importance of sleep in a military population. *US Army Med Dep J.* 2013 Oct-Dec 98-108.
16. Kraemer WJ, Szivak TK. Strength training for the warfighter. *J Strength Cond Res* 2012 Jul;26 (7):S107-118.
17. Welsh TT, Alemany JA, Montain SJ et al. Effects of intensified military field training on jumping performance. *Int J Sports Med.* 2008; 29: 45-52.
18. Friedl KE, Knapik JJ, Häkkinen K et al. Perspectives on Aerobic and Strength Influences on Military Physical Readiness: Report of an International Military Physiology Roundtable. *J Strength Cond Res.* 2015 Nov;29 Suppl 11:S10-23.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
19. Harman, E., Gutkunst, D., Frykman, D. et al. Effects of two different eight-week training programs on military physical performance. *J Strength Cond Res* 2008 22 (2), 524.
 20. Piirainen JM, Salmi JA, Avela J, Linnamo V. Effect of body composition on the neuromuscular function of Finnish conscripts during an 8-week basic training period. *J Strength Cond Res*. 2008 Nov;22(6): 1916-25.
 21. Friedl KE, Moore RJ, Hoyt RW, Marchitelli LJ, Martinez-Lopez LE, Askew WE. Endocrine markers of semistarvation in healthy lean men in a multistressor environment. *J Appl Physiol*. 2000 8:1820-1830.
 22. McCauley, G. O., McBride, J. M., Cormie, P. et al. Acute hormonal and neuromuscular responses to hypertrophy, strength and power type resistance exercise. *European journal of applied physiology* 2009 105(5), 695-704.
 23. Tanskanen MM, Kyröläinen H, Uusitalo AL et al. Serum sex-hormone binding globulin and cortisol concentrations are associated with overreaching during strenuous military training. *J Strength Cond Res*. 2011 Mar;25(3):787-97.
 24. Gomez-Merino D, Chennaoui M, Burnat P, Drogou C, Guezennec CY. Decrease in serum leptin after prolonged physical activity in men. *Med Sci Sports Exerc*. 2002 Oct;34(10):1594-9.
 25. Gomez-Merino D, Chennaoui M, Burnat P, Drogou C, Guezennec CY. Immune and hormonal changes following intense military training. *Mil Med*. 2003 Dec;168(12):1034-8.
 26. Nindl BC, Castellani JW, Young AJ et al. Differential responses of IGF-I molecular complexes to military operational field training. *J Appl Physiol*. 2003 Sep;95(3):1083-9.
 27. Nindl BC, Rarick KR, Castellani JW et al. Altered secretion of growth hormone and luteinizing hormone after 84 h of sustained physical exertion superimposed on caloric and sleep restriction. *J Appl Physiol*. 2006 Jan;100(1):120-8.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
28. Vaara JP, Kallioma R, Hynninen P, Kyröläinen H. Physical fitness and hormonal profile during an 11-week paratroop training period. *J Strength Cond Res.* 2015 Nov 29(S11):S168-72.
 29. Vickers JN, Williams AM. Performing under pressure: The effects of physiological arousal, cognitive anxiety and gaze control in biathlon. *Journal of Motor Behaviour* 2007 39, 381.
 30. Swain DP, Ringleb SI, Naik DN, Butowicz CM. Effect of Training with and without a Load on Military Fitness Tests and Marksmanship. *J Strength Cond Res.* 2011 Jul;25(7):1857-65.
 31. Frykman PN, Merullo DJ, Banderet LE, Gregorczyk K, Hasselquist L. Marksmanship Deficits Caused by an Exhaustive Whole-Body Lifting Task With and Without Torso-Borne Loads. *J Strength Cond Res.* 2012;26(7):S30-36.
 32. Jaworski RL, Jensen A, Niederberger B, Congalton R, Kelly KR. Changes in Combat Task Performance Under Increasing Loads in Active Duty Marines. *Mil Med.* 2015 Mar;180(3S):179- 186.
 33. Tenan MS, LaFiandra ME, Ortega SV. The Effect of Soldier Marching, Rucksack Load, and Heart Rate on Marksmanship. *Human Factors.* 2017 March;59(2):259-267.
 34. Nibbeling N, Oudejans R. Anxiety and Perceptual-Motor Performance: Towards an integrated model of concepts, mechanisms, and processes. *2014 Psychological Research* 76, 747.
 35. Häkkinen K, Kallinen M, Izquierdo M et al. Changes in agonist-antagonist EMG, muscle CSA, and force during strength training in middle-aged and older people. *J Appl Physiol.* 1998 Apr;84(4):1341-9.
 36. Nindl BC, Santtila M, Vaara J, Häkkinen K, Kyröläinen H. Circulating IGF-1 is associated with fitness and health outcomes in a population of 846 young healthy men. *Growth Horm IGF Res.* 2011 Jun;21(3):124-8.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
37. Nindl BC. Insulin-Like Growth Factor-I as a Candidate Metabolic Biomarker: Military Relevance and Future Directions for Measurement. *J Diabetes Sci Technol.* 2009 Mar 1;3(2):371-6.
38. Rosendal, L., Henning, L., Flyvbjerg, A., Frystyk, J., Ørskov, H., & Kjær, M. Physical capacity influences the response of insulin-like growth factor and its binding proteins to training. *J Appl Physiol* 2002 93, 1669–1675.
39. Evans RK, Scoville CR, Ito MA, Mello RP. Upper body fatiguing exercise and shooting performance. *Mil Med.* 2003 Jun;168(6):451-6.
40. Knapik JJ, Ang P, Meiselman H et al. Soldier performance and strenuous road marching: influence of load mass and load distribution. *Mil Med.* 1997 Jan;162(1):62-7.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65






Figure 1. Correlations between the lower body maximal isometric strength and shooting standing score between the PRE-MID, PRE-POST and PRE-RECO measurement points

Abbreviations: PRE, Before training measurements; POST, Post training measurements; RECO, Recovery measurements

Isometric strength	PRE	MID		POST		RECO	
Bench Press (N)	890±181	929±179	***	900±179	†	873±178	*†††‡‡‡‡
Leg Press (N)	3495±931	3406±923	-	3532±1011	†	3424±913	-

Shooting	PRE	MID		POST		RECO	
Prone (Points)	84.3±11.7	85.5±9.1	-	84.8±11.5	-	87.2±11.0	-
Standing (Points)	58.2±12.3	45.2±10.4	***	61.4±10.8	†††	56.8±13.6	†††‡

Table 1. Mean (\pm SD) values of isometric strength and shooting tests during MFT (*, †, ‡ = $p < 0.05$, ***, †††, ‡‡‡ = $p < 0.001$; *, ***, ††† = compared to PRE values, ††† = compared to MID values, ‡, ‡‡‡ = compared to POST values)

Abbreviations: PRE, Before training measurements; POST, Post training measurements; RECO, Recovery measurements

Hormones	PRE	MID		POST		RECO	
TES (nmol/l)	18.4±4.5	13.8±4.9	***	16.0±4.2	***††	19.9±3.7	*†††††††
COR (nmol/l)	301±86	355±76	*	396±69	***††	385±85	***
IGF-1 (nmol/l)	40.6±7.7	32.5±8.9	***	32.5±7.7	***	39.4±7.8	†††††††
SHBG (pmol/l)	30.1±7.6	32.8±7.9	**	34.3±9.1	***	31.5±8.1	†††

Table 2. Mean (\pm SD) serum hormone concentrations and SHBG concentrations during MFT (* = $p < 0.05$, **, †† = $p < 0.01$, ***, ††††, ††††† = $p < 0.001$; *, **, *** = compared to PRE values, †, ††, ††† = compared to MID values, †, ††, ††† = compared to POST values)

Abbreviations: PRE, Before training measurements; POST, Post training measurements; RECO, Recovery measurements; TES, Testosterone; COR, Cortisol; IGF-1, Insulin-like growth factor 1; SHBG, Sex hormone-binding globulin