

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Helenius, Ilkka; Paukkunen, Hannu

Title: Double D-meson production in proton-proton and proton-lead collisions at the LHC

Year: 2020

Version: Published version

Copyright: © 2019 The Authors. Published by Elsevier B.V.

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Helenius, I., & Paukkunen, H. (2020). Double D-meson production in proton-proton and proton-lead collisions at the LHC. *Physics Letters B*, 800, Article 135084.

<https://doi.org/10.1016/j.physletb.2019.135084>



Double D-meson production in proton-proton and proton-lead collisions at the LHC

Ilkka Helenius^{a,b}, Hannu Paukkunen^{a,b,*}

^a University of Jyväskylä, Department of Physics, P.O. Box 35, FI-40014 University of Jyväskylä, Finland

^b Helsinki Institute of Physics, P.O. Box 64, FI-00014 University of Helsinki, Finland



ARTICLE INFO

Article history:

Received 20 June 2019

Received in revised form 4 October 2019

Accepted 2 November 2019

Available online 6 November 2019

Editor: J.-P. Blaizot

Keywords:

Open heavy-flavour production

Double-parton scattering

Proton-nucleus scattering

Perturbative QCD

ABSTRACT

We consider the simultaneous production of two heavy-flavoured hadrons – particularly D mesons – at the LHC. We base our calculations on collinearly factorized QCD at next-to-leading order, using the contemporary parton distribution functions and D-meson fragmentation functions. The contributions of double-parton scatterings are included in the approximation of independent partonic interactions. Our framework benchmarks well with the available proton-proton data from the LHCb collaboration giving us confidence to make predictions for proton-lead collisions. Our results indicate that the double D-meson production in proton-lead collisions should be measurable at the LHCb kinematics with the already collected Run-II data, and should provide evidence for double-parton scattering at perturbative scales with a nuclear target.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The recent measurements of inclusive open heavy-flavour – particularly D and B mesons – in proton-proton (p-p) collisions at the CERN Large Hadron Collider (LHC) [1–8] provide opportunities to expose different facets of Quantum Chromodynamics (QCD) [9]. On one hand, due to the heavy-quark mass which serves as a hard interaction scale, the perturbative QCD calculations [10–15] can be extended e.g. to very small transverse momenta (p_T) where the calculations with massless quarks become inherently invalid. As the measurements at low p_T are statistically very precise, they offer an ideal testbed to benchmark perturbative calculations at low interaction scales. On the other hand, open heavy-flavour production can be used as a tool to probe non-perturbative aspects of heavy-quark fragmentation [16] and the quark-gluon structure of protons and nuclei [17–24]. The low- p_T open heavy-flavour production in proton-lead (p-Pb) collisions [25–28] may also open prospects to disentangle non-linear saturation [29,30] and collinearly factorized QCD pictures in a regime where both should be valid descriptions.

The inclusive production of two D mesons provides exciting further opportunities. While the heavy-quarks are predominantly

produced in pairs, the experimental overall reconstruction efficiencies for two D-meson final states are low and, roughly, only one out of million primarily produced double-D events can be reconstructed. Nevertheless, simultaneous production of two D mesons has been observed in p-p [31] and p- \bar{p} [32] collisions. This offers possibilities to test e.g. the heavy-quark vs. heavy-antiquark asymmetries [33] and, in particular, to study the double-parton scattering (DPS) [34–38]. While the formal theory of factorization in DPS has recently advanced significantly [39–41], we still know relatively little of the non-perturbative structure of e.g. the double-parton distributions (dPDFs) [42] which would be required in precise phenomenological applications. Thus, simplifying assumptions concerning DPS have to be made which can lead to apparent shortcomings. For instance, measurements are often interpreted in terms of an *effective cross section* σ_{eff} whose inverse is proportional to the DPS probability. Its value has been observed to differ significantly depending from which observable it is extracted [43]. It is conceivable that this is due to overly simplifying the problem of DPS by ignoring partonic correlations [44–47] or, alternatively, overlooking the contributions of single-parton scattering (SPS) [48]. A complementary approach to hard DPS is provided by Monte-Carlo event generators in which the soft or semi-hard multiparton interactions (MPIs) give rise to the underlying event found necessary to describe the multiplicity distributions in hadronic high-energy collisions [41,49].

The generic prediction is that in proton-nucleus (p-A) collisions the DPS signal gets enhanced in comparison to p-p case, due to

* Corresponding author.

E-mail addresses: ilkka.m.helenius@jyu.fi (I. Helenius), hannu.paukkunen@jyu.fi (H. Paukkunen).

the possibility of proton to interact with two or more nucleons simultaneously [50–54,47,55]. As in p-p collisions, multiple interactions are necessary to explain the multiplicity distributions in collisions involving heavy nuclei [56,57], but a clean experimental confirmation for hard DPS processes is still lacking. As we will conclude later in this letter, it seems realistic that the double D-meson production could provide the first direct evidence of DPS in p-Pb collisions at clearly perturbative scales, and that the signal should be visible already in the collected Run-II data. In reaching this conclusion we have first confronted our QCD framework with the LHCb p-p data and a reasonable agreement there encourages us to apply it to p-Pb collisions. Before getting into the actual results we will first describe our theoretical framework in the next two sections.

2. Double-inclusive production in nuclear collisions

We will estimate the double-inclusive cross sections in collision of two nuclei, A and B , in terms of inclusive per-nucleon SPS cross sections $\sigma_{nn \rightarrow C+X}^{\text{SPS}}$ as

$$\frac{d\sigma_{AB \rightarrow a+b+X}}{d^3\vec{p}^a d^3\vec{p}^b} = AB \left[\frac{d\sigma_{nn \rightarrow a+b+X}^{\text{SPS}}}{d^3\vec{p}^a d^3\vec{p}^b} + \frac{m}{\sigma_{\text{eff}}^{AB}} \frac{d\sigma_{nn \rightarrow a+X}^{\text{SPS}}}{d^3\vec{p}^a} \frac{d\sigma_{nn \rightarrow b+X}^{\text{SPS}}}{d^3\vec{p}^b} \right] \quad (1)$$

where \vec{p}^a and \vec{p}^b refer to the momenta of the produced particles a and b . If a and b are identical particles $m = 1/2$, and $m = 1$ otherwise. In the case of independent partonic interactions, the effective cross section σ_{eff}^{AB} in A - B collision is process independent and can be interpreted as a purely geometric object,

$$\frac{1}{\sigma_{\text{eff}}^{AB}} \equiv \left\{ \frac{1}{\sigma_{\text{eff}}} + \frac{(B-1)}{B^2} \int d^2\vec{B} [T_{nB}(\vec{B})]^2 + \frac{(A-1)}{A^2} \int d^2\vec{B} [T_{nA}(\vec{B})]^2 + \frac{(A-1)(B-1)}{(AB)^2} \int d^2\vec{B} [T_{AB}(\vec{B})]^2 \right\}. \quad (2)$$

Here,

$$\frac{1}{\sigma_{\text{eff}}} = \int d^2\vec{b} [t_{nn}(\vec{b})]^2, \quad (3)$$

where $t_{nn}(\vec{b})$ is the overlap function between two nucleons at fixed impact parameter \vec{b} . In geometric sense, we would write

$$t_{nn}(\vec{b}) \equiv \int_{-\infty}^{\infty} d^2\vec{s} t_n(\vec{s} + \vec{b}/2) t_n(\vec{s} - \vec{b}/2), \quad (4)$$

where $t_n(\vec{s})$ is the transverse profile of nucleons obtained by integrating the density of nucleons ρ^n over the longitudinal spatial component,

$$t_n(\vec{s}) \equiv \int_{-\infty}^{\infty} dz \rho^n(\vec{s}, z). \quad (5)$$

The overlap functions $T_{nA}(\vec{B})$ and $T_{AB}(\vec{B})$ at fixed impact parameter \vec{B} are here defined as [58]

$$T_{nA}(\vec{B}) \equiv \int_{-\infty}^{\infty} d^2\vec{s} t_{nn}(\vec{s} + \vec{B}/2) T_A(\vec{s} - \vec{B}/2) \approx T_A(\vec{B}), \quad (6)$$

where the approximation holds for point-like nucleons, and

$$T_{AB}(\vec{B}) \equiv \int_{-\infty}^{\infty} d^2\vec{s}^A d^2\vec{s}^B T_A(\vec{s}^A) T_B(\vec{s}^B) t_{nn}(\vec{B} + \vec{s}^B - \vec{s}^A) \approx \int_{-\infty}^{\infty} d^2\vec{s} T_A(\vec{s} + \vec{B}/2) T_B(\vec{s} - \vec{B}/2), \quad (7)$$

where $T_A(\vec{S})$ is the standard nuclear thickness function

$$T_A(\vec{S}) \equiv \int_{-\infty}^{\infty} dz \rho^A(\vec{S}, z), \quad (8)$$

and ρ^A denotes the density of nuclei. In our notation, the normalization is

$$\int d^2\vec{s} T_A(\vec{s}) = A. \quad (9)$$

Typically, the DPS contribution in Eq. (1) is derived [51–53,59] by writing the DPS cross section in terms of dPDFs, and assuming that the dPDFs factorize into a product of single-parton PDFs and that the partonic cross sections for the two subprocesses are unrelated and spatially independent. Here, we present a complementary method to obtain Eq. (1) in the approximation where all partonic correlation are neglected. Indeed, in a Glauber-type approach, the total cross section for a collision of nuclei A and B can be written in the impact-parameter space as

$$\sigma_{AB}^{\text{total}} = \int d^2\vec{B} \sum_{k=1}^{AB} \mathcal{P}_k(\vec{B}), \quad (10)$$

where $\mathcal{P}_k(\vec{B})$ is the probability of exactly k nucleon-nucleon interactions at fixed impact parameter \vec{B} ,

$$\mathcal{P}_k(\vec{B}) = \int \left[\prod_{i=1}^A d\vec{S}_i^A \frac{T_A(\vec{S}_i^A)}{A} \right] \left[\prod_{i=1}^B d\vec{S}_i^B \frac{T_B(\vec{S}_i^B)}{B} \right] \times \sum_{\alpha_{11}=0}^1 \dots \sum_{\alpha_{AB}=0}^1 [p_{11}(\alpha_{11}) p_{12}(\alpha_{12}) \dots p_{AB}(\alpha_{AB})] \delta_{k, \alpha_{11} + \dots + \alpha_{AB}}. \quad (11)$$

In this expression, we have defined

$$p_{ij}(\alpha_{ij}) \equiv (t_{ij} \sigma_{nn}^{\text{total}})^{\alpha_{ij}} (1 - t_{ij} \sigma_{nn}^{\text{total}})^{1 - \alpha_{ij}} \quad (12)$$

where t_{ij} is an abbreviation for the overlap function between two nucleons

$$t_{ij} \equiv t_{nn}(\vec{B} + \vec{S}_i^A - \vec{S}_j^B). \quad (13)$$

The second line in Eq. (11) thus corresponds to the probability of getting exactly k nucleon-nucleon interactions (and $AB - k$ missing ones) at fixed geometric configuration. The total cross section in a single nucleon-nucleon collisions $\sigma_{nn}^{\text{total}}$ is given by

$$\sigma_{nn}^{\text{total}} = \int d^2\vec{b} \sum_{k=1}^{\infty} p_k(\vec{b}), \quad (14)$$

where the probability $p_k(\vec{b})$ for k partonic interactions is considered to be Poissonian,

$$p_k(\vec{b}) = \exp\left[-t_{nn}(\vec{b})\sigma_{nn}\right] \frac{\left[t_{nn}(\vec{b})\sigma_{nn}\right]^k}{k!}. \quad (15)$$

The quantity σ_{nn} appearing in Eq. (15) is the integrated inclusive cross sections,

$$\sigma_{nn} = \sum_f \int d\text{PS}_f \frac{d\sigma_{nn \rightarrow f}^{\text{SPS}}}{d\text{PS}_f}, \quad d\text{PS}_f = \prod_{i \in f} d^3\vec{p}_i \quad (16)$$

where the summation is over all *exclusive* final states f . We will always make a distinction between the (intensive) total cross section like $\sigma_{nn}^{\text{total}}$, and (extensive) integrated cross section like σ_{nn} . The double-inclusive cross section can now be written as

$$\frac{d\sigma_{AB \rightarrow a+b+X}}{d\vec{p}^a d\vec{p}^b} = \int d\vec{B} \sum_{k=1}^{AB} \mathcal{P}_k(\vec{B}) \quad (17)$$

$$\prod_{r=1}^k \int d^2\vec{b}_r \sum_{k_r=1}^{\infty} \frac{p_{k_r}(\vec{b}_r)}{\sigma_{nn}^{\text{total}}} \prod_{\ell=1}^{k_r} \left[\frac{1}{\sigma_{nn}} \sum_{f_{r\ell}} \int d\text{PS}_{f_{r\ell}} \frac{d\sigma_{nn \rightarrow f_{r\ell}}^{\text{SPS}}}{d\text{PS}_{f_{r\ell}}} \right] \left[\sum_{i=1}^k \sum_{j=1}^{k_r} \sum_n \delta^{(3)}(\vec{p}^a - \vec{p}_{ij}^{a_n}) \times \begin{cases} 1, & \text{if } a_n \in f_{ij} \\ 0, & \text{if } a_n \notin f_{ij} \end{cases} \right] \left[\sum_{i=1}^k \sum_{j=1}^{k_r} \sum_n \delta^{(3)}(\vec{p}^b - \vec{p}_{ij}^{b_n}) \times \begin{cases} 1, & \text{if } b_n \in f_{ij} \\ 0, & \text{if } b_n \notin f_{ij} \end{cases} \right].$$

In the equation above, each term is a product of the form

$$\mathcal{P}_k(\vec{B}) \times \left[\prod_{r=1}^k p_{k_r}(\vec{b}_r) \right] \times \left[\prod_{\ell=1}^{k_r} \frac{1}{\sigma_{nn}} \frac{d\sigma_{nn \rightarrow f_{r\ell}}^{\text{SPS}}}{d\text{PS}_{f_{r\ell}}} \right], \quad (18)$$

corresponding to the total probability density of having k nucleon-nucleon interactions, each with exactly $k_{r=1, \dots, k}$ partonic interactions resulting with a specific (exclusive) final state $f_{r\ell}$. The last two lines in Eq. (17) simply select those final states which contain the desired particles carrying the momenta \vec{p}^a and \vec{p}^b , and the summation over n accounts for the fact that the final state can contain several a or b particles. With some combinatorics, Eq. (17) simplifies to Eq. (1) when we identify

$$\frac{d\sigma_{nn \rightarrow a+X}^{\text{SPS}}}{d^3\vec{p}^a} \equiv \sum_f \int d\text{PS}_f \frac{d\sigma_{nn \rightarrow f}^{\text{SPS}}}{d\text{PS}_f} \times \sum_i \delta^{(3)}(\vec{p}^a - \vec{p}^{a_i}) \times \begin{cases} 1, & \text{if } a_i \in f \\ 0, & \text{if } a_i \notin f \end{cases}, \quad (19)$$

and

$$\frac{d\sigma_{nn \rightarrow a+b+X}^{\text{SPS}}}{d^3\vec{p}^a d^3\vec{p}^b} \equiv \sum_f \int d\text{PS}_f \frac{d\sigma_{nn \rightarrow f}^{\text{SPS}}}{d\text{PS}_f} \times \sum_i \delta^{(3)}(\vec{p}^a - \vec{p}^{a_i}) \times \begin{cases} 1, & \text{if } a_i \in f \\ 0, & \text{if } a_i \notin f \end{cases} \times \sum_i \delta^{(3)}(\vec{p}^b - \vec{p}^{b_i}) \times \begin{cases} 1, & \text{if } b_i \in f \\ 0, & \text{if } b_i \notin f \end{cases}. \quad (20)$$

From the same formalism, also three-particle (in general n -particle) inclusive cross sections [60,61,53] can be derived.

3. Perturbative-QCD framework for open heavy flavour

In this paper we will be mostly concerned in the D-meson production at $p_T > 3$ GeV, which is the kinematic region considered in the LHCb double-D measurement [31]. In this region the inclusive production of D mesons can be reliably described within general-mass variable-flavour-number scheme (GM-VNFS). Schematically, the cross sections are convolutions of PDFs $f_i(x, \mu_{\text{fact}}^2)$, partonic cross sections $d\hat{\sigma}$, and fragmentation functions (FFs) $D_{k \rightarrow h}(z, \mu_{\text{frag}}^2)$,

$$d\sigma_{nn \rightarrow a+X}^{\text{SPS}} = \sum_{ijk} f_i(\mu_{\text{fact}}^2) \otimes d\hat{\sigma}_{ij \rightarrow k+X}(\mu_{\text{fact}}^2, \mu_{\text{ren}}^2, \mu_{\text{frag}}^2) \otimes f_j(\mu_{\text{fact}}^2) \otimes D_{k \rightarrow a}(\mu_{\text{frag}}^2). \quad (21)$$

For single-inclusive D-meson production this has been considered at next-to-leading order (NLO) QCD first in Ref. [14] within the so-called SACOT scheme [62,63]. In the SACOT scheme, the partonic cross sections for contributions in which the partonic subprocess is initiated by a charm quark or the fragmenting parton is a light one, are independent of the charm-quark mass m_{charm} . This leads, in general, to diverging cross sections towards $p_T \rightarrow 0$. In an alternative SACOT- m_T scheme [15] this unphysical behaviour is resolved by accounting for the underlying kinematic constraint of heavy-quark production. In this work we use the SACOT- m_T variant, albeit in the considered $p_T > 3$ GeV region, both schemes should be equivalent within the scale uncertainties. Our default choice for factorization (μ_{fact}), fragmentation (μ_{frag}) and renormalization (μ_{ren}) scales is $\mu_{\text{fact}}^2 = \mu_{\text{frag}}^2 = \mu_{\text{ren}}^2 = p_T^2 + m_{\text{charm}}^2$, where p_T refers to the D-meson transverse momentum.

The SPS contribution in which the two D mesons, h_1 and h_2 , are simultaneously produced is of the form,

$$d\sigma_{pp \rightarrow a+b+X}^{\text{SPS}} = \sum_{ijkl} f_i(\mu_{\text{fact}}^2) \otimes d\hat{\sigma}_{ij \rightarrow k+l+X}(\mu_{\text{fact}}^2, \mu_{\text{ren}}^2, \mu_{\text{frag}}^2) \otimes f_j(\mu_{\text{fact}}^2) \otimes D_{k \rightarrow a}(\mu_{\text{frag}}^2) \otimes D_{l \rightarrow b}(\mu_{\text{frag}}^2). \quad (22)$$

For this process, no GM-VNFS calculation is available. Thus, we will resort to the zero-mass approximation available in the NLO DIPHOX [64] (v.1.2) code. Taking into account the large scale uncertainties, this approximation should be sufficiently precise in the considered $p_T > 3$ GeV region. However, the kinematical cuts applied in the considered LHCb measurement [31] ($p_T > 3$ GeV and $2 < y < 4$) include also a problematic configuration in which the two D mesons are collinear. In a full SACOT- m_T description this contribution would be finite, scaling as $\log(m_{\text{charm}}^2)$ where the remaining $\log(m_{\text{charm}}^2)$ terms would still need to be resummed via *di-hadron FFs* [65]. In a zero-mass calculation, however, the cross sections diverge in the collinear configuration. Here, as a proxy for the full SACOT- m_T treatment, we have regulated our calculations by imposing a physical cut $(\hat{p}_1 + \hat{p}_2)^2 > 4m_{\text{charm}}^2$ for the fragmenting partons' four momenta $\hat{p}_{1,2}$. Our central choice for the QCD scales here is the average p_T of the produced two D mesons.

The dominant uncertainty in our calculations comes from the unknown higher-order (NNLO and beyond) contributions. As usual, we estimate the potential size of these corrections by varying the QCD scales as

$$0.5 \leq \frac{\mu_{\text{fact}}}{\mu_{\text{ren}}} \leq 2, \quad 0.5 \leq \frac{\mu_{\text{frag}}}{\mu_{\text{ren}}} \leq 2, \quad (23)$$

around the central scale choices and finding the combinations that give the highest and lowest prediction for each considered observable. As default, we do the scale variations in sync for




Fig. 1. The LHCb inclusive D-meson and Λ_c^\pm data [1] in $\sqrt{s} = 7$ TeV p-p collisions compared with SACOT- m_T calculation using the KKK08 [66] and BKK05 [67] FFs. The gray bands show the scale uncertainty with central KKK08/BKK05 FFs. The central predictions with FFs based on BELLE (purple short dashed), OPAL (green dashed dotted), and CLEO (red dotted) data are displayed as well. The PYTHIA predictions (blue long dashed) are shown also.

the two contributions in Eq. (1), the single-inclusive and double-inclusive SPS cross sections (17 scale configurations in total). We use NNPDF3.1pch PDFs [68] in which the intrinsic charm component is zero at the mass threshold $\mu_{\text{fact}} = m_c = 1.51$ GeV. The fragmentation functions for D^0 and D^+ are taken from the KKK08 analysis [66] (see Ref. [69] for a very recent alternative). The KKK08 FFs have been fitted to e^+e^- data from different experiments. We have checked that while the fits to BELLE [70] and OPAL [71] data give essentially equally good descriptions of the inclusive LHCb D^0 and D^+ cross sections at $\sqrt{s} = 7$ TeV [1], the FFs fitted to CLEO data [72] clearly overshoot the LHCb data at high p_T . This is demonstrated in the upper panels of Fig. 1. However, we have found that the D^0 -to- D^+ ratios which are almost exclusively sensitive to the FFs are clearly best described by the OPAL variant, which also gives a better description than the BELLE FFs of the CMS midrapidity data [8] at very-high p_T [73]. Thus, in this paper, we adopt the OPAL FFs from the KKK08 package. For D_s^\pm and Λ_c^\pm FFs we use BKK05 [67] analysis. While the LHCb and ALICE single-inclusive D_s^\pm data [1,6] are well consistent with these FFs, the Λ_c^\pm data [1] are underestimated by the BKK05 Λ_c^\pm FFs.

Our comparisons with the LHCb data on D_s^\pm and Λ_c^\pm are shown in the bottom panels of Fig. 1.

The KKK08 and BKK05 FFs do not discriminate between charge-conjugate states, but are given as a sum. (e.g. $D_i^{D^0+D^0}$). In what follows, however, we will need the D-meson FFs one by one. Taking the D^0 states here as an example, we will use the following prescription for the charm-quark containing state,

$$D_c^{D^0} = D_{c/\bar{c}}^{D^0+\bar{D}^0}, \quad (24)$$

$$D_{\bar{c}}^{D^0} = 0, \quad (25)$$

$$D_i^{D^0} = \frac{1}{2} D_i^{D^0+\bar{D}^0}, \quad i \neq c/\bar{c}, \quad (26)$$

and an analogous one for the antiquark-containing state,

$$D_c^{\bar{D}^0} = 0, \quad (27)$$

$$D_{\bar{c}}^{\bar{D}^0} = D_{c/\bar{c}}^{D^0+\bar{D}^0}, \quad (28)$$

$$D_i^{\bar{D}^0} = \frac{1}{2} D_i^{D^0+\bar{D}^0}, \quad i \neq c/\bar{c}. \quad (29)$$

In addition to the NLO QCD framework described above, we present the predictions from PYTHIA 8 Monte-Carlo event generator using the standard ‘‘Monash 2013 tune’’ [74]. A sample of minimum-bias events were generated, including also MPIs, from which the different D-meson combinations within the LHCb acceptance were picked up to obtain the cross sections for each pair. In line with the LHCb measurements, each pair of D mesons is counted separately. In Fig. 1 we also show the PYTHIA predictions for the inclusive D mesons and Λ_c^\pm , generated with the provided Rivet analysis [75]. In general, the PYTHIA setup overpredicts the LHCb D-meson measurements, and the disagreement is stronger for D^\pm and D_S^\pm than for D^0 . A similar behaviour has been recently observed in the case of jets containing a D^0 meson [76]. The measured Λ_c^\pm cross sections are, in turn, underestimated by PYTHIA. In the Monash tune the parameters related to charm fragmentation were constrained using a limited set of LEP data. Partly the interpretation of these data is hindered by the large feed-down from B-mesons. Furthermore, the data is not sensitive to $g \rightarrow c\bar{c}$ branchings that are abundant at the LHC. Thus the observed disagreement could potentially be cured by re-tuning the relevant parameters using a larger sample of charm-production data from LEP, HERA and LHC.

4. Results

We will now compare our results for double D-meson production with the LHCb p-p data [31], and make predictions for p-Pb collisions. As for σ_{eff} , we will consider the variation $10 \text{ mb} < \sigma_{\text{eff}} < 25 \text{ mb}$ which is roughly the range deduced from jet, W^\pm and photon measurements [43]. The uncertainty estimates shown in the plots combine the scale uncertainty and the variation in σ_{eff} .

4.1. p-p collisions

In the case of p-p collision, Eqs. (1) and (2) reduce to

$$\frac{d\sigma_{pp \rightarrow a+b+X}}{d\bar{p}^a d\bar{p}^b} = \frac{d\sigma_{pp \rightarrow a+b+X}^{\text{sps}}}{d\bar{p}^a d\bar{p}^b} + \frac{m}{\sigma_{\text{eff}}} \frac{d\sigma_{pp \rightarrow a+X}^{\text{sps}}}{d\bar{p}^a} \frac{d\sigma_{pp \rightarrow b+X}^{\text{sps}}}{d\bar{p}^b}. \quad (30)$$

Our results for the integrated cross sections within the LHCb acceptance are shown in Fig. 2. For the opposite-sign D mesons (upper panel), the SPS contribution is clearly larger than the DPS one, and the agreement with the data is very good. The measured systematics among different combinations of D (and Λ_c) species is well reproduced by the used set of FFs. As the cross section accumulates from the lower end of the considered p_T range, the scale uncertainty is sizable and dominates over the variation in σ_{eff} . For like-sign final states (lower panel) the DPS becomes the dominant production mechanism. Again, the calculation agrees with the data within the scale uncertainties, though our central scale choice seem to somewhat overestimate the cross sections. The disagreement between the data and PYTHIA results is considerably larger than in the single-inclusive case (Fig. 1). Apart from pairs including Λ_c the predicted cross sections are 4–8 times higher than the data. For pairs including Λ_c the systematic is again the opposite.

More insight can be obtained from Fig. 3 where we show cross-section ratios. The upper panel shows ratios between the double like-sign vs. opposite-sign cross sections,

$$\sigma^{ab} / \sigma^{a\bar{b}} \equiv \frac{\sigma_{pp \rightarrow a+b+X}}{\sigma_{pp \rightarrow a+\bar{b}+X}}. \quad (31)$$

These measure essentially the ratio between the DPS and SPS contributions. There is clearly a fair data-to-theory agreement within




Fig. 2. The integrated double-D cross sections for opposite-sign (upper panel) and like-sign (lower panel) cases. The coloured bands denote the combined scale and σ_{eff} uncertainty in NLO calculations. The inner darker bands include only the variation in σ_{eff} . The PYTHIA predictions are shown as blue dashed lines. The data are from Ref. [31].

the scale and σ_{eff} uncertainties. Our central predictions somewhat overestimate the measured values which is consistent with Fig. 2. The scale uncertainties do not cancel out since the partonic channels for like-sign and opposite-sign production are different (e.g. $c\bar{c}$ pair production is significant for $D^0 D^0$ final state but not for $D^0 D^0$). Interestingly, the PYTHIA results are in excellent agreement with the LHCb data even though the absolute cross sections are way off. Since the numerator in Eq. (31) is sensitive to DPS (or MPIs in general), we conclude that the good agreement here suggest that the inconsistencies observed in Fig. 2 are indeed due to poorly-constrained charm fragmentation, rather than the MPI modelling in PYTHIA [77–79].

The bottom panel of Fig. 3 shows ratios

$$\sigma^a \sigma^b / \sigma^{ab} \equiv m \frac{\sigma_{pp \rightarrow a+X} \times \sigma_{pp \rightarrow b+X}}{\sigma_{pp \rightarrow a+b+X}}. \quad (32)$$

From Eq. (30) we see that in the absence of SPS, this ratio would be equal to σ_{eff} , but if there is a contribution from SPS, the ratio will be below σ_{eff} . In general, our predictions for the opposite-sign case match very well with the data, but tend to underestimate the measured like-sign ratios. This is well in line with our earlier observations and also here a better overall agreement would be obtained if the DPS cross section would be somewhat smaller. Thus, the double-charm production data would prefer a somewhat larger phenomenological σ_{eff} than what other measurements indicate [43]. We would like to stress here that the scale uncertainties have been computed by varying the scale choices (17 combinations) simultaneously in the numerator and denominator of Eq. (32). However, since the processes in the numerator and denominator are not the same it is not out of question that the scale choices/variations could be made independently. This would sig-




Fig. 3. Upper panel: like-sign vs. opposite-sign ratios, see Eq. (31). Lower panel: Product of two single-inclusive D-meson cross sections divided by the double-D cross sections, see Eq. (32). The coloured bars denote the combined scale and σ_{eff} uncertainty, and the inner darker bands include only the variation in σ_{eff} . The dashed lines correspond to what PYTHIA predicts. The upper set of bands/lines/data points correspond to like-sign D mesons, and the lower set to opposite-sign combinations.

nificantly increase the scale uncertainties in the bottom panel of Fig. 3. In our framework presented above the underlying idea is to rather take the variation in σ_{eff} extracted from other processes as an estimate for the associated uncertainty. The PYTHIA predictions are here well compatible with our NLO calculations, though they somewhat undershoot the measured ratios both for like- and opposite-sign ratios. This further supports our conclusion that the disagreement observed in Figs. 1 and 2 arise from the fragmentation scheme in PYTHIA.

4.2. p-Pb collisions

The reasonable description of the p-p data gives us confidence to apply the framework in p-A collisions. We note that the GM-VFNS framework used here has recently been contrasted against LHCb data [26] on single-inclusive D-meson production [17] finding and a good agreement when modern nuclear PDFs are used. This indicates that such data can be well described in a collinear-factorization based framework without e.g. photoproduction contributions arising from an amplified photon flux of Pb, or any non-linear saturation effects. We take this as a further evidence that our framework of double-inclusive cross sections with no photoproduction component nor saturation can be applied also to p-Pb collisions. In this case, Eqs. (1) and (2) reduce to

$$\frac{d\sigma_{pA \rightarrow a+b+X}}{d\vec{p}^a d\vec{p}^b} = A \left[\frac{d\sigma_{nn \rightarrow a+b+X}^{\text{sps}}}{d\vec{p}^a d\vec{p}^b} + \frac{m}{\sigma_{\text{eff}}^{\text{pA}}} \frac{d\sigma_{nn \rightarrow a+X}^{\text{sps}}}{d\vec{p}^a} \frac{d\sigma_{nn \rightarrow b+X}^{\text{sps}}}{d\vec{p}^b} \right] \quad (33)$$




Fig. 4. Upper panel: Integrated cross sections for $D^0\bar{D}^0$ and D^0D^0 cross sections in p-Pb collisions within the LHCb acceptance at $\sqrt{s} = 8.16$ TeV. The coloured bars denote the combined scale and σ_{eff} uncertainty, and the inner darker bands include only the variation in σ_{eff} . The LHCb projections correspond to 12.2 nb^{-1} (forward) and 18.6 nb^{-1} (backward) luminosities assuming the predicted central value and overall efficiency of 1.2×10^{-6} . Lower panel: A sketch of the relative azimuthal-angle dependence in p-p and projected p-Pb collisions.

with

$$\frac{1}{\sigma_{\text{eff}}^{\text{pA}}} \equiv \frac{1}{\sigma_{\text{eff}}} \times \left\{ 1 + \sigma_{\text{eff}} \frac{A-1}{A^2} \int d^2\vec{B} \left[T_{nA}(\vec{B}) \right]^2 \right\}. \quad (34)$$

The impact-parameter integral for $A = 208$ (Pb) gives

$$\int d^2\vec{B} \left[T_{n\text{Pb}}(\vec{B}) \right]^2 \approx 31.66 \text{ mb}^{-1} \quad (35)$$

taking $d = 0.54$ fm and $r = 6.49$ fm in the Woods-Saxon profile,

$$\rho^A(\vec{s}, z) = n_0 \left[1 + \exp \left(\frac{\sqrt{s^2 + z^2} - r}{d} \right) \right]^{-1}, \quad (36)$$

and fixing n_0 by the normalization condition of Eq. (9). With $\sigma_{\text{eff}} = 10 \dots 25$ mb, we find

$$\frac{1}{\sigma_{\text{eff}}^{\text{pPb}}} \approx \frac{2.5 \dots 4.8}{\sigma_{\text{eff}}} \quad (37)$$

in full consistency e.g. with Ref. [51]. That is, the DPS signal is enhanced approximately by a factor of three in comparison to p-p scattering. We note that effects of nucleon-nucleon correlations on the integral in Eq. (35) are expected to be small for a large nucleus like Pb [80]. Our results for the integrated cross sections within the LHCb kinematics are shown in Fig. 4. Here, we have only considered D^0 production which has the largest cross sections, see Fig. 2, and the y acceptance refers to that in the center-of-mass frame of the p-Pb collision. When computing the per-nucleon cross

sections $\sigma_{nn \rightarrow a+b+X}^{\text{sps}}$ and $\sigma_{nn \rightarrow a/b+X}^{\text{sps}}$, we have used the EPPS16 nuclear modifications [81] for Pb. At the LHCb kinematics this leads to a $\sim 20\%$ suppression for p-Pb (forward) SPS cross sections, but since this is squared in DPS contribution, the suppression can reach $\sim 40\%$ in DPS case. For Pb-p configuration (backward) the nuclear-PDF effects are smaller. In comparison to the p-p case in Fig. 2 the impact of enhanced DPS contribution is clear: Whereas in p-p case the DPS contribution to the opposite-sign yield was rather small in comparison to SPS, in p-Pb collisions the two are comparable. For the like-sign yields the SPS contribution in p-Pb collisions is entirely overpowered by the DPS part, whereas in the p-p case the SPS still had a 20% contribution or so. Due to the additional contribution from the $T_{nA}^2(\vec{B})$ integral in Eq. (34), the variation in σ_{eff} plays only a minor role as indicated in Fig. 4. The $\sim 30\%$ differences between forward and backward cross sections are due to the EPPS16 nuclear effects. Thus, by a suitable measurement where other theoretical uncertainties would cancel out, e.g. a forward-to-backward ratio for double D-meson production, further constraints for nuclear PDFs could, perhaps, be obtained.

An interesting question is whether these cross sections are large enough to be measured with the already collected Run-II data. In Run-II data taking the luminosities collected by the LHCb were 12.2 nb^{-1} for p-Pb (forward) and 18.6 nb^{-1} Pb-p (backward) collisions [27]. The overall detection efficiency ϵ for $D^0\bar{D}^0$ and D^0D^0 final states in the LHCb p-p measurement [31] was approximately $\epsilon \approx 1.2 \times 10^{-6}$. Using these luminosities and efficiencies with our central theoretical predictions we calculate the expected number of events N from which the statistical uncertainty is obtained as \sqrt{N}/N . These estimates are also shown in Fig. 4. Within the scale uncertainties we expect approximately $10 \dots 40 D^0\bar{D}^0$ pairs in p-Pb collisions (forward), and $20 \dots 80$ in Pb-p configuration (backward). For the like-sign case the corresponding numbers are $2 \dots 20 D^0D^0$ pairs in p-Pb collisions (forward), and $4 \dots 40$ in Pb-p configuration (backward). Thus, we are led to conclude that the double D-meson production – at least the opposite-sign case – should be observable at the LHCb with the Run-II luminosity. Lowering the minimum- p_T cut below 3 GeV would easily increase the yields to a definitely measurable level, but towards lower p_T our predictions become increasingly uncertain.

As is well known, the increased importance of the DPS contribution in p-Pb collisions may significantly affect the kinematic distributions [82,83]. Particularly interesting observable is the relative azimuthal-angle $\Delta\phi$ distribution of the two D mesons [84]. In p-p collisions [31] the $\Delta\phi$ distribution for $D^0\bar{D}^0$ peaks at $\Delta\phi = 0$ for the logarithmically enhanced $g \rightarrow c\bar{c}$ splitting, and at $\Delta\phi = \pi$ due to the leading-order contributions that are back-to-back in transverse plane. These are commonly referred to as the *near-side peak* and the *away-side peak*, respectively. The disappearance of the away-side peak has long been predicted to be the smoking gun of saturation physics [85,86]. However, the enhanced DPS contribution in p-Pb collisions will generate a $\Delta\phi$ -independent contribution which levels off these peaks. Thus, a flatter away-side peak in p-Pb in comparison to p-p cannot directly be interpreted as being due to saturation physics. What we would like to estimate here is the difference between the $\Delta\phi$ dependence in p-Pb and p-p collisions just due to the enhanced $\Delta\phi$ -independent DPS background in p-Pb which shrouds the nontrivial $\Delta\phi$ dependence originating from the SPS contribution. Unfortunately our GM-VFNS framework cannot reliably predict the $\Delta\phi$ dependence near $\Delta\phi = \pi$ but a soft-gluon resummation encoded e.g. in parton showers, would be required. To estimate the effect, we have fitted the $\Delta\phi$ dependence of the LHCb $D^0\bar{D}^0$ data [31] in p-p collisions assuming a negligible contribution from DPS. This assumption is consistent both with

our results (see Fig. 2) and also with Ref. [84] where it has been shown that for $0 \lesssim \Delta\phi \lesssim \pi/2$ the fixed-order QCD quite correctly predicts the $\Delta\phi$ dependence. Our estimate for the $\Delta\phi$ dependence in p-Pb is then a linear combination

$$\frac{d\sigma^{\text{pPb}}}{d\Delta\phi} \propto \frac{d\sigma^{\text{pp}}}{d\Delta\phi} \Big|_{\text{fitted}} + \beta. \quad (38)$$

The non-trivial $\Delta\phi$ dependence obviously comes from the first term (representing the SPS contribution) while β stands for the DPS background and its value is determined by the relative importance of integrated SPS and DPS cross sections (given by our GM-VFNS calculation) in p-Pb collisions. Depending on the scale choices, we estimate the DPS contribution to be roughly between $20\% \dots 40\%$. The resulting projection for the $\Delta\phi$ dependence is shown in the lower panel of Fig. 4 where the coloured band comes from the scale and σ_{eff} uncertainties. We see that both the near- and away-side peaks become less pronounced in p-Pb than what they are in p-p. Thus, we can confirm that the DPS contributions should be considered when interpreting the possible (probable?) weakening of the away-side two-particle correlations in terms of e.g. saturation physics. In fact, only stronger suppression than what our estimate in Fig. 4 indicates could be interpreted to originate from saturation phenomena. Since the DPS contribution should be nearly the same for $D^0\bar{D}^0$ and D^0D^0 final states (in our calculations they are equal), the difference

$$\frac{d\sigma_{\text{pPb} \rightarrow D^0\bar{D}^0+X}}{d\Delta\phi} - \frac{d\sigma_{\text{pPb} \rightarrow D^0D^0+X}}{d\Delta\phi} \approx \frac{d\sigma_{\text{pPb} \rightarrow D^0\bar{D}^0+X}^{\text{sps}}}{d\Delta\phi} \quad (39)$$

should serve to subtract the ‘‘pedestal’’ DPS yield in a rather model-independent way and, as indicated above, correspond very closely to the SPS contribution in $D^0\bar{D}^0$ production. Even though the presented calculations are for double D-meson production, we would expect a similar reduction of the near- and away-side peaks also for light-flavour hadrons (such as $\pi^+\pi^-$) in due to enhanced DPS contribution in p-Pb collisions.

5. Summary

We have explored the double-inclusive D-meson production at the LHC with focus on the forward LHCb kinematics. The contributions of double-parton scatterings were included in the approximation of independent parton-parton collisions, and the required single-parton cross sections were computed within the collinearly factorized QCD at an NLO level. We confronted our predictions with the LHCb p-p data finding a good, or least an acceptable agreement within the QCD scale uncertainties and reasonable variation in the effective cross σ_{eff} . As a whole, the LHCb data would prefer a rather large σ_{eff} compared to values derived from other final states. We also compared the LHCb p-p data with PYTHIA predictions. We found that the absolute cross sections for single- and double-inclusive open-charm production are not well reproduced by the widely used Monash tune. However, the cross-section ratios, which are less sensitive to the details in charm fragmentation, are described equally well or even better than what our NLO calculations do. Since the ratios are more sensitive to the multi-parton dynamics than the heavy-quark fragmentation model, it seems that the latter will need some further tuning to establish an agreement with the absolute cross sections.

In addition, we applied our framework to the case of p-Pb collisions in which the contribution from double-parton scattering is predicted to get significantly enhanced due to multiple nucleon-nucleon interactions. Our calculations, accounting for realistic reconstruction efficiencies, indicate that the yields should be high

enough to be measured with the already-collected LHC Run-II data at the LHCb. This should provide a clear evidence for the hard double-parton scattering in p-Pb. As the contributions from single- and double-parton scatterings to opposite-sign double-D pair become comparable, also the azimuthal correlations are significantly altered. Therefore it seems necessary to take the double-parton scattering component into account when interpreting e.g. the possible complete or probable partial disappearance of the away-side peak.

Acknowledgements

We thank Michael Winn for clarifying us details of the LHCb p-p measurements and Peter Skands for discussions related to the Monash-tune applied in PYTHIA simulations. Our work was financed by the Academy of Finland, projects n.o. 297058 and 308301. The Finnish IT Center for Science (CSC) is acknowledged for computing resources within the project jyy2580 of T. Lappi.

References

- [1] R. Aaij, et al., Prompt charm production in pp collisions at $\sqrt{s}=7$ TeV, Nucl. Phys. B 871 (2013) 1–20, <https://doi.org/10.1016/j.nuclphysb.2013.02.010>, arXiv:1302.2864.
- [2] R. Aaij, et al., Measurements of prompt charm production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV, J. High Energy Phys. 03 (2016) 159, [https://doi.org/10.1007/JHEP03\(2016\)159](https://doi.org/10.1007/JHEP03(2016)159), arXiv:1510.01707, Erratum: J. High Energy Phys. 05 (2017) 074, [https://doi.org/10.1007/JHEP05\(2017\)074](https://doi.org/10.1007/JHEP05(2017)074), [https://doi.org/10.1007/JHEP09\(2016\)013](https://doi.org/10.1007/JHEP09(2016)013).
- [3] R. Aaij, et al., Measurements of prompt charm production cross-sections in pp collisions at $\sqrt{s} = 5$ TeV, J. High Energy Phys. 06 (2017) 147, [https://doi.org/10.1007/JHEP06\(2017\)147](https://doi.org/10.1007/JHEP06(2017)147), arXiv:1610.02230.
- [4] R. Aaij, et al., Measurement of the B^\pm production cross-section in pp collisions at $\sqrt{s} = 7$ and 13 TeV, J. High Energy Phys. 12 (2017) 026, [https://doi.org/10.1007/JHEP12\(2017\)026](https://doi.org/10.1007/JHEP12(2017)026), arXiv:1710.04921.
- [5] S. Acharya, et al., Measurement of D-meson production at mid-rapidity in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 77 (8) (2017) 550, <https://doi.org/10.1140/epjc/s10052-017-5090-4>, arXiv:1702.00766.
- [6] S. Acharya, et al., Measurement of D^0 , D^+ , D^{*+} and D_s^\pm production in pp collisions at $\sqrt{s} = 5.02$ TeV with ALICE, arXiv:1901.07979, 2019.
- [7] G. Aad, et al., Measurement of $D^{*\pm}$, D^\pm and D_s^\pm meson production cross sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, Nucl. Phys. B 907 (2016) 717–763, <https://doi.org/10.1016/j.nuclphysb.2016.04.032>, arXiv:1512.02913.
- [8] A.M. Sirunyan, et al., Nuclear modification factor of D^0 mesons in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 782 (2018) 474–496, <https://doi.org/10.1016/j.physletb.2018.05.074>, arXiv:1708.04962.
- [9] A. Andronic, et al., Heavy-flavour and quarkonium production in the LHC era: from proton–proton to heavy-ion collisions, Eur. Phys. J. C 76 (3) (2016) 107, <https://doi.org/10.1140/epjc/s10052-015-3819-5>, arXiv:1506.03981.
- [10] P. Nason, S. Dawson, R.K. Ellis, The one particle inclusive differential cross-section for heavy quark production in hadronic collisions, Nucl. Phys. B 327 (1989) 49–92, [https://doi.org/10.1016/0550-3213\(89\)90286-1](https://doi.org/10.1016/0550-3213(89)90286-1), Erratum: Nucl. Phys. B 335 (1990) 260, [https://doi.org/10.1016/0550-3213\(90\)90180-L](https://doi.org/10.1016/0550-3213(90)90180-L).
- [11] M.L. Mangano, P. Nason, G. Ridolfi, Heavy quark correlations in hadron collisions at next-to-leading order, Nucl. Phys. B 373 (1992) 295–345, [https://doi.org/10.1016/0550-3213\(92\)90435-E](https://doi.org/10.1016/0550-3213(92)90435-E).
- [12] M. Cacciari, M. Greco, P. Nason, The P(T) spectrum in heavy flavor hadroproduction, J. High Energy Phys. 05 (1998) 007, <https://doi.org/10.1088/1126-6708/1998/05/007>, arXiv:hep-ph/9803400.
- [13] S. Frixione, P. Nason, B.R. Webber, Matching NLO QCD and parton showers in heavy flavor production, J. High Energy Phys. 08 (2003) 007, <https://doi.org/10.1088/1126-6708/2003/08/007>, arXiv:hep-ph/0305252.
- [14] B.A. Kniehl, G. Kramer, I. Schienbein, H. Spiesberger, Inclusive D^{*+} production in p anti-p collisions with massive charm quarks, Phys. Rev. D 71 (2005) 014018, <https://doi.org/10.1103/PhysRevD.71.014018>, arXiv:hep-ph/0410289.
- [15] I. Helenius, H. Paukkunen, Revisiting the D-meson hadroproduction in general-mass variable flavour number scheme, J. High Energy Phys. 05 (2018) 196, [https://doi.org/10.1007/JHEP05\(2018\)196](https://doi.org/10.1007/JHEP05(2018)196), arXiv:1804.03557.
- [16] D.P. Anderle, T. Kaufmann, M. Stratmann, F. Ringer, I. Vitev, Using hadron-in-jet data in a global analysis of D^* fragmentation functions, Phys. Rev. D 96 (3) (2017) 034028, <https://doi.org/10.1103/PhysRevD.96.034028>, arXiv:1706.09857.
- [17] K.J. Eskola, I. Helenius, P. Paakkinen, H. Paukkunen, A QCD analysis of LHCb D-meson data in p+Pb collisions, arXiv:1906.02512, 2019.
- [18] O. Zenaiev, et al., Impact of heavy-flavour production cross sections measured by the LHCb experiment on parton distribution functions at low x, Eur. Phys. J. C 75 (8) (2015) 396, <https://doi.org/10.1140/epjc/s10052-015-3618-z>, arXiv:1503.04581.
- [19] R. Gauld, J. Rojo, L. Rottoli, J. Talbert, Charm production in the forward region: constraints on the small-x gluon and backgrounds for neutrino astronomy, J. High Energy Phys. 11 (2015) 009, [https://doi.org/10.1007/JHEP11\(2015\)009](https://doi.org/10.1007/JHEP11(2015)009), arXiv:1506.08025.
- [20] R. Gauld, J. Rojo, Precision determination of the small-x gluon from charm production at LHCb, Phys. Rev. Lett. 118 (7) (2017) 072001, <https://doi.org/10.1103/PhysRevLett.118.072001>, arXiv:1610.09373.
- [21] R. Gauld, Forward D predictions for pPb collisions, and sensitivity to cold nuclear matter effects, Phys. Rev. D 93 (1) (2016) 014001, <https://doi.org/10.1103/PhysRevD.93.014001>, arXiv:1508.07629.
- [22] J.-P. Lansberg, H.-S. Shao, Towards an automated tool to evaluate the impact of the nuclear modification of the gluon density on quarkonium, D and B meson production in proton-nucleus collisions, Eur. Phys. J. C 77 (1) (2017) 1, <https://doi.org/10.1140/epjc/s10052-016-4575-x>, arXiv:1610.05382.
- [23] A. Kusina, J.-P. Lansberg, I. Schienbein, H.-S. Shao, Gluon shadowing in heavy-flavor production at the LHC, Phys. Rev. Lett. 121 (5) (2018) 052004, <https://doi.org/10.1103/PhysRevLett.121.052004>, arXiv:1712.07024.
- [24] M. Cacciari, M.L. Mangano, P. Nason, Gluon PDF constraints from the ratio of forward heavy-quark production at the LHC at $\sqrt{s} = 7$ and 13 TeV, Eur. Phys. J. C 75 (12) (2015) 610, <https://doi.org/10.1140/epjc/s10052-015-3814-x>, arXiv:1507.06197.
- [25] J. Adam, et al., D-meson production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. C 94 (5) (2016) 054908, <https://doi.org/10.1103/PhysRevC.94.054908>, arXiv:1605.07569.
- [26] R. Aaij, et al., Study of prompt D^0 meson production in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV, J. High Energy Phys. 10 (2017) 090, [https://doi.org/10.1007/JHEP10\(2017\)090](https://doi.org/10.1007/JHEP10(2017)090), arXiv:1707.02750.
- [27] R. Aaij, et al., Measurement of B^+ , B^0 and Λ_b^0 production in pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV, Phys. Rev. D 99 (5) (2019) 052011, <https://doi.org/10.1103/PhysRevD.99.052011>, arXiv:1902.05599.
- [28] R. Aaij, et al., First measurement of charm production in fixed-target configuration at the LHC, Phys. Rev. Lett. 122 (13) (2019) 132002, <https://doi.org/10.1103/PhysRevLett.122.132002>, arXiv:1810.07907.
- [29] H. Fujii, K. Watanabe, Heavy quark pair production in high energy pA collisions: open heavy flavors, Nucl. Phys. A 920 (2013) 78–93, <https://doi.org/10.1016/j.nuclphysa.2013.10.006>, arXiv:1308.1258.
- [30] B. Ducloué, T. Lappi, H. Mäntysaari, Forward J/ψ and D meson nuclear suppression at the LHC, Nucl. Part. Phys. Proc. 289–290 (2017) 309–312, <https://doi.org/10.1016/j.nuclphysbps.2017.05.071>, arXiv:1612.04585.
- [31] R. Aaij, et al., Observation of double charm production involving open charm in pp collisions at $\sqrt{s} = 7$ TeV, J. High Energy Phys. 06 (2012) 141, [https://doi.org/10.1007/JHEP06\(2012\)141](https://doi.org/10.1007/JHEP06(2012)141), Addendum: J. High Energy Phys. 03 (2014) 108, [https://doi.org/10.1007/JHEP03\(2014\)108](https://doi.org/10.1007/JHEP03(2014)108), arXiv:1205.0975.
- [32] B. Reiser, J. Lewis, T. Liu, M.J. Kim, K. Chung, J. Russ, P. Lukens, Charm production studies at CDF, Nucl. Phys. Proc. Suppl. 170 (2007) 243–247, <https://doi.org/10.1016/j.nuclphysbps.2007.05.053>.
- [33] R. Gauld, U. Haisch, B.D. Pecjak, E. Re, Beauty-quark and charm-quark pair production asymmetries at LHCb, Phys. Rev. D 92 (2015) 034007, <https://doi.org/10.1103/PhysRevD.92.034007>, arXiv:1505.02429.
- [34] I. Belyaev, D. Savrina, Study of double parton scattering processes with heavy quarks, Adv. Ser. Dir. High Energy Phys. 29 (2018) 141–157, arXiv:1711.10877.
- [35] B. Blok, M. Strikman, Open charm production in Double Parton Scattering processes in the forward kinematics, Eur. Phys. J. C 76 (12) (2016) 694, <https://doi.org/10.1140/epjc/s10052-016-4551-5>, arXiv:1608.00014.
- [36] E.R. Cazaroto, V.P. Gonçalves, F.S. Navarra, Heavy quark production in pA collisions: the double parton scattering contribution, Mod. Phys. Lett. A 33 (25) (2018) 1850141, <https://doi.org/10.1142/S0217732318501419>, arXiv:1607.04023.
- [37] R. Maciula, V.A. Saleev, A.V. Shipilova, A. Szczurek, New mechanisms for double charmed meson production at the LHCb, Phys. Lett. B 758 (2016) 458–464, <https://doi.org/10.1016/j.physletb.2016.05.052>, arXiv:1601.06981.
- [38] A. van Hameren, R. Maciula, A. Szczurek, Single-parton scattering versus double-parton scattering in the production of two $c\bar{c}$ pairs and charmed meson correlations at the LHC, Phys. Rev. D 89 (9) (2014) 094019, <https://doi.org/10.1103/PhysRevD.89.094019>, arXiv:1402.6972.
- [39] M. Diehl, J.R. Gaunt, Double parton scattering theory overview, Adv. Ser. Dir. High Energy Phys. 29 (2018) 7–28, arXiv:1710.04408.
- [40] M. Diehl, J.R. Gaunt, K. Schönwald, Double hard scattering without double counting, J. High Energy Phys. 06 (2017) 083, [https://doi.org/10.1007/JHEP06\(2017\)083](https://doi.org/10.1007/JHEP06(2017)083), arXiv:1702.06486.
- [41] P. Bartalini, J.R. Gaunt, Multiple Parton Interactions at the LHC, Adv. Ser. Dir. High Energy Phys. 29 (2018) 1–450, <https://doi.org/10.1142/10646>.
- [42] J.R. Gaunt, W.J. Stirling, Double parton distributions incorporating perturbative QCD evolution and momentum and quark number sum rules, J. High Energy Phys. 03 (2010) 005, [https://doi.org/10.1007/JHEP03\(2010\)005](https://doi.org/10.1007/JHEP03(2010)005), arXiv:0910.4347.

- [43] M. Aaboud, et al., Study of the hard double-parton scattering contribution to inclusive four-lepton production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Lett. B* 790 (2019) 595–614, <https://doi.org/10.1016/j.physletb.2019.01.062>, arXiv:1811.11094.
- [44] B. Blok, Yu. Dokshitzer, L. Frankfurt, M. Strikman, The four jet production at LHC and Tevatron in QCD, *Phys. Rev. D* 83 (2011) 071501, <https://doi.org/10.1103/PhysRevD.83.071501>, arXiv:1009.2714.
- [45] B. Blok, Yu. Dokshitzer, L. Frankfurt, M. Strikman, pQCD physics of multiparton interactions, *Eur. Phys. J. C* 72 (2012) 1963, <https://doi.org/10.1140/epjc/s10052-012-1963-8>, arXiv:1106.5533.
- [46] B. Blok, Yu. Dokshitzer, L. Frankfurt, M. Strikman, Perturbative QCD correlations in multi-parton collisions, *Eur. Phys. J. C* 74 (2014) 2926, <https://doi.org/10.1140/epjc/s10052-014-2926-z>, arXiv:1306.3763.
- [47] B. Blok, M. Strikman, U.A. Wiedemann, Hard four-jet production in pA collisions, *Eur. Phys. J. C* 73 (6) (2013) 2433, <https://doi.org/10.1140/epjc/s10052-013-2433-7>, arXiv:1210.1477.
- [48] A. Karpishkov, M. Nefedov, V. Saleev, Evidence in favor of Single Parton Scattering mechanism in Υ and D associated production at the LHC, arXiv:1904.05004, 2019.
- [49] P. Bartalini, et al., Multi-Parton Interactions at the LHC, arXiv:1111.0469, 2011.
- [50] M. Strikman, D. Treleani, Measuring double parton distributions in nucleons at proton nucleus colliders, *Phys. Rev. Lett.* 88 (2002) 031801, <https://doi.org/10.1103/PhysRevLett.88.031801>, arXiv:hep-ph/0111468.
- [51] D. d'Enterria, A.M. Snigirev, Same-sign W production in proton-nucleus collisions at the LHC as a signal for double parton scattering, *Phys. Lett. B* 718 (2013) 1395–1400, <https://doi.org/10.1016/j.physletb.2012.12.032>, arXiv:1211.0197.
- [52] D. d'Enterria, A.M. Snigirev, Enhanced J/Ψ -pair production from double parton scatterings in nucleus-nucleus collisions at the Large Hadron Collider, *Phys. Lett. B* 727 (2013) 157–162, <https://doi.org/10.1016/j.physletb.2013.10.004>, arXiv:1301.5845.
- [53] D. d'Enterria, A. Snigirev, Double, triple, and n -parton scatterings in high-energy proton and nuclear collisions, *Adv. Ser. Dir. High Energy Phys.* 29 (2018) 159–187, arXiv:1708.07519.
- [54] E. Huayra, E.G. de Oliveira, R. Pasechnik, Probing double parton scattering via associated open charm and bottom production in ultraperipheral pA collisions, arXiv:1905.03294, 2019.
- [55] B. Blok, M. Strikman, Multiparton pp and pA collisions: from geometry to parton-parton correlations, *Adv. Ser. Dir. High Energy Phys.* 29 (2018) 63–99.
- [56] W.-T. Deng, X.-N. Wang, R. Xu, Hadron production in $p+p$, $p+Pb$, and $Pb+Pb$ collisions with the HIJING 2.0 model at energies available at the CERN Large Hadron Collider, *Phys. Rev. C* 83 (2011) 014915, <https://doi.org/10.1103/PhysRevC.83.014915>, arXiv:1008.1841.
- [57] C. Bierlich, G. Gustafson, L. Lönnblad, H. Shah, The Angantyr model for heavy-ion collisions in PYTHIA8, *J. High Energy Phys.* 10 (2018) 134, [https://doi.org/10.1007/JHEP10\(2018\)134](https://doi.org/10.1007/JHEP10(2018)134), arXiv:1806.10820.
- [58] W. Florkowski, Phenomenology of Ultra-Relativistic Heavy-Ion Collisions, 2010.
- [59] M. Diehl, D. Ostermeier, A. Schafer, Elements of a theory for multiparton interactions in QCD, *J. High Energy Phys.* 03 (2012) 089, [https://doi.org/10.1007/JHEP03\(2012\)089](https://doi.org/10.1007/JHEP03(2012)089), Erratum: *J. High Energy Phys.* 03 (2016) 001, [https://doi.org/10.1007/JHEP03\(2016\)001](https://doi.org/10.1007/JHEP03(2016)001), arXiv:1111.0910.
- [60] D. d'Enterria, A.M. Snigirev, Triple parton scatterings in high-energy proton-proton collisions, *Phys. Rev. Lett.* 118 (12) (2017) 122001, <https://doi.org/10.1103/PhysRevLett.118.122001>, arXiv:1612.05582.
- [61] D. d'Enterria, A.M. Snigirev, Triple-parton scatterings in proton-nucleus collisions at high energies, *Eur. Phys. J. C* 78 (5) (2018) 359, <https://doi.org/10.1140/epjc/s10052-018-5687-2>, arXiv:1612.08112.
- [62] M. Krämer, F.I. Olness, D.E. Soper, Treatment of heavy quarks in deeply inelastic scattering, *Phys. Rev. D* 62 (2000) 096007, <https://doi.org/10.1103/PhysRevD.62.096007>, arXiv:hep-ph/0003035.
- [63] M. Guzzi, P.M. Nadolsky, H.-L. Lai, C.P. Yuan, General-mass treatment for deep inelastic scattering at two-loop accuracy, *Phys. Rev. D* 86 (2012) 053005, <https://doi.org/10.1103/PhysRevD.86.053005>, arXiv:1108.5112.
- [64] T. Binoth, J.P. Guillet, E. Pilon, M. Werlen, A next-to-leading order study of pion pair production and comparison with E706 data, *Eur. Phys. J. C* 24 (2002) 245–260, <https://doi.org/10.1007/s100520200921>, arXiv:hep-ph/0111043.
- [65] A. Majumder, X.-N. Wang, The Dihadron fragmentation function and its evolution, *Phys. Rev. D* 70 (2004) 014007, <https://doi.org/10.1103/PhysRevD.70.014007>, arXiv:hep-ph/0402245.
- [66] T. Kneesch, B.A. Kniehl, G. Kramer, I. Schienbein, Charmed-meson fragmentation functions with finite-mass corrections, *Nucl. Phys. B* 799 (2008) 34–59, <https://doi.org/10.1016/j.nuclphysb.2008.02.015>, arXiv:0712.0481.
- [67] B.A. Kniehl, G. Kramer, Charmed-hadron fragmentation functions from CERN LEP1 revisited, *Phys. Rev. D* 74 (2006) 037502, <https://doi.org/10.1103/PhysRevD.74.037502>, arXiv:hep-ph/0607306.
- [68] R.D. Ball, et al., Parton distributions from high-precision collider data, *Eur. Phys. J. C* 77 (10) (2017) 663, <https://doi.org/10.1140/epjc/s10052-017-5199-5>, arXiv:1706.00428.
- [69] M. Salajegheh, S.M. Moosavi Nejad, M. Soleymaninia, H. Khanpour, S. Atashbar Tehrani, NNLO charmed-meson fragmentation functions and their uncertainties in the presence of meson mass corrections, arXiv:1904.09832, 2019.
- [70] R. Seuster, et al., Charm hadrons from fragmentation and B decays in e^+e^- annihilation at $\sqrt{s} = 10.6$ -GeV, *Phys. Rev. D* 73 (2006) 032002, <https://doi.org/10.1103/PhysRevD.73.032002>, arXiv:hep-ex/0506068.
- [71] G. Alexander, et al., A Study of charm hadron production in $Z^0 \rightarrow c\bar{c}$ and $Z^0 \rightarrow b\bar{b}$ decays at LEP, *Z. Phys. C* 72 (1996) 1–16, <https://doi.org/10.1007/s002880050218>.
- [72] M. Artuso, et al., Charm meson spectra in e^+e^- annihilation at 10.5-GeV c.m.e, *Phys. Rev. D* 70 (2004) 112001, <https://doi.org/10.1103/PhysRevD.70.112001>, arXiv:hep-ex/0402040.
- [73] I. Helenius, H. Paukkunen, Hard probes 2018 conference, <https://indico.cern.ch/event/634426/contributions/3090548/>, (Accessed 17 June 2019).
- [74] P. Skands, S. Carrazza, J. Rojo, Tuning PYTHIA 8.1: the Monash 2013 Tune, *Eur. Phys. J. C* 74 (8) (2014) 3024, <https://doi.org/10.1140/epjc/s10052-014-3024-y>, arXiv:1404.5630.
- [75] A. Buckley, J. Butterworth, L. Lonnblad, D. Grellscheid, H. Hoeth, J. Monk, H. Schulz, F. Siegert, Rivet user manual, *Comput. Phys. Commun.* 184 (2013) 2803–2819, <https://doi.org/10.1016/j.cpc.2013.05.021>, arXiv:1003.0694.
- [76] S. Acharya, et al., Measurement of the production of charm jets tagged with D^0 mesons in pp collisions at $\sqrt{s} = 7$ TeV, arXiv:1905.02510, 2019.
- [77] T. Sjöstrand, M. van Zijl, A multiple interaction model for the event structure in hadron collisions, *Phys. Rev. D* 36 (1987) 2019, <https://doi.org/10.1103/PhysRevD.36.2019>.
- [78] T. Sjöstrand, P.Z. Skands, Multiple interactions and the structure of beam remnants, *J. High Energy Phys.* 03 (2004) 053, <https://doi.org/10.1088/1126-6708/2004/03/053>, arXiv:hep-ph/0402078.
- [79] T. Sjöstrand, The development of MPI modeling in Pythia, *Adv. Ser. Dir. High Energy Phys.* 29 (2018) 191–225, <https://doi.org/10.1142/9789813227767-0010>, arXiv:1706.02166.
- [80] M. Alvioli, M. Azarkin, B. Blok, M. Strikman, Revealing minijet dynamics via centrality dependence of double parton interactions in proton-nucleus collisions, *Eur. Phys. J. C* 79 (6) (2019) 482, <https://doi.org/10.1140/epjc/s10052-019-6998-7>, arXiv:1901.11266.
- [81] K.J. Eskola, P. Paakkinen, H. Paukkunen, C.A. Salgado, EPPS16: nuclear parton distributions with LHC data, *Eur. Phys. J. C* 77 (3) (2017) 163, <https://doi.org/10.1140/epjc/s10052-017-4725-9>, arXiv:1612.05741.
- [82] M. Strikman, W. Vogelsang, Multiple parton interactions and forward double pion production in pp and dA scattering, *Phys. Rev. D* 83 (2011) 034029, <https://doi.org/10.1103/PhysRevD.83.034029>, arXiv:1009.6123.
- [83] T. Lappi, H. Mantysaari, Forward dihadron correlations in deuteron-gold collisions with the Gaussian approximation of JIMWLK, *Nucl. Phys. A* 908 (2013) 51–72, <https://doi.org/10.1016/j.nuclphysa.2013.03.017>, arXiv:1209.2853.
- [84] R. Vogt, Heavy flavor azimuthal correlations in cold nuclear matter, *Phys. Rev. C* 98 (3) (2018) 034907, <https://doi.org/10.1103/PhysRevC.98.034907>, arXiv:1806.01904.
- [85] J.L. Albacete, C. Marquet, Azimuthal correlations of forward di-hadrons in $d+Au$ collisions at RHIC in the Color Glass Condensate, *Phys. Rev. Lett.* 105 (2010) 162301, <https://doi.org/10.1103/PhysRevLett.105.162301>, arXiv:1005.4065.
- [86] J.L. Albacete, C. Marquet, Gluon saturation and initial conditions for relativistic heavy ion collisions, *Prog. Part. Nucl. Phys.* 76 (2014) 1–42, <https://doi.org/10.1016/j.pnpnp.2014.01.004>, arXiv:1401.4866.