

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Pihlainen, Kai; Vaara, Jani; Ojanen, Tommi; Santtila, Matti; Vasankari, Tommi; Tokola, Kari; Kyröläinen, Heikki

Title: Effects of Baseline Fitness and BMI Levels on Changes in Physical Fitness During Military Service

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2020 Elsevier

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Pihlainen, K., Vaara, J., Ojanen, T., Santtila, M., Vasankari, T., Tokola, K., & Kyröläinen, H. (2020). Effects of Baseline Fitness and BMI Levels on Changes in Physical Fitness During Military Service. *Journal of Science and Medicine in Sport*, 23(9), 841-845.
<https://doi.org/10.1016/j.jsams.2020.02.006>

Journal Pre-proof

Effects of Baseline Fitness and BMI Levels on Changes in Physical Fitness During Military Service

Kai Pihlainen, Jani Vaara, Tommi Ojanen, Matti Santtila, Tommi Vasankari, Kari Tokola, Heikki Kyröläinen



PII: S1440-2440(19)31497-5
DOI: <https://doi.org/10.1016/j.jsams.2020.02.006>
Reference: JSAMS 2260

To appear in: *Journal of Science and Medicine in Sport*

Received Date: 16 November 2019
Revised Date: 13 January 2020
Accepted Date: 7 February 2020

Please cite this article as: Pihlainen K, Vaara J, Ojanen T, Santtila M, Vasankari T, Tokola K, Kyröläinen H, Effects of Baseline Fitness and BMI Levels on Changes in Physical Fitness During Military Service, *Journal of Science and Medicine in Sport* (2020), doi: <https://doi.org/10.1016/j.jsams.2020.02.006>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Effects of Baseline Fitness and BMI Levels on Changes in Physical Fitness During Military Service

Kai Pihlainen ¹, Jani Vaara ², Tommi Ojanen ³, Matti Santtila ², Tommi Vasankari ⁴, Kari Tokola ⁴, and Heikki Kyröläinen ^{2,5}

¹ Training Division, Defence Command, Finnish Defence Forces, P.O. Box 919, 00131 Helsinki, Finland

² Department of Military Pedagogy and Leadership, National Defence University, P.O. Box 7, 00861, Helsinki, Finland

³ Finnish Defence Research Agency, Finnish Defence Forces, P.O. Box 5, 04401 Järvenpää, Finland

⁴ UKK Institute for Health Promotion Research, Kaupinpuistonkatu 1, 33500 Tampere, Finland

⁵ Faculty of Sport and Health Sciences, University of Jyväskylä, P.O. Box 35 (VIV), 40014 University of Jyväskylä, Finland

Corresponding author

Kai Pihlainen, MSc, Marunakuja 6 A 28, 00840 Helsinki

Tel. +358 40 024 8910

E-mail. kai.pihlainen@gmail.com

Word count: 2994

Abstract word count: 239

Number of tables: 2 (+2 supplement tables)

Number of figures: 1

ABSTRACT

Objectives. The purpose of the present study was to investigate how aerobic fitness, muscle fitness and body mass index (BMI) change in relation to their baseline levels during 6-12 months of military service.

Design: Retrospective longitudinal follow-up study.

Methods. The study group consisted of 249 279 healthy young male conscripts (age 19.1 ± 0.4 yrs.) who completed their military service between the years 2005-2015. Anthropometrics (body mass, height, BMI), aerobic fitness (12-minute running test) and muscle fitness (sit-ups, push-ups, standing long jump) were measured.

Results. A 12-minute running test improved by 5% (107 ± 292 m), standing long jump 1% (2.1 ± 16.2 cm), 1-min sit-ups 19% (4 ± 8 repetitions/min) and 1-min push-ups 33% (5 ± 10 repetitions/min) ($p < 0.001$ for all). Baseline fitness and baseline BMI levels were inversely associated with their changes ($r = -0.37 - -0.47$, $p < 0.001$). Performance improved in conscripts in the lowest two baseline fitness quartiles in all tests, while it decreased in conscripts in the highest fitness quartiles. In addition, in conscripts who were obese at baseline, body mass decreased on average by 4.9 ± 7.0 kg ($p < 0.001$).

Conclusions. On average, the physical fitness of conscripts improved during their compulsory military service. In particular, conscripts with a lower baseline fitness level or higher BMI showed the largest improvements, which may be significant findings from both a military readiness and national health perspective. However, the decline in physical performance of high-fit conscripts

highlights the importance of individualization of physical training and military training load during military service.

Keywords: Physical performance, exercise, readiness, public health, conscript.

Journal Pre-proof

INTRODUCTION

It is generally recognized that successful management of military duties requires high levels of aerobic fitness and muscular strength¹. Aerobic fitness is particularly important in prolonged military tasks consisting of physical activity at varying intensities². Muscular strength and power are also essential physical performance attributes during physically demanding tasks such as sprinting short distances, jumping over obstacles, and lifting, carrying or dragging heavy loads and materials³. Adequate physical fitness level is also desirable since it may decrease the risk of musculoskeletal injuries and premature discharge from military service^{4,5}. From a health perspective, physical fitness and physical activity have been linked to a number of central and peripheral adaptations that decrease the risk of cardiovascular diseases⁶, which are currently the leading causes of death in males globally⁷. In addition, physical activity is a significant contributor to reductions in excess body weight, which is a risk factor for all-cause mortality and disability⁸.

While some studies have observed positive adaptations to a standardized military training^{9,10,11}, many of them have focused on one training phase such as basic training^{4,12,13}. Currently, there are limited data concerning the effects of longer follow-ups, e.g. 6-12 months, which report physical performance and body composition changes with a large representative sample size and throughout the complete military service time^{14,15}. Finland is one of the very few countries where conscription is compulsory for all healthy males, which results in about 75% of young men performing their military service annually. Currently, the military service is divided into three main phases, known as basic training, special training and unit training period. The length of each period is approximately 8 weeks. In addition, roughly one third of the conscripts are trained for non-commissioned officer and reserve officer tasks for 16 to 22 weeks. During the 6–12 months of military service, conscripts perform progressive physical training that includes combat training and close-order drills, marching and sports-related training for more than 20 hours per week⁹. However,

the execution of optimal military training has become more challenging because an increasing proportion of young Finnish men are overweight and have poor aerobic fitness¹⁶. With this in mind, the purpose of the present study was to investigate military service-induced changes in physical fitness and body mass index (BMI) of male conscripts in relation to their baseline levels.

METHODS

The present retrospective dataset consisted of individual fitness test and body anthropometry results of 249 279 healthy young male conscripts (19.1±0.4 yrs., 179.2±6.6 cm, 77.0±13.3 kg) during 2005–2015 (Table 1). The baseline fitness tests were conducted during the first two weeks of military service by educated fitness instructors, while the post-measurements were performed approximately 4–6 weeks before the end of service. The average duration of the military service was 38 (range 24-52) weeks. After the tests, fitness instructors imported the results to the central database according to the standards determined by the Training Division of the Defence Command. Results concerning temporal changes in the initial body anthropometrics and physical fitness of conscripts from the same dataset were recently published by Santtila et al.¹⁶.

Table 1 here

The present fitness tests and anthropometric measurements were conducted as part of military service, after a physical examination by medical doctors. Safety instructions were given to conscripts before each fitness test, and they were advised of their right to voluntarily interrupt the test at any time. The test termination criteria included the following indications: onset of angina-like symptoms, shortness of breath, wheezing, leg cramps, claudication, light-headedness, confusion, or nausea¹⁷. All subjects were fully informed of the procedures and possible risks associated with the

fitness tests. This study was approved by the Defence Command Finland and conducted according to the 1975 Declaration of Helsinki. The data were anonymized before scientific use.

The measurements of the present study have been described in detail by Santtila et al. ¹⁶. Briefly, anthropometric measurements were conducted by a physician during a standardized medical examination as part of the military service. Body mass was measured to the nearest 100 g and stature with an accuracy of 5 mm. Thereafter, BMI was calculated by dividing body mass (kg) by the square of stature (m²).

Physical fitness tests, protocols and techniques were standardized according to the guidelines of the Defence Command Finland ¹⁷, and the tests were supervised and demonstrated by educated fitness instructors. Aerobic fitness of the conscripts was assessed using the 12-min running test. Conscripts were instructed to perform the test with maximal effort at a progressively increasing running speed and the results were recorded with an accuracy of 10 m.

Separated from the 12-min running test by at least two days, muscle fitness was assessed using standing long jump, 1-min sit-up and 1-min push-up tests. Standing long jump (SLJ) was used to assess explosive power production of the lower extremities. The result of this test was expressed in centimeters from the longest jump of three trials. The sit-up test assessed dynamic performance of abdominal and hip flexor muscles, while the push-up test assessed arm and shoulder extensor muscle performance. The outcome measure of the sit-up and push-up tests was the number of repetitions in one minute.

Commercial software (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp.) was used for statistical analyses. Data are presented as means and standard deviations (\pm SD) where appropriate. Significance of changes between baseline and follow-up measures were analyzed using paired samples T-tests. Pearson correlation coefficients were calculated to examine associations between body composition and physical fitness variables and their changes. The data were additionally pooled into combined BMI categories and quartiles for each fitness test result, and the group differences were tested using analysis of covariance (ANCOVA). In the ANCOVA models, the contributions of categorized baseline fitness and BMI levels to changes in a given fitness variable were examined. Similarly, the contributions of categorized baseline BMI and categorized baseline fitness test results to changes in BMI were determined. Further adjustment for service duration only had a very small effect on comparative values and is therefore not presented in this article. Statistical significance was defined as $p < 0.05$.

RESULTS

The average changes in stature and body mass were 0.5 ± 1.5 cm ($p < 0.001$) and 0.0 ± 4.9 kg ($p = 0.602$), respectively, during the follow-up period. The respective change in BMI was -0.1 ± 1.5 kg/m² ($p < 0.001$). Change in body mass explained 93% of the change in BMI ($R^2 = 0.93$, $p < 0.001$). Despite the modest overall changes, the most marked reductions in body mass were observed in initially obese conscripts. Whereas the underweight (BMI < 18.5) conscripts gained body mass on average by 4.0 ± 4.6 kg ($p < 0.001$), their overweight (BMI 25–29.9) and obese (BMI ≥ 30) counterparts lost -2.0 ± 5.9 ($p < 0.001$) and -4.9 ± 7.0 kg ($p < 0.001$), respectively. Conscripts in the normal BMI category (18.5–24.9) gained body mass by an average of 1.1 ± 3.8 kg ($p < 0.001$).

Mean running distance in the 12-min running test improved by 107 ± 292 m ($p < 0.001$). Respective improvements in SLJ (2.1 ± 16.2 cm), 1-min sit-ups (3.8 ± 8.4 repetitions/min) and 1-min push-ups (5.0 ± 10.1 repetitions/min) also reached statistical significance ($p < 0.001$ for all). In general, one third (34%) of the conscripts improved their results in all fitness tests (12-min run, SLJ, 1-min push-up and sit-up), and another third (34%) improved in three out of four tests. On the other hand, 11% of conscripts performed worse in at least three out of four tests after military service. With the exception of stature, baseline anthropometrics and fitness were moderately associated with their changes ($r = -0.37 - -0.49$, $p < 0.001$) during military service (table 2).

Table 2 here

When the data were pooled into combined baseline fitness quartiles and BMI categories, the largest increases in all fitness variables were observed in conscripts who were initially in the least fit quartiles, irrespective of BMI category (figure 1). Conversely, conscripts who were initially in the highest fitness quartiles showed a decline in performance in all assessed fitness variables, especially in SLJ and the 12-min run.

Figure 1 here

Positive fitness adaptations to military service were more pronounced in the lowest baseline fitness categories and the magnitude of change decreased linearly with increasing baseline fitness, independent of the baseline BMI category (supplement table 3). Baseline fitness had a stronger effect on change in physical fitness than baseline BMI category (supplement table 4). In addition, when compared to the obese BMI category, the most positive adaptations in fitness were observed in conscripts with initially normal BMI.

DISCUSSION

Compulsory military service generally induced beneficial adaptations to physical performance and body composition. A significant proportion of conscripts improved most of the assessed fitness outcomes, and 97.6% of conscripts improved in at least one of the four fitness tests. The most marked improvements were observed in conscripts with the lowest baseline fitness levels. Similarly, the greatest decreases in body mass were observed in conscripts who were obese at the beginning of military service. Hence, compulsory military service led to improvements in the fitness of young men with the greatest need for improvements, while body mass decreased in obese men by nearly 5 kg and their physical fitness improved significantly. On the other hand, conscripts initially in the highest fitness quartile exhibited decreased physical performance in every test, irrespective of baseline BMI. Therefore, military training should be optimized more individually, especially for initially high-fit individuals, to improve their performance during the service.

Although the average changes in body mass were modest overall, conscripts who were initially underweight gained weight and the initially obese conscripts lost weight during military service. While the BMI of underweight conscripts increased by 7%, it declined in obese conscripts by 5%. This is an important finding given that being underweight or overweight is associated with negative health consequences¹⁸. Being of a normal weight is also beneficial from an operative perspective since excess weight increases oxygen cost and may impair performance in critical military tasks¹⁹. These results support previous findings^{14, 15, 20} which have also documented military service-induced decreases in fat mass and changes in fat distribution in obese conscripts, as well as improvements in cardiovascular risk factors in a smaller sample of the same study population. Furthermore, Santtila et al.¹¹ showed that among conscripts with normal BMI, military service increased fat free mass and decreased fat mass without significant changes in body mass.

The improvement of physical fitness, as well as BMI, was presumably due to an increase in total physical activity during the military service and the endurance nature of military training, compared to the habitual physical activity patterns of conscripts before entering the service. Tudor-Locke et al.²¹ reported that over 10 000 steps per day can be classified as a physically active lifestyle. Typically, Finnish conscripts take $13\,937 \pm 2276$ steps per day during military field training²². Furthermore, conscripts often wear combat gear (25-35 kg) and carry other equipment, which increases the total work load of training. The results of the present study support previous findings of improved aerobic capacity and muscular fitness induced by military service^{9, 14, 20}. On the other hand, while conscripts in the lowest baseline fitness quartile showed the largest improvements in test results, conscripts in the highest baseline fitness quartile showed the largest decreases in test performance, irrespective of the baseline BMI category. An inverse relationship between the initial fitness level and its change during the basic training phase of military training has also been found in other studies^{11, 13}. Furthermore, Santtila et al.⁹ showed that most of the positive adaptations occurred during the basic training period, with only small or even no further changes during the following 8 weeks of special training. This suggests that a plateau in training adaptations is reached already during the first eight weeks of military service. Therefore, more variation in training stimulus (intensity and volume) as well as progression of the balance between training load and recovery should be implemented in the later stages of military training in order to induce continuous development of physical fitness throughout military service, especially for conscripts with high initial fitness levels.

More positive adaptations to military training in low-fit subjects and decrements in aerobic fitness of high-fit subjects have also been observed in other countries^{13, 23}. In a 10-month follow-up, Dyrstad et al.²³ found that the aerobic capacity of 107 recruits improved by >1% during the first 10

weeks of military training. However, by the end of military service, the average aerobic capacity had slightly decreased from baseline. It should be noted that the initial mean aerobic capacity was clearly higher in the Norwegian study, while the sample size was not as high as in the present study. Nonetheless, Dyrstad et al.²³ observed an increase in aerobic capacity in the lowest fitness category group. More recently, Burley et al.¹³ compared changes in physical performance of initially low- and high-fit recruits who completed a 12-week basic training regimen. In response to training, initially high-fit recruits showed a greater tendency towards a decrease in aerobic fitness, as well as decreased performance in several military task simulations, whereas most of their low-fit counterparts showed positive changes.

SLJ was used to assess lower-body explosive power production in the present study, and this fitness outcome decreased in the two highest baseline fitness quartiles. Rosendal et al.⁴ found a decrease in unloaded and loaded jump performance after a 12-week basic training period, and concluded that a high volume of endurance activity during military training may have a negative influence on rapid force development of the extensor muscles. It has been shown that high volumes of concurrent strength and endurance training involving the same muscle groups may interfere with the development of explosive power production²⁴. Military training typically consists of a high overall volume of unilateral prolonged, low-intensity endurance-type activities, often with inadequate recovery, which may lead to symptoms of overtraining^{12,25}. Similar findings of baseline fitness-related changes in vertical jump performance have been observed during a 9-month combat operation²⁶. Sharp et al.²⁶ observed an increase in vertical jump performance in initially low-fit soldiers, while jump performance decreased in high-fit soldiers. In this case, the suggested mechanism for the negative adaptation was detraining in higher-fit subjects who were already closer to their maximum potential at the beginning of the deployment. Based on differences in training adaptations of low- and high-fit conscripts in the present study, detraining may be one possible

explanation for the diminished physical performance of high-fit conscripts. On the other hand, excessive monotonous low-intensity endurance-type physical activity, especially for the lower extremities, during military training may induce overtraining symptoms²⁵. These two opposing theories could be tested by using more sophisticated methods such as blood biomarkers or muscle biopsies during longitudinal military training research.

Previous studies have shown significant inverse associations between cardiorespiratory fitness and mortality²⁷. It has also been documented that 6-12 months of military service induces beneficial changes in traditional cardiovascular risk factors, such as blood pressure and cholesterol concentrations¹⁵. Importantly, the observed improvement in aerobic fitness in the present study most likely induced a cardioprotective effect, especially in the obese subjects who improved their 12-min run test result significantly. Furthermore, muscle fitness has been shown to be associated with health outcomes and the association may even be independent of aerobic fitness²⁸. Thus, overall improvements in muscle fitness may offer an additive cardioprotective effect. Collectively, combined improvements in physical fitness and body composition during military service may be considered to induce positive health effects in a large number of young men. This highlights the significant role of military service in public health promotion.

This is the first study to present changes in physical fitness and body anthropometrics during military service in large, nationally representative cohorts of young adult men. Nevertheless, this study also has some limitations. For example, the number of subjects varied for different variables due to missing data. However, our results from an eleven-year period describe rather reliably the average changes in male conscripts during 6-12 months of military training between 2005 and 2015. Covariates such as motivation, diet and smoking, all of which may have affected the changes, were not controlled and may be regarded as limitations of the study. The conscripts lived mainly in the

garrisons, and the same diet (3200-3600 kcal/day) was maintained throughout their structured military service²⁰. Assumedly, diet and energy balance varied between individuals before the military service and therefore, the standardized diet during conscription may also have accounted for changes in body mass as well as fitness variables.

CONCLUSION

Compulsory military service in Finland improves the average aerobic and muscle fitness of conscripts. The positive adaptations are more pronounced in young men who initially have a lower fitness level. Furthermore, obese conscripts and those who are overweight also benefit from military service via its positive effects on body composition. Together, these adaptations may have clinical significance from a public health perspective if they can be maintained to some extent after military service. For the initially high-fit conscripts, increased intensity in endurance and strength training may enhance positive adaptations during military service¹³. One time-efficient solution could be implementation of high-intensity functional training that can be performed at the individual effort level or by using fitness level groups. This concurrent strength and endurance training modality has been shown to induce positive adaptations simultaneously to aerobic and muscular fitness²⁹. Another solution might be dividing conscripts into training groups (inactive, moderate, and active) based on their preservice physical activity as Jurvelin et al.³⁰ recently suggested.

PRACTICAL IMPLICATIONS

- Positive physiological adaptations to military service serve the public health, but also, the aims of the national defense from an operative readiness perspective.
- Some adjustments to physical training such as more progressive training programs and replacement of some of the low-intensity components with higher intensity activities or

strength training are encouraged, especially for conscripts who belong to the highest baseline fitness quartiles.

- These adjustments may help to ensure that the fitness level of the abovementioned subgroups continues to improve or is at least maintained during military service.

ACKNOWLEDGEMENTS

There are no conflicts of interest, financially or otherwise, among any of the authors of this article. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The views, opinions, and/or findings contained in this publication are those of the authors and should not be construed as an official position, policy, or decision of the Finnish Defence Forces, unless so designated by other documentation.

REFERENCES

1. Sharp M, Patton J, Vogel J. A database of physically demanding tasks performed by U.S. army soldiers. Natick: Military Performance Division, U.S. Army Research Institute of Environmental Medicine; 1998. Available at: <http://www.dtic.mil/dtic/tr/fulltext/u2/a338922.pdf>. Accessed 5 November 2019
2. Friedl KE, Knapik JJ, Häkkinen K, et al. Perspectives on aerobic and strength influences on military physical readiness: report of an international military physiology roundtable. *J Strength Cond Res* 2015; 29 Suppl 11: S10-23.
3. Nindl BC, Alvar BA, Dudley J, et al. Executive Summary From the National Strength and Conditioning Association's Second Blue Ribbon Panel on Military Physical Readiness: Military Physical Performance Testing. *J Strength Cond Res*. 2015; 29 Suppl 11: S216-20.
4. Rosendal L, Langberg H, Skov-Jensen A, et al. Incidence of injury and physical performance adaptations during military training. *Clin J Sport Med*. 2003; 13(3): 157-63.
5. Taanila H, Hemminki AJ, Suni JH, et al. Low physical fitness is a strong predictor of health problems among young men: a follow-up study of 1411 male conscripts. *BMC Public Health*. 2011; 25; 11:590.
6. Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. 2009; 20; 301: 2024-35.
7. Naghavi M, Abajobir AA, Abbafati C, et al. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017 Sep 16; 390(10100):1151-1210.
8. Afshin A, Forouzanfar MH, Reitsma MB, et al. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med*. 2017 Jul 6; 377(1): 13-27.

9. Santtila M; Häkkinen K; Nindl BC, et al. Cardiovascular and neuromuscular performance responses induced by 8 weeks of basic training followed by 8 weeks of specialized military training. *J Strength Cond Res* 2012; 26: 745-751.
10. Hofstetter MC, Mäder U, Wyss T. Effects of a 7-week outdoor circuit training program on Swiss Army recruits. *J Strength Cond Res.* 2012; 26(12): 3418-25.
11. Santtila M, Häkkinen K, Karavirta L, et al. Changes in cardiovascular performance during an 8-week military basic training period combined with added endurance or strength training. *Mil Med* 2008; 173: 1173-1179.
12. Booth CK, Probert B, Forbes-Ewan C, et al. Australian army recruits in training display symptoms of overtraining. *Mil Med.* 2006; 171(11): 1059-64.
13. Burley SD, Drain J, Sampson JA, et al. Positive, limited and negative responders: The variability in physical fitness adaptation to basic military training. *J Sci Med Sport* 2018; 21(11): 1168–1172.
14. Mikkola I, Keinänen-Kiukaanniemi S, Jokelainen J, et al. Aerobic performance and body composition changes during military service. *Scand J Prim Health Care.* 2012; 30(2): 95-100.
15. Cederberg H, Mikkola I, Jokelainen J, et al. Exercise during military training improves cardiovascular risk factors in young men. *Atherosclerosis.* 2011; 216: 489-495.
16. Santtila M, Pihlainen K, Koski H, et al. Physical Fitness in Young Men between 1975 and 2015 with a Focus on the Years 2005-2015. *Med Sci Sports Exerc.* 2018; 50(2): 292-298.
17. Pihlainen K, Santtila M, Ohrankämmen O, et al. Fitness Test Manual of the Finnish Defence Forces. Prima Edita. 2011; ISBN 978951220534; 11-12.
18. Flegal KM, Graubard BI, Williamson DF, et al. Excess deaths associated with underweight, overweight, and obesity. *JAMA.* 2005 Apr 20; 293(15): 1861-7.
19. Lyons J, Allsopp A, Bilzon J. Influences of body composition upon the relative metabolic and cardiovascular demands of load-carriage. *Occup Med (Lond).* 2005; 55(5): 380–384.

20. Mikkola I, Jokelainen JJ, Timonen MJ, et al. Physical activity and body composition changes during military service. *Med Sci Sports Exerc.* 2009; 41(9): 1735-42.
21. Tudor-Locke C, Hatano Y, Pangrazi RP, et al. Revisiting “How many steps are enough?” *Med Sci Sports Exerc* 2008; 40 (7 Suppl): S537-43.
22. Ojanen T, Häkkinen K, Vasankari T, et al. Changes in physical performance during 21 days of Military Field Training in Warfighters. *Mil Med* 2018 May 1; 183(5-6):e174-e181.
23. Dyrstad SM, Soltvedt R, Hallén J. Physical fitness and physical training during Norwegian military service. *Mil Med.* 2006; 171(8): 736-41.
24. Häkkinen K, Alen M, Kraemer WJ, et al. Neuromuscular adaptations during concurrent strength and endurance training versus strength training. *Eur J Appl Physiol.* 2003; 89(1): 42-52.
25. Tanskanen M, Uusitalo AL, Häkkinen K, et al. Aerobic fitness, energy balance, and body mass index are associated with training load assessed by activity energy expenditure. *Scand J Med Sci Sports* 2009; 19: 871-878.
26. Sharp M, Knapik J, Walker L et al. Physical fitness and body composition after a 9-month deployment to Afghanistan. *Med Sci Sports Exerc.* 2008; 40(9): 1687–1692.
27. Katzmarzyk P, Church T, Blair S. Cardiorespiratory Fitness Attenuates the Effects of the Metabolic Syndrome on All-Cause and Cardiovascular Disease Mortality in Men. *Arch. Intern. Med.* 2004; 164: 1092–1097.
28. Grøntved A, Ried-Larsen M, Møller NC, et al. Muscle strength in youth and cardiovascular risk in young adulthood (the European Youth Heart Study). *Br J Sports Med.* 2015; 49(2): 90-4.
29. Kyröläinen H, Pihlainen K, Vaara JP, Ojanen T, Santtila M. Optimising training adaptations and performance in military environment. *J Sci Med Sport.* 2018; 21(11): 1131-1138.

30. Jurvelin H, Tanskanen-Tervo M, Kinnunen H, Santtila M, Kyröläinen H. Training Load and Energy Expenditure during Military Basic Training Period. *Med Sci Sports Exerc.* 2020 Jan;52(1):86-93. doi: 10.1249/MSS.0000000000002092.

Journal Pre-proof

TABLES

Table 1. Study sample descriptive statistics at baseline.

Variable	n (pre-post)	mean±SD (pre)
Body mass (kg)	63 366	77+13
Stature (cm)	56 803	179+7
BMI (kg/m ²)	56 444	24.0±3.8
12-min run (m)	218 810	2461±366
Standing long jump (cm)	220 318	218±26
1-min sit-up (reps/min)	221 276	37±11
1-min push-up (reps/min)	220 028	32±14

Table 2. Pearson correlation coefficients between the baseline test results and their changes (Δ) during military service

Variable	<i>n</i>	<i>R</i> (baseline vs. Δ)	<i>p</i>
Body mass	60 102	-0.43	<0.001
Stature	56 679	-0.15	<0.001
Body mass index	56 444	-0.47	<0.001
12-min run	218 810	-0.46	<0.001
Standing long jump	220 318	-0.37	<0.001
1-min sit-up	221 276	-0.44	<0.001
1-min push-up	220 028	-0.39	<0.001

FIGURE LEGENDS

Figure 1. Mean changes in 12-min running distance (a), standing long jump (b), 1-min push-up (c) and 1-min sit-up (d) performance of conscripts in the combined baseline BMI category and fitness quartile groups during military service.

