

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Ikonen, Jenni N.; Joro, Raimo; Uusitalo, Arja L.T.; Kyröläinen, Heikki; Kovanen, Vuokko; Atalay, Mustafa; Tanskanen-Tervo, Minna M.

**Title:** Effects of military training on plasma amino acid concentrations and their associations with overreaching

**Year:** 2020

**Version:** Accepted version (Final draft)

**Copyright:** © Authors, 2020

**Rights:** CC BY-NC-ND 4.0

**Rights url:** <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Please cite the original version:**

Ikonen, J. N., Joro, R., Uusitalo, A. L., Kyröläinen, H., Kovanen, V., Atalay, M., & Tanskanen-Tervo, M. M. (2020). Effects of military training on plasma amino acid concentrations and their associations with overreaching. *Experimental Biology and Medicine*, 245(12), 1029-1038.  
<https://doi.org/10.1177/1535370220923130>

## Effects of military training on plasma amino acid concentrations and their associations with overreaching

Journal:	<i>Experimental Biology and Medicine</i>
Manuscript ID	EBM-19-RM-0645.R1
Manuscript Type:	Original Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Ikonen, Niina; University of Eastern Finland School of Medicine, Institute of Biomedicine Joro, Raimo ; University of Eastern Finland School of Medicine, Institute of Biomedicine Uusitalo, Arja ; University of Helsinki Faculty of Medicine, Clinic for Sports and Exercise Medicine, Kyröläinen, Heikki ; University of Jyväskylä Faculty of Sports and Health Sciences Kovanen, Vuokko ; University of Jyväskylä Faculty of Sports and Health Sciences Atalay, Mustafa; University of Eastern Finland School of Medicine, Physiology Tanskanen-Tervo, Minna ; University of Jyväskylä Faculty of Sports and Health Sciences
Keywords:	overreaching, overtraining, military training, amino acids, glutamine-glutamate ratio, METABOLISM
Abstract:	<p>Amino acids are thought to have a key role in the processes contributing to overreaching development through their metabolic properties and neuronal functions. In the present study, the effects of 10-week military training on the concentrations of 19 amino acids were investigated. Plasma amino acid concentrations were measured at rest from 53 healthy male conscripts on weeks 1, 4, 7 and 9 of their military service. Conscripts were classified as overreached and non-overreached. Overreaching classification was based on fulfilling at least 3 of 5 criteria: greater than 5 % decrease in maximal oxygen uptake, increased Rating Perceived Exertion (RPE) and decreased lactate-RPE ratio in submaximal marching test, admitting feeling overloaded and both increased scores in fatigue and decreased scores in vigor in the Profile of Mood States Adolescents. Eight conscripts (18 %) were classified as overreached; their glutamine-glutamate ratio and alanine and arginine levels were significantly lower (<math>p &lt; 0.05</math>) and glutamate concentration significantly higher (<math>p &lt; 0.05</math>) in comparison to their non-overreached counterparts. The levels of arginine increased (<math>p &lt; 0.05</math>) and tryptophan (<math>p &lt; 0.001</math>) decreased in both groups throughout the study. The tyrosine concentration increased in non-overreached but, in contrast, remained at the same level in overreached individuals (<math>p &lt; 0.05</math>). The results suggest that alterations in the levels of three metabolically important amino acids, alanine, glutamate and arginine, and the possibly neuroactive tyrosine and tryptophan might explain some of the physical and psychological symptoms of overreaching. The present study also confirms the potential use of glutamine-glutamate ratio as a tool for detecting overreaching.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



SCHOLARONE™  
Manuscripts

# Effects of military training on plasma amino acid concentrations and their associations with overreaching

Niina J. J. Ikonen<sup>1</sup>, Raimo I. A. Joro<sup>1</sup>, Arja L. Uusitalo<sup>2a,b</sup>, Heikki Kyröläinen<sup>3</sup>,  
Vuokko Kovanen<sup>3</sup>, Mustafa Atalay<sup>1\*,\*\*</sup>, Minna M. Tanskanen-Tervo<sup>3\*\*</sup>

<sup>1</sup>Institute of Biomedicine, University of Eastern Finland, PL 1627, 70211 Kuopio, Finland

<sup>2a</sup>Department of Sports and Exercise Medicine, Clinicum, University of Helsinki, Alppikatu 2, 00530 Helsinki, Finland

<sup>2b</sup>Clinic for Sports and Exercise Medicine, Foundation for Sports and Exercise Medicine, Alppikatu 2, 00530 Helsinki, Finland

<sup>3</sup>Faculty of Sport and Health Sciences, University of Jyväskylä, PL 35, 40014 Jyväskylä, Finland

\*Corresponding author: Mustafa Atalay. Email: mustafa.atalay@uef.fi. Institute of Biomedicine, University of Eastern Finland, PL 1627, 70211 Kuopio, Finland

\*\* Minna M. Tanskanen-Tervo and Mustafa Atalay contributed equally to this work and are shared last authors

**Running Title:** Military training and plasma amino acids concentrations

## Abstract

Amino acids are thought to have a key role in the processes contributing to overreaching development through their metabolic properties and neuronal functions. In the present study, the effects of 10-week military training on the concentrations of 19 amino acids were investigated. Plasma amino acid concentrations were measured at rest from 53 healthy male conscripts on weeks 1, 4, 7 and 9 of their military service. Conscripts were classified as overreached and non-overreached. Overreaching classification was based on fulfilling at least 3 of 5 criteria: greater than 5 % decrease in maximal oxygen uptake, increased Rating Perceived Exertion (RPE) and decreased lactate-RPE ratio in submaximal marching test, admitting feeling overloaded and both increased scores in fatigue and decreased scores in vigor in the Profile of Mood States Adolescents. Eight conscripts (15 %) were classified as overreached; their glutamine-glutamate ratio and alanine and arginine levels were significantly lower ( $p < 0.05$ ) and glutamate concentration significantly higher ( $p < 0.05$ ) in comparison to their non-overreached counterparts. The levels of arginine increased ( $p < 0.05$ ) and tryptophan ( $p < 0.001$ ) decreased in both groups throughout the study. The tyrosine concentration increased in non-overreached but, in contrast, remained at the same level in overreached individuals ( $p < 0.05$ ). The results suggest that alterations in the levels of three metabolically important amino acids, **alanine, glutamate and arginine**, and the possibly neuroactive tyrosine and tryptophan might explain some of the physical and psychological symptoms of overreaching. The present study also confirms the potential use of glutamine-glutamate ratio as a tool for detecting overreaching.

1  
2  
3 **Keywords:** overreaching; overtraining; military training; amino acids; glutamine-  
4  
5 glutamate ratio; metabolism  
6  
7

## 8 9 10 **Impact statement**

11  
12 The diagnosis of overtraining syndrome and overreaching poses a great challenge.  
13  
14 Military training aims at improving the physical performance of the conscripts, but an  
15  
16 excessive training load could also lead to overreaching. This study of Finnish conscripts  
17  
18 provides new insights into the pathophysiology of overreaching and overtraining through  
19  
20 amino acids concentrations. In addition to confirming the possible use of the plasma  
21  
22 glutamine/glutamate concentration to indicate and predict overreaching, we made a novel  
23  
24 finding i.e. low alanine and arginine concentrations might have a role in the performance  
25  
26 decrement and fatigue related to overreaching. Moreover, this study is the first to show  
27  
28 the possible association between amino acids with putative neuronal properties and  
29  
30 overreaching. Thus, the present findings might help to detect and prevent overreaching  
31  
32 and offer a reliable diagnostic approach in the future. In order to avoid overreaching,  
33  
34 military training should be planned more periodically and individually, especially during  
35  
36 the first four weeks of military service.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

A primary target of basic military training (BT) programs is to develop physical fitness, strength and endurance of the conscripts within a short time to enable them to perform combat-specific tasks. Imbalanced excessive training and recovery, however, can lead to overreaching (OR), a short-term decrement of performance, which may take several days or even weeks for recovery<sup>1</sup>. A prolonged imbalance, in turn, may lead to the so-called overtraining syndrome (OTS), a long-term decrease in performance despite continuous training, from which recovery may take months or years<sup>1</sup>.

Despite intensive research, the mechanism underlying the development of OTS and OR has not yet been clarified nor have any reliable and specific biomarkers been identified. However, both OTS and OR may also include several individually varying physiological and psychological symptoms<sup>1</sup>. Amino acids are involved in many crucial biochemical processes i.e. brain function and various metabolic pathways including protein turnover and energy metabolism during exercise. Therefore, they could be associated with the symptoms of OTS and OR. The glutamine-glutamate (Gln/Glu) ratio has been reported to decline in overreached athletes<sup>2-4</sup> whereas increased glutamate (Glu) levels in OR may reflect excessive training stress while a decreased glutamine (Gln) level has been associated with a deteriorated capacity of work<sup>5</sup> and an impaired immune system<sup>3, 4, 6</sup>. Nevertheless, not all studies have reported altered Glu<sup>3, 4</sup> and Gln concentrations in overreached athletes<sup>3, 4, 6</sup> nor found a link between Gln and immune depression<sup>7, 8</sup>. On the other hand, the branched chain amino acids (BCAAs) leucine, isoleucine and valine are known to compete with tryptophan for the same carrier in the blood-brain barrier and

1  
2  
3 furthermore, these energy yielding BCAAs are utilized when exercising. Hence, a lower  
4 BCAA concentration has been suggested to increase tryptophan intake to the brain during  
5 exercise. Tryptophan, is a precursor of the neurotransmitter serotonin, and therefore, its  
6 excess could be involved in the development of centrally-mediated fatigue related to OR<sup>9</sup>,  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

<sup>10</sup>. Thus, strenuous training evoked alterations in amino acid metabolism may trigger the development of OR.

There is a paucity of knowledge on whether changes in the levels of amino acids can be involved in OR. Military training provides a relevant model to study OR and OTS because the conscripts represent a healthy homogenous age group undergoing strenuous training programs while having a controlled diet and an active lifestyle. Moreover, intolerance of conscripts to BT represents a major challenge to military training and therefore the mechanisms underlying this condition need to be clarified. Therefore, the aim of this study was to investigate the effect of military training on the concentrations of 19 plasma amino acid in overreached (OR) and non-overreached (nOR) conscripts during the BT period and to evaluate the role of amino acids as a tool for indicating OR.

## Materials and Methods

### Subjects

Finnish male conscripts (N=53, age  $19.6 \pm 0.3$  years) volunteered to take part in the study. They were a part of 60 participants who had been selected according to specific criteria including willingness and not having any musculoskeletal and cardiorespiratory



1  
2  
3 disorders<sup>11</sup>. All the conscripts were informed of the study design and experiments, gave  
4 their written informed consent and were given permission to withdraw from the study at  
5 any point. The study was approved by the Finnish Defence Forces and the Ethical  
6 Committees of the University of Jyväskylä and the Kainuu region of Finland. This study  
7 was funded by the Finnish Ministry of Education, Finnish Cultural Foundation, Polar  
8 Electro Oy and the Scientific Advisory Board for Defence.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

## 19 **Military training protocol**

20  
21  
22 Finnish conscripts performed a standard 8-week BT period and a consequent 2 weeks of  
23 their specialized military training in wintertime where the outdoor temperatures varied  
24 between -30 and +1 °C. Military service included gradually increasing physical exercise  
25 from 2 h per day to approximately 4 h at the end of the study consisting of both sports-  
26 related endurance and strength type of physical training and military-related physical  
27 training such as combat training and other forms of military training including shooting,  
28 and standard military education<sup>11</sup>. Conscripts marched to the dining rooms four times a  
29 day (approximately 5 km in total) to consume standard military meals and **performed**  
30 **other daily living activities as described previously<sup>11</sup>. The dietary intake and energy**  
31 **balance have been assessed in a previous study, where the reported intake indicated that**  
32 **daily carbohydrate consumption in relation to the total reported energy intake was 58 ±**  
33 **4% (5.2 ± 1.6 g·kg<sup>-1</sup>), fat 27 ± 4% (1.1 ± 0.4 g·kg<sup>-1</sup>), protein 14 ± 2 % (1.3 ± 0.4 g·kg<sup>-1</sup>),**  
34 **and alcohol 0.3 ± 0.9% (0.01 ± 0.04 g·kg<sup>-1</sup>)<sup>12</sup>. During the study period, there were 4**  
35 **marches with battle equipment and an overnight combat training exercise.**  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Study protocol

To measure the symptoms of overtraining, conscripts performed physical and psychological tests, which have been utilized in previous studies investigating overtraining or training adaptation among endurance and strength athletes and soldiers, since military training is a combination of various stressors<sup>11</sup>. These tests involved performance tests such as a submaximal marching test and a maximal treadmill test, Profile of Mood States Adolescents (POMS-A) questionnaire<sup>13</sup>, an inquiry of upper respiratory infections (URTI), and direct question on fatigue and assessment of Rating Perceived Exertion (RPE). Blood samples were also drawn and analyzed. The overall study protocol is presented in Table 1.

Conscripts performed a 45-minute-submaximal marching test with battle equipment on an outdoor track at the speed of their 70 % maximal workload to determine their general aerobic performance in a military environment. This was calculated by using the American College of Sports Medicine estimation formula for maximal oxygen uptake ( $VO_{2max}$ ) in running and taking into account the 20-kg combat gear<sup>14</sup>. The marching test was chosen since it represented the daily routine well and was available for a large group of soldiers. RPE was assessed 3 times, every 15 minutes using a 6-to 20-point scale, and their averaged value was utilized in the further analyses. The maximal treadmill test for evaluating  $VO_{2max}$ , in turn, included a specific warming up, walking and running at pre-determined velocities, and a gradually increasing speed and inclination of the treadmill until the point of exhaustion<sup>14</sup>. RPE was measured just before each increase in the

1  
2  
3 workload. Blood lactate was determined 1 min after the completion of the exercise from  
4 fingertip blood using a lactate analyzer (LactatePro, Arkray, Japan). Lactate-RPE ratio  
5  
6 being determined subsequently. Pulmonary ventilation and respiratory gas exchange data  
7  
8 were measured by a breath-by-breath method (Jaeger Oxygen Pro, VIASYS Healthcare  
9  
10 GmbH, Hoechberg, Germany). Heart rate was recorded at 5-s intervals (Polar810i, Polar  
11  
12 Electro Oy, Kempele, Finland). The maximal treadmill tests and submaximal tests were  
13  
14 performed four times in total (Table 1). The exercise protocol remained the same during  
15  
16 the entire study.  
17  
18  
19  
20  
21  
22  
23

### 24 **Physical activity**

25  
26 The subjects reported their physical activity during one month before military service by  
27  
28 a self-administered short version of international physical activity questionnaire (IPAQ)  
29  
30 covering the previous seven days<sup>15</sup>. IPAQ outcome and physical activity were reported as  
31  
32 minutes per week calculated in five different categories 1) Light PA: daily minutes x days  
33  
34 per week with walking x 3.3 MET 2) Moderate PA: daily minutes x days per week with  
35  
36 moderate-intensity physical activity x 4.0 MET 3) Vigorous PA: daily minutes x days per  
37  
38 week with physical, vigorous activity x 8.0 MET 4) Moderate to vigorous PA (ModVig  
39  
40 PA): Moderate PA + Vigorous PA. 5) Total PA: Light PA + Moderate PA + Vigorous  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
PA.

### 51 **POMS-A questionnaire**

52  
53 On weeks 1, 2, 4, 7, 8 and 9, the conscripts filled in a POMS-A questionnaire for  
54  
55 evaluating their mood state by answering questions about it over the past week as well as  
56  
57  
58  
59  
60

1  
2  
3 the current day by using a scale from 1 to 5, (1 refers to not at all and 5 extremely). Points  
4  
5 were summed up and mood scores determined for each conscript. Conscripts were also  
6  
7 directly asked “Do you feel mentally or physically overloaded?” on weeks 7, 8 and 9.  
8  
9

### 10 11 12 **Inquiry about URTIs and sickness-related absences**

13  
14 Conscripts filled in a form enquiring about the symptoms of URTIs at baseline and on  
15  
16 weeks 4, 6, 7, 8 and 9. Their responses were rated from 0 to 5 and a maximum sum of  
17  
18 points was 55 on each survey week. Points were added up, and cumulative sums for  
19  
20 weeks 2, 4, 7, 8 and 9 were created to represent the total burden of URTIs per week.  
21  
22 Sickness-related absences were defined as an attendance / not-attendance in service,  
23  
24 because of illnesses or injuries examined by physician, and as previously reported<sup>11</sup>, the  
25  
26 average number of sickness absences was 3 days; the main reason for sickness absence  
27  
28 was URTI (71 %) while musculoskeletal disorders (13 %) and other disorders (13 %)  
29  
30 were the second most common reasons with digestion disorders accounting for the  
31  
32 remainder of the sick leaves. In this study, sickness absence days were added up and  
33  
34 divided into tertiles (from 0 to 2 days, from 3 to 6 days, and from 7 to 18 days).  
35  
36  
37  
38  
39  
40  
41

### 42 **Anthropometric measurements and fitness level**

43  
44 Body composition measurements of body mass (BM), fat free mass (FFM), fat mass  
45  
46 (FM) and percentage of body fat (F%) were determined using an eight-point bioelectrical  
47  
48 impedance (Inbody 720, Biospace Co. Ltd, Seoul, Korea). For each subject, the  
49  
50 measurements were performed between 6 a.m. and 7 a.m. after an overnight fast and after  
51  
52 voiding, with no exercise for 12 hours before the test. Conscripts wore T-shirts and  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 trousers during the measurements. Height was measured by using a wall-mounted  
4  
5 stadiometer and rounded to the nearest 0.5 cm.  
6  
7  
8  
9

## 10 **Blood samples**

11  
12 Blood samples were drawn from an antecubital vein after an overnight fast (10 hours)  
13  
14 each time in the seated position, between 6:30 and 7:30 a.m. The first sample was taken  
15  
16 on day 7, followed by measurements on weeks 4, 7 and 9. Fingertip lactate samples were  
17  
18 taken before and after the submaximal exercise between 9 and 12 a.m. after standard food  
19  
20 ingestion as described in the previous study<sup>11</sup>. After protein precipitation, the  
21  
22 concentrations of free amino acids in plasma were determined by reversed phase high  
23  
24 performance liquid chromatography (RPHPLC). The HPLC system included a  
25  
26 Quaternary Gradient Pump unit, PU-2089 Plus, an Intelligent Autosampler AS-2057 Plus,  
27  
28 Intelligent Fluorescence Detector, FP-2020 by Jasco, and data processing software Jasco  
29  
30 Chrompass as previously described. Briefly, plasma samples were extracted with  
31  
32 acetonitrile and derivatized to form fluorescent o-phthalaldehyde conjugates. Individual  
33  
34 amino acids were separated on a reversed phase Zorbax C18 column (3.0 mm x 150 mm  
35  
36 x 3.5  $\mu$ m, Agilent Technologies, Santa Clara, CA) and detected at wavelengths 338 nm  
37  
38 for excitation and 455 nm for emission<sup>16</sup>.  
39  
40  
41  
42  
43  
44  
45  
46

## 47 **Overreaching classification**

48  
49 Conscripts were classified as overreached (OR) and non-overreached (nOR) according to  
50  
51 five criteria from which they had to fulfill at least three in order to be classified as OR to  
52  
53 ensure the most accurate classification. Since there are no exact diagnostic tools to  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 identify either OR or OTS and the symptoms may vary among individuals, the OR  
4  
5 criteria were chosen based on the literature regarding accepted physiological changes in  
6  
7 OR athletes or soldiers. According to the literature, the most typical symptoms related to  
8  
9 OR are performance decline with a reported increase in perceived exertion<sup>17-20</sup>, fatigue<sup>1,</sup>  
10  
11 <sup>19, 21</sup> and mood changes<sup>1, 22, 23</sup> and these signs were taken into consideration in the  
12  
13 classification. Table 2 presents the descriptive data between the nOR and OR conscripts  
14  
15 on week 1 of the present study.  
16  
17  
18  
19  
20

21  
22 Criterion 1 was a decreased  $VO_{2max}$  ranging from greater than 5 %<sup>19</sup> of the lowest value  
23  
24 on week 1 or 5 to the lowest value on week 7 or 9. Criterion 2 was an increase greater  
25  
26 than 1.0 in mean RPE during the sub-maximal marching test<sup>17-19</sup> from the lowest value on  
27  
28 week 1 or 4 until weeks 7 and 9. Criterion 3 was a decrease greater than 0.27 mmol in the  
29  
30 lactate-RPE ratio. Two times the standard deviation was set as a threshold since it was  
31  
32 clearly different compared to the values from healthy athletes, and the literature  
33  
34 illustrates similar changes in OR athletes<sup>24</sup>. Criterion 4 was feeling mentally or physically  
35  
36 overloaded on weeks 7, 8 and 9. Criterion 5 was a decrease greater than 1.0 in vigor in  
37  
38 POMS-A from the highest score on weeks 1, 2 or 4 until weeks 7, 8 and 9 and an increase  
39  
40 greater than 1.0 in fatigue from the lowest score on week 1, 2 or 4 until weeks 7, 8 and 9.  
41  
42 Both criteria had to be valid and the change had to remain during the following weeks  
43  
44 until the end of the 10-week study period. In order to avoid the possible impact of  
45  
46 individual mental stress caused by the change to a military environment, the baseline  
47  
48 value was taken from the lowest value in the early weeks. Only vigor and fatigue were  
49  
50 considered in our study since they have been reported to clearly change in overreached  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 conditions while the other negative moods have increased or presented no change<sup>4, 19, 25,</sup>  
4  
5 <sup>26</sup>. We did not take into account URTIs as a criterion of OR since the military  
6  
7 environment poses a high risk for even otherwise healthy adults to develop respiratory  
8  
9 illnesses <sup>27, 28</sup>, and possible epidemics in wintertime could lead to an overestimation of  
10  
11 OR among conscripts.  
12  
13  
14  
15  
16

## 17 **Statistical analysis**

18  
19 Statistical analyses were performed using IBM Statistics SPSS 20. To determine the  
20  
21 possible differences in the descriptive statistics on week 1 between the OR and nOR, an  
22  
23 independent sample T-test was performed if normality was evident. Total physical  
24  
25 activity, height, fat mass and Body Mass Index (BMI) were not normally distributed and,  
26  
27 therefore, Mann-Whitney U test was used. Pearson's  $\chi^2$  was performed to evaluate  
28  
29 possible differences between the groups in the fitness level and in the three-point scale of  
30  
31 sickness absences. Possible differences in the scores of the URTI inquiry between nOR  
32  
33 and OR were tested by Mann-Whitney U test as data was skewed. Generalized linear  
34  
35 mixed model was used to identify significant differences, for the effect of training  
36  
37 between the groups and training x group interactions. The least significance difference  
38  
39 (LSD) post hoc test was used to examine the differences between the groups on a specific  
40  
41 week if a main effect of group on any specific amino acid was detected. In order to fulfill  
42  
43 the assumption of normally distributed residuals, glutamate was log transformed. The  
44  
45 estimates of the mixed model (E) are reported. Significance was set at  $p < 0.05$ .  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Results

Out of a total of 53 conscripts, 8 (15 %) were classified as OR with the remaining 45 designated as nOR. Criterion 1 was met by 47 % of the conscripts, while criteria 2, 3, 4, 5 were met by 16 %, 20 %, 49 % and 18 %, respectively. There were no statistical differences in anthropometric measurements nor in the level of activity between nOR and OR at the beginning of the study (Table 2). The response rate for URTI inquiry was 100 %, 96%, 87 %, 68 % and 51 %, on weeks 2, 4, 7, 8 and 9, respectively. All conscripts reported URTI symptoms during BT. There was no significant difference in the symptoms of URTI between the groups apart from week 8 when the OR individuals had a significantly higher cumulative score in the URTI inquiry ( $p<0.05$ ). Sickness absence data was available for 98 % of the conscripts; 38 % of the conscripts had sickness absence of 0 to 2 days, 34 % of 3 to 6 days and 26 % of 7 to 18 days. When compared to nOR, OR conscripts also had significantly more sickness absence days in the subscale 7-18 days ( $p<0.05$ ). There was a main effect of group for Gln/Glu ratio, and levels of Gln, alanine and arginine ( $p<0.05$ ) (Figure 1). Compared to nOR, Gln/Glu, and the levels of alanine and arginine were significantly lower in OR ( $E -10.1\pm 4.3$ ,  $-119.1\pm 49.1$ ,  $-65.0\pm 25.0$ , respectively) during the entire 10-week study period ( $p<0.05$ ) (Figure 1). Moreover, the Glu concentration was higher ( $E 0.2\pm 0.1$ ,  $p<0.05$ ) in OR than in nOR during the first four weeks of military training (Figure 1). The arginine level also increased in both groups ( $E 4.6\pm 2.5$ ) ( $p<0.05$ ). There were no significant changes in the amounts of Gln or BCAAs among nOR and OR individuals over the study period. The tyrosine concentration exhibited a group x training interaction with no change in OR but an increase in nOR ( $E 1.4\pm 0.6$ ,  $p<0.05$ ) on weeks 4, 7 and 9 compared to the



1  
2  
3 measurement at day 7. Tryptophan levels decreased ( $E -1.2 \pm 0.4$ ,  $p < 0.001$ ) throughout the  
4  
5 study period in both groups. Otherwise, there were no significant changes in the  
6  
7 concentrations of the other amino acids (Table 3).  
8  
9  
10  
11  
12  
13  
14  
15

## 16 Discussion

17  
18  
19 As far as we are aware, the present study is the first to have evaluated the role of 19  
20 amino acids in overreaching. The present study provides further evidence that the  
21  
22 Gln/Glu ratio is a reliable tool for evaluating overreaching, mainly due to significant  
23  
24 increase in Glu in OR. One notable new finding was the result that alanine, arginine and  
25  
26 tyrosine might be involved in overreaching. In addition, military training, which includes  
27  
28 not only endurance and strength-type of training but also other stressors like sleep  
29  
30 deprivation, negative energy balance, as well as stressful environmental factors like  
31  
32 working, living and sleeping outdoors and, occasionally, a restricted social life, seems to  
33  
34 have an impact on tryptophan and arginine concentrations.  
35  
36  
37  
38  
39  
40  
41  
42

43 As there are no specific diagnostic criteria for OR and the existing classifications have  
44  
45 not been clinically validated, we created a 5-scale categorization based on the literature  
46  
47 regarding typical findings on overreaching and overtraining. It is, however, recognized  
48  
49 that several markers may indicate OR<sup>1</sup>. Therefore, in an attempt to ensure the most  
50  
51 accurate classification, we required that at least three of five criteria had to be valid in  
52  
53 order for an individual to be classified as OR. The selected criteria values were clearly  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 outside the normal range and represented the following changes in OR. Criterion 1,  
4 decrease in  $VO_{2max}$ , which is the most important and most widely applied parameter as a  
5 critical criterion of overtraining, and criterion 4, feelings of being mentally or physically  
6 overloaded, were fulfilled by almost every second conscript while the rest of the fulfilled  
7 criteria represented approximately a prevalence of OR (15 %) among conscripts. Since all  
8 conscripts had reported symptoms of URTI, the observed decrease in  $VO_{2max}$  in nOR  
9 might be an effect of mild or recent URTI. As military training consists of specified  
10 training programs combined with environmental stressors, conscripts are under pressure  
11 to learn new combat training skills and feelings of fatigue are subjective, almost half of  
12 the conscripts might have felt themselves mentally or physically overloaded, although  
13 classical OR might not have been present.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 Interestingly, we also observed differences in the levels of alanine, in the Gln/Glu ratio,  
32 as well as in arginine and Glu levels between the groups on the 7<sup>th</sup> day of military  
33 training when OR was expected to be present. However, the OR conscripts had reported  
34 higher but non-significant, physical activity levels compared to nOR before starting their  
35 military service. Although the  $VO_{2max}$  level slightly differed between groups, an elevated  
36 physical activity level (even if statistically non-significantly) might be a predisposing  
37 factor for developing OR later during BT and might have an impact on the results  
38 measured during the 1<sup>st</sup> week. Alterations in amino acid profiles may represent early  
39 biochemical changes related to the onset of OR, although OR was not yet clinically  
40 evident. On the other hand, an altered amino acid profile might further aggravate OR  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 through the impacts of these changes in amino acid levels on several metabolic, neuronal  
4 and hormonal pathways which will be discussed in the following paragraphs.  
5  
6  
7  
8  
9

10 Previously, several studies have suggested that a lower Gln/Glu ratio may be a promising  
11 marker for overreaching due to either lower Gln or higher Glu<sup>2-4</sup>. Smith and Norris<sup>5</sup> even  
12 proposed overreaching to occur when Gln/Glu would be less than 3.58 but higher ratios  
13 were observed in the present study. The differences might be due to the different  
14 analytical methods, since in our study, amino acids were analyzed by HPLC while a  
15 colorimetric assay was used in the study of Smith and Norris<sup>5</sup>. Nevertheless, also in the  
16 present study, the Gln/Glu ratio was significantly reduced in OR thus confirming the  
17 results from previous studies that the Gln/Glu ratio may be beneficial for evaluating OR.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 In some studies, the Gln concentration has been observed to decrease in overreached  
32 athletes<sup>8, 29, 30</sup> and this has been proposed to reflect a response to the training volume<sup>5</sup>,  
33 representing a negative effect of exercise stress<sup>29</sup>, which may also result in overload  
34 training-induced impairment of the immune system. Monocytes and lymphocytes require  
35 Gln for their activity and, therefore a lower Gln concentration could explain the more  
36 common URTIs in OR<sup>6, 31, 32</sup>. We also detected that the OR conscripts had more  
37 symptoms of URTI by week 8 than their nOR counterparts. The difference disappeared  
38 on the 9<sup>th</sup> week; however, the response rate also declined which may have affected the  
39 results. Nevertheless, we did not observe any difference of Gln levels between OR and  
40 nOR which is in agreement with other studies<sup>3, 4</sup>. Therefore, we did not find any  
41 association of low glutamine levels with URTIs in OR. However, URTIs may have  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 contributed to the development of OR and exposed the OR conscripts to a higher  
4  
5 likelihood of sickness absence compared with nOR.  
6  
7  
8  
9

10 Consistent with our results, earlier studies have reported a significant increase in Glu  
11 levels in OR humans<sup>4,5</sup> and rats<sup>33</sup> due to excessive training stress. Therefore, as there was  
12 no difference in Gln levels between nOR and OR, this indicates that the lower Gln/Glu  
13 ratio in OR is probably due to increased Glu levels although marginal and insignificant  
14 changes in Gln may alter this ratio. The increase in Glu concentration may precede OTS  
15 rather than being specific for OTS since in the present study there was no difference  
16 between OR and nOR in the post hoc tests on the week 7 and 9. In support of this  
17 proposal, Coutts et al.<sup>3</sup> reported that Glu levels tended to increase in response to intensive  
18 training but did not detect a significant difference between OR and nOR. In addition, our  
19 findings agree with Smith and Norris<sup>5</sup> suggesting that the Glu level could potentially be  
20 used to evaluate an individual's tolerance to training and risk of OR. Thus, our study  
21 indicates that the ability to cope with training load was mostly impaired in the first four  
22 weeks of basic military training period.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 There is little in the current literature regarding the effect of training on the plasma  
43 arginine concentrations. However, exercise stimulates the production of nitric oxide<sup>34</sup>,  
44 while arginine has an important role as its precursor<sup>35,36</sup>. A recent study demonstrated  
45 that aerobic exercise training decreased arginine concentrations but increased nitric oxide  
46 generation in a heterogeneous population of subjects<sup>37</sup>. In contrast to this observation, we  
47 found that military training increased arginine levels in both the OR and nOR groups.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Thus, the possible underlying mechanism may not be related solely to the nitric oxide  
4 metabolism. The conflicting results may also be due to the characteristics of the subjects  
5 and the different type and intensity of the training program as military training is a  
6 combination of strength and endurance training, which might differentially affect the  
7 arginine levels. Moreover, arginine also has multiple functions in the body and an  
8 increased arginine concentration may be associated with several exercise adaptations  
9 including the synthesis of creatine<sup>38</sup>, enhanced ammonia removal during exercising<sup>39</sup> and  
10 increased growth hormone secretion<sup>40</sup> and release<sup>41</sup>.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Furthermore, we detected a difference in the arginine concentrations between nOR and  
25 OR. Arginine is a conditionally essential amino acid and under stressful conditions,  
26 including serious illnesses and intensive physical activity, there is elevated utilization of  
27 arginine for physiological processes<sup>42, 43</sup>. Thus, the increased utilization rate of arginine  
28 might be a possible response to stressful conditions and this could account for the  
29 difference in OR, even though a training response was detected in both groups. Arginine  
30 metabolism might be involved in the development of OR and contribute to some of the  
31 symptoms via metabolic pathways.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 Interestingly, we found a reduced alanine concentration in OR. Alanine has a central role  
46 in energy metabolism during exercise through its role in the glucose-alanine cycle, which  
47 is a crucial step of gluconeogenesis in the liver<sup>44</sup>. In support of this mechanism, previous  
48 studies have reported that the plasma alanine concentration increases in response to acute  
49 exercise<sup>44-46</sup> but decreases if it is prolonged<sup>47</sup>. Nevertheless, since alanine is closely  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 related to energy metabolism and we found a decreased alanine concentration in OR, it is  
4 possible to postulate that OR is linked with an increased gluconeogenic demand.  
5  
6

7 Furthermore, there is a consensus that OR represents a maladaptation of the  
8 hypothalamic-pituitary-adrenal axis (HPA)<sup>1</sup> indicating that there might be a  
9  
10 hypersensitivity of the pituitary subsequently followed by insensitivity and a blunted  
11  
12 hormone response<sup>48, 49</sup>. Thus, in the early stages of OR, there might be elevated cortisol  
13  
14 and catecholamine responses as a result of abnormalities in the HPA axis, leading to  
15  
16 increased gluconeogenesis which can also be connected with the lower alanine levels. In  
17  
18 addition, we previously reported a non-significant negative energy balance among the  
19  
20 present conscripts<sup>12</sup>, which can be a consequence of strenuous military training with  
21  
22 high-energy expenditure and the lack of sleep and nutritional deprivation. Energy  
23  
24 deficiency is a known risk factor for OR<sup>1</sup> and might be associated with alterations in  
25  
26 alanine levels.  
27  
28  
29  
30  
31  
32  
33  
34

35 According the central fatigue theory, the changes in tryptophan, which is a precursor of  
36 serotonin, could explain the exhaustion related to OR through increased serotonin  
37  
38 synthesis in the brain. It is known that free tryptophan competes with BCAAs for the  
39  
40 same carrier to pass through the blood-brain barrier and, on the other hand, with fatty  
41  
42 acids for albumin. Thus exercise-induced energy metabolism has been suggested to favor  
43  
44 the transport of free tryptophan to the brain as the utilization of BCAAs and fatty acids  
45  
46 increases. This, in turn, could elevate the rate of serotonin synthesis in the brain, leading  
47  
48 to a feeling of fatigue which is a characteristic symptom of OR<sup>9, 10, 50</sup>. In fact, both  
49  
50 mechanisms would lead to increased unbound tryptophan concentrations in the plasma. In  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 contrast, we did not observe any significant increase in tryptophan in OR **nor any**  
4 **significant changes in the levels of BCAAs.** Thus, tryptophan does not seem to be a  
5  
6 sensitive marker for the early changes of OR.  
7  
8  
9

10  
11  
12 **Although we did not observe any increase in tryptophan levels in OR, we found that its**  
13 **levels did decline in both groups throughout the training. Acute exercise has been**  
14 **reported to decrease the tryptophan levels in trained athletes<sup>51, 52</sup>. However, as far as we**  
15 **are aware, there are only two studies, which have investigated the effect of chronic**  
16 **exercise on tryptophan levels among healthy subjects<sup>53</sup> or athletes<sup>54</sup>. Melancon et. al**  
17 **found that plasma tryptophan availability was elevated by 30 and 60 minutes of acute**  
18 **exercise after 16-week aerobic exercise training among 60 aged men<sup>53</sup>. However, free**  
19 **tryptophan availability measured during the bouts of acute exercise was attenuated**  
20 **following training. The researchers suggested this occurred possibly due to a lower**  
21 **serotonergic and sympathetic response due to regular exercising<sup>53</sup>. Thus, a similar**  
22 **process might explain the decreased tryptophan levels in our study. Additionally, a**  
23 **randomized double-blinded placebo-controlled trial of a probiotic among trained athletes**  
24 **found that tryptophan levels were lower after 12 weeks of normal winter training in the**  
25 **placebo group. The placebo group also experienced more upper respiratory infections**  
26 **than the probiotic group indicating that tryptophan may have a possible role in the**  
27 **regulation of immune system, although confounding factors could not be ruled out in that**  
28 **trial<sup>54</sup>. Endurance training has also been reported to cause adaptations to the kynurenine**  
29 **pathway and in the activity of indoleamine 2,3-dioxygenase, which is the enzyme**  
30 **responsible for the breakdown of tryptophan<sup>55</sup>. Activation of kynurenine pathway and**  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 decreased tryptophan levels were also reported during acute exercise<sup>52</sup>. Therefore, the  
4 reduction in the tryptophan levels observed in the present study could also be a result of  
5 intense training activating the kynurenine pathway and might contribute to the high  
6 prevalence of symptoms of URTI among conscripts.  
7  
8  
9  
10  
11  
12

13  
14 Tyrosine is a precursor of the neurotransmitter dopamine, which takes part in excitatory  
15 neurotransmission in the reward-motivation system of the brain and has an important role  
16 in motivation, reward, memory and attention<sup>56</sup>. A reduction of the dopamine  
17 concentration in the brain as a result of prolonged exercise has been suggested to be  
18 linked with the onset of central fatigue based on the studies conducted with rats<sup>57-59</sup> or  
19 humans<sup>60, 61</sup>. Already in the 1990s, Davis and Bailey<sup>56</sup> proposed the central fatigue theory  
20 of BCAAs and serotonin, and postulated that a high serotonin-dopamine ratio could be  
21 related to impaired physical performance, tiredness and onset of fatigue whereas a low  
22 ratio was associated with improved performance as a consequence of maintenance of  
23 motivation and arousal through dopaminergic neuronal pathways. Furthermore, previous  
24 studies have also shown that long-term exercise modifies the activity of the dopaminergic  
25 system in patients with Parkinson disease<sup>62-64</sup>. Thus, the observed increase in tyrosine  
26 levels throughout training in nOR might be a physiological response to the alterations of  
27 neuronal pathways in the dopaminergic system whereas in OR, the exercise-induced  
28 plasticity might be disrupted due to pathophysiological changes in brain neurotransmitter  
29 levels which also lead to a sensation of fatigue.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Tyrosine is also a precursor of the neurotransmitter noradrenaline, which plays a role in  
4 learning and memory, attention, arousal, anxiety and pain<sup>65</sup>. Studies performed with  
5 noradrenaline reuptake inhibitors have demonstrated that the noradrenergic system  
6 decreases performance rather than improves it in humans<sup>66-68</sup>. Although the earlier studies  
7 have not reported any improvement in physical performance after tyrosine  
8 supplementation<sup>69-73</sup>, some trials suggest that ingestion of tyrosine can enhance working  
9 memory, cognitive tasks and tolerance to stress under stressed situations, such as military  
10 operations<sup>74, 75</sup> or sleep deprivation<sup>76, 77</sup>, through adrenergic pathways. Consequently, our  
11 findings of no change in tyrosine in OR, in contrast to an increase in nOR throughout  
12 training, may reflect a maladaptation to the increased exercise stress. The increased  
13 tyrosine levels in nOR, in turn, might also represent a physiological adaptation in the  
14 conscripts to the challenges posed by stressful military conditions resulting in better  
15 cognitive functioning.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35 The determination of OR and OTS is challenging and, therefore, our findings should be  
36 considered with caution. Due to logistic reasons, we were unable to take the baseline  
37 blood samples on day 1 with the first samples being taken on day 7, which might slightly  
38 affect the present results. Furthermore, OR is a consequence of complex set of stressors  
39 and individually altered symptoms are also affected by other factors including  
40 environmental and social circumstances, which makes it challenging to detect even using  
41 a variety of criteria. Furthermore, amino acids form a complex metabolic network which  
42 might be changed not only on strenuous exercise but also by many other factors such as  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 nutrition, therefore the possible clarification of the mechanism behind the  
4  
5 pathophysiology of OR and OTS still remains unclear.  
6  
7  
8  
9

## 10 11 **Conclusion** 12 13

14  
15 In conclusion, the present study indicates that two metabolically active amino acids,  
16  
17 **alanine and arginine**, might be predisposing factors for OR due to early observed  
18  
19 decrease in their levels in OR. In addition, changes in the levels of the neuronal  
20  
21 precursors, tryptophan and tyrosine, might be involved in the pathophysiology of OR  
22  
23 although it does seem that tryptophan might not be a sensitive marker for early changes  
24  
25 of OR. Moreover, an increase in the arginine concentration in response to military  
26  
27 training may be indicative of a positive training adaptation. The results of our study also  
28  
29 confirm the possible use of the Gln/Glu ratio as a tool for evaluating OR due to  
30  
31 significant increase in the Glu concentration which may reflect the exercise induced  
32  
33 fatigue and OR. **However, further investigations will be needed with a determination of**  
34  
35 **other biochemical parameters, which might allow a clear interpretation of the changes in**  
36  
37 **the levels of the amino acids.** Moreover, because approximately 15 % of conscripts were  
38  
39 classified as OR, more attention should be paid to the individual response to military  
40  
41 training, for example to have more tolerable training intensities, especially during the first  
42  
43 four weeks of BT.  
44  
45  
46  
47  
48  
49  
50

51  
52 **Authors' contributions:** MMT and HK designed the study, MMT performed tests and  
53  
54 collected the data. VK analyzed the concentrations of amino acids in blood. NJJI, RIAJ,  
55  
56  
57  
58  
59  
60

1  
2  
3 MA, AU and MMT chose and reviewed the OR criteria based on the literature. NJJI and  
4  
5 RIAJ performed the statistical analysis. NJJI wrote the manuscript and all the authors  
6  
7 participated in reviewing it.  
8  
9

## 10 11 12 **ACKNOWLEDGEMENTS** 13 14 15

16  
17 We would like to thank the Finnish Ministry of Education, Finnish Cultural Foundation,  
18  
19 Polar Electro Oy and the Scientific Advisory Board for Defense for their financial  
20  
21 support of the research.  
22  
23  
24  
25

## 26 **DECLARATION OF CONFLICTING INTERESTS** 27 28 29

30 The author(s) declared no potential conflicts of interest with respect to the research,  
31  
32 authorship and/or publication of this article.  
33  
34  
35  
36

## 37 **FUNDING** 38 39 40

41 This work was supported by the Finnish Ministry of Education; Finnish Cultural  
42  
43 Foundation; Polar Electro Oy; and the Scientific Advisory Board for Defense. The  
44  
45 funders had no role in study design, data collection and analysis, decision to publish or  
46  
47 preparation of the manuscript.  
48  
49  
50

## 51 52 53 **ORCID iD** 54 55 56 57 58 59 60

Mustafa Atalay <https://orcid.org/0000-0001-8999-1426>

### Figure legend

Figure 1. Mean  $\pm$  SD concentration at rest for significantly changed amino acids i.e. glutamine-glutamate ratio (Gln/Glu) and levels of glutamate (Glu), arginine, alanine, tyrosine and tryptophan in non-overreached (nOR) and overreached (OR) conscripts at weeks 1, 4, 7 and 9 of the military service. Significant difference between groups \* $p < 0.05$ . Significant change due to exercise # $p < 0.05$ , ### $p < 0.001$ . Group x training interaction + $p < 0.05$ .

### REFERENCES

1. Meeusen R, Duclos M, Foster C, Fry A, Gleeson M, Nieman D, Raglin J, Rietjens G, Steinacker J, Urhausen A, European College of Sport Science, American College of Sports Medicine. Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc* 2013;**45**:186-205
2. Coutts A, Reaburn P, Piva TJ, Murphy A. Changes in selected biochemical, muscular strength, power, and endurance measures during deliberate overreaching and tapering in rugby league players. *Int J Sports Med* 2007;**28**:116-24
3. Coutts AJ, Reaburn P, Piva TJ, Rowsell GJ. Monitoring for overreaching in rugby league players. *Eur J Appl Physiol* 2007;**99**:313-24
4. Halson SL, Lancaster GI, Jeukendrup AE, Gleeson M. Immunological responses to overreaching in cyclists. *Med Sci Sports Exerc* 2003;**35**:854-61
5. Smith DJ, Norris SR. Changes in glutamine and glutamate concentrations for tracking training tolerance. *Med Sci Sports Exerc* 2000;**32**:684-9
6. Walsh NP, Blannin AK, Robson PJ, Gleeson M. Glutamine, exercise and immune function. Links and possible mechanisms. *Sports Med* 1998;**26**:177-91

- 1  
2  
3 7. Gabriel HH, Urhausen A, Valet G, Heidelbach U, Kindermann W. Overtraining and  
4 immune system: a prospective longitudinal study in endurance athletes. *Med Sci Sports*  
5 *Exerc* 1998;**30**:1151-7  
6
- 7  
8 8. Mackinnon LT, Hooper SL. Plasma glutamine and upper respiratory tract infection  
9 during intensified training in swimmers. *Med Sci Sports Exerc* 1996;**28**:285-90  
10
- 11 9. Newsholme EA, Blomstrand E. Branched-chain amino acids and central fatigue. *J Nutr*  
12 2006;**136**:274S-6S  
13
- 14 10. Meeusen R, Watson P, Hasegawa H, Roelands B, Piacentini MF. Brain  
15 neurotransmitters in fatigue and overtraining. *Appl Physiol Nutr Metab* 2007;**32**:857-64  
16  
17
- 18 11. Tanskanen MM, Kyrolainen H, Uusitalo AL, Huovinen J, Nissila J, Kinnunen H,  
19 Atalay M, Hakkinen K. Serum sex hormone-binding globulin and cortisol concentrations  
20 are associated with overreaching during strenuous military training. *J Strength Cond Res*  
21 2011;**25**:787-97  
22
- 23 12. Tanskanen M, Uusitalo AL, Hakkinen K, Nissila J, Santtila M, Westerterp KR,  
24 Kyrolainen H. Aerobic fitness, energy balance, and body mass index are associated with  
25 training load assessed by activity energy expenditure. *Scand J Med Sci Sports*  
26 2009;**19**:871-8  
27  
28
- 29 13. Terry PC, Lane AM, Lane HJ, Keohane L. Development and validation of a mood  
30 measure for adolescents. *J Sports Sci* 1999;**17**:861-72  
31  
32
- 33 14. American College of Sports Medicine. Guidelines for exercise testing and  
34 prescription, 6th ed. Baltimore: Lippincott Williams & Wilkins, 2001, pp. 68,117,303  
35  
36
- 37 15. Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, Pratt  
38 M, Ekelund U, Yngve A, Sallis JF, Oja P. International physical activity questionnaire:  
39 12-country reliability and validity. *Med Sci Sports Exerc* 2003;**35**:1381-95  
40
- 41 16. Mero A, Leikas A, Knuutinen J, Hulmi JJ, Kovanen V. Effect of strength training  
42 session on plasma amino acid concentration following oral ingestion of leucine, BCAAs  
43 or glutamine in men. *Eur J Appl Physiol* 2009;**105**:215-23  
44  
45
- 46 17. Decroix L, Lamberts RP, Meeusen R. Can the Lamberts and Lambert Submaximal  
47 Cycle Test Reflect Overreaching in Professional Cyclists?. *Int J Sports Physiol Perform*  
48 2018;**13**:23-8  
49
- 50 18. Siegl A, M Kosel E, Tam N, Koschnick S, Langerak NG, Skorski S, Meyer T,  
51 Lamberts RP. Submaximal Markers of Fatigue and Overreaching; Implications for  
52 Monitoring Athletes. *Int J Sports Med* 2017;**38**:675-82  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 19. Halson SL, Bridge MW, Meeusen R, Busschaert B, Gleeson M, Jones DA,  
4 Jeukendrup AE. Time course of performance changes and fatigue markers during  
5 intensified training in trained cyclists. *J Appl Physiol (1985)* 2002;**93**:947-56  
6  
7  
8 20. Uusitalo AL. Overtraining: making a difficult diagnosis and implementing targeted  
9 treatment. *Phys Sportsmed* 2001;**29**:35-50  
10  
11 21. Budgett R. Fatigue and underperformance in athletes: the overtraining syndrome. *Br J*  
12 *Sports Med* 1998;**32**:107-10  
13  
14 22. Kentta G, Hassmen P, Raglin JS. Training practices and overtraining syndrome in  
15 Swedish age-group athletes. *Int J Sports Med* 2001;**22**:460-5  
16  
17 23. Matos NF, Winsley RJ, Williams CA. Prevalence of nonfunctional  
18 overreaching/overtraining in young English athletes. *Med Sci Sports Exerc*  
19 2011;**43**:1287-94  
20  
21 24. Garcin M, Fleury A, Billat V. The ratio HLa : RPE as a tool to appreciate  
22 overreaching in young high-level middle-distance runners. *Int J Sports Med* 2002;**23**:16-  
23 21  
24  
25 25. Cadegiani FA, Kater CE. Body composition, metabolism, sleep, psychological and  
26 eating patterns of overtraining syndrome: Results of the EROS study (EROS-PROFILE).  
27 *J Sports Sci* 2018;**36**:1902-10  
28  
29 26. Comotto S, Bottoni A, Moci E, Piacentini MF. Analysis of session-RPE and profile  
30 of mood states during a triathlon training camp. *J Sports Med Phys Fitness* 2015;**55**:361-  
31 7  
32  
33 27. Korzeniewski K, Nitsch-Osuch A, Konior M, Lass A. Respiratory tract infections in  
34 the military environment. *Respir Physiol Neurobiol* 2015;**209**:76-80  
35  
36 28. Ryan MA, Christian RS, Wohlrabe J. Handwashing and respiratory illness among  
37 young adults in military training. *Am J Prev Med* 2001;**21**:79-83  
38  
39 29. Rowbottom DG, Keast D, Goodman C, Morton AR. The haematological, biochemical  
40 and immunological profile of athletes suffering from the overtraining syndrome. *Eur J*  
41 *Appl Physiol Occup Physiol* 1995;**70**:502-9  
42  
43 30. Parry-Billings M, Budgett R, Koutedakis Y, Blomstrand E, Brooks S, Williams C,  
44 Calder PC, Pilling S, Baigrie R, Newsholme EA. Plasma amino acid concentrations in the  
45 overtraining syndrome: possible effects on the immune system. *Med Sci Sports Exerc*  
46 1992;**24**:1353-8  
47  
48 31. Castell LM, Poortmans JR, Newsholme EA. Does glutamine have a role in reducing  
49 infections in athletes?. *Eur J Appl Physiol Occup Physiol* 1996;**73**:488-90  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




- 1  
2  
3 32. Kingsbury KJ, Kay L, Hjelm M. Contrasting plasma free amino acid patterns in elite  
4 athletes: association with fatigue and infection. *Br J Sports Med* 1998;**32**:25,32;  
5 discussion 32-3  
6  
7  
8 33. Hohl R, Ferrareso RL, De Oliveira RB, Lucco R, Brenzikofer R, De Macedo DV.  
9 Development and characterization of an overtraining animal model. *Med Sci Sports Exerc*  
10 2009;**41**:1155-63  
11  
12 34. Moncada S. Nitric oxide in the vasculature: physiology and pathophysiology. *Ann N*  
13 *Y Acad Sci* 1997;**811**:60,7; discussion 67-9  
14  
15 35. Wu G, Morris SM,Jr. Arginine metabolism: nitric oxide and beyond. *Biochem J*  
16 1998;**336(Pt 1)**:1-17  
17  
18  
19 36. Reid MB. Role of nitric oxide in skeletal muscle: synthesis, distribution and  
20 functional importance. *Acta Physiol Scand* 1998;**162**:401-9  
21  
22  
23 37. Tsukiyama Y, Ito T, Nagaoka K, Eguchi E, Ogino K. Effects of exercise training on  
24 nitric oxide, blood pressure and antioxidant enzymes. *J Clin Biochem Nutr* 2017;**60**:180-  
25 6  
26  
27  
28 38. Brosnan JT, Brosnan ME. Creatine: endogenous metabolite, dietary, and therapeutic  
29 supplement. *Annu Rev Nutr* 2007;**27**:241-61  
30  
31 39. Schaefer A, Piquard F, Geny B, Doutreleau S, Lampert E, Mettauer B, Lonsdorfer J.  
32 L-arginine reduces exercise-induced increase in plasma lactate and ammonia. *Int J Sports*  
33 *Med* 2002;**23**:403-7  
34  
35  
36 40. Wideman L, Weltman JY, Hartman ML, Veldhuis JD, Weltman A. Growth hormone  
37 release during acute and chronic aerobic and resistance exercise: recent findings. *Sports*  
38 *Med* 2002;**32**:987-1004  
39  
40  
41 41. Kanaley JA. Growth hormone, arginine and exercise. *Curr Opin Clin Nutr Metab*  
42 *Care* 2008;**11**:50-4  
43  
44  
45 42. Zhou M, Martindale RG. Arginine in the critical care setting. *J Nutr* 2007;**137**:1687S-  
46 92S  
47  
48  
49 43. Morris SM,Jr. Recent advances in arginine metabolism: roles and regulation of the  
50 arginases. *Br J Pharmacol* 2009;**157**:922-30  
51  
52 44. Felig P, Wahren J. Amino acid metabolism in exercising man. *J Clin Invest*  
53 1971;**50**:2703-14  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 45. Einspahr KJ, Tharp G. Influence of endurance training on plasma amino acid  
4 concentrations in humans at rest and after intense exercise. *Int J Sports Med*  
5 1989;**10**:233-6  
6  
7  
8 46. Brodan V, Kuhn E, Pechar J, Tomkova D. Changes of free amino acids in plasma of  
9 healthy subjects induced by physical exercise. *Eur J Appl Physiol Occup Physiol*  
10 1976;**35**:69-77  
11  
12 47. Refsum HE, Gjessing LR, Stromme SB. Changes in plasma amino acid distribution  
13 and urine amino acids excretion during prolonged heavy exercise. *Scand J Clin Lab*  
14 *Invest* 1979;**39**:407-13  
15  
16  
17 48. Buyse L, Decroix L, Timmermans N, Barbe K, Verrelst R, Meeusen R. Improving the  
18 Diagnosis of Nonfunctional Overreaching and Overtraining Syndrome. *Med Sci Sports*  
19 *Exerc* 2019;**51**:2524-30  
20  
21 49. Meeusen R, Nederhof E, Buyse L, Roelands B, de Schutter G, Piacentini MF.  
22 Diagnosing overtraining in athletes using the two-bout exercise protocol. *Br J Sports Med*  
23 2010;**44**:642-8  
24  
25  
26 50. Meeusen R, Watson P, Hasegawa H, Roelands B, Piacentini MF. Central fatigue: the  
27 serotonin hypothesis and beyond. *Sports Med* 2006;**36**:881-909  
28  
29  
30 51. Areces F, Gonzalez-Millan C, Salinero JJ, Abian-Vicen J, Lara B, Gallo-Salazar C,  
31 Ruiz-Vicente D, Del Coso J. Changes in Serum Free Amino Acids and Muscle Fatigue  
32 Experienced during a Half-Ironman Triathlon. *PLoS One* 2015;**10**:e0138376  
33  
34 52. Strasser B, Geiger D, Schauer M, Gatterer H, Burtscher M, Fuchs D. Effects of  
35 Exhaustive Aerobic Exercise on Tryptophan-Kynurenine Metabolism in Trained  
36 Athletes. *PLoS One* 2016;**11**:e0153617  
37  
38  
39 53. Melancon MO, Lorrain D, Dionne IJ. Changes in markers of brain serotonin activity  
40 in response to chronic exercise in senior men. *Appl Physiol Nutr Metab* 2014;**39**:1250-6  
41  
42  
43 54. Strasser B, Geiger D, Schauer M, Gostner JM, Gatterer H, Burtscher M, Fuchs D.  
44 Probiotic Supplements Beneficially Affect Tryptophan-Kynurenine Metabolism and  
45 Reduce the Incidence of Upper Respiratory Tract Infections in Trained Athletes: A  
46 Randomized, Double-Blinded, Placebo-Controlled Trial. *Nutrients* 2016;**8**:752  
47  
48 55. Schlittler M, Goigny M, Agudelo LZ, Venckunas T, Brazaitis M, Skurvydas A,  
49 Kamandulis S, Ruas JL, Erhardt S, Westerblad H, Andersson DC. Endurance exercise  
50 increases skeletal muscle kynurenine aminotransferases and plasma kynurenic acid in  
51 humans. *Am J Physiol Cell Physiol* 2016;**310**:C836-40  
52  
53  
54 56. Davis JM, Bailey SP. Possible mechanisms of central nervous system fatigue during  
55 exercise. *Med Sci Sports Exerc* 1997;**29**:45-57  
56  
57  
58  
59  
60



- 1  
2  
3 57. Gerald MC. Effects of (+)-amphetamine on the treadmill endurance performance of  
4 rats. *Neuropharmacology* 1978;**17**:703-4  
5  
6  
7 58. Heyes MP, Garnett ES, Coates G. Central dopaminergic activity influences rats  
8 ability to exercise. *Life Sci* 1985;**36**:671-7  
9  
10 59. Zheng X, Hasegawa H. Administration of caffeine inhibited adenosine receptor  
11 agonist-induced decreases in motor performance, thermoregulation, and brain  
12 neurotransmitter release in exercising rats. *Pharmacol Biochem Behav* 2016;**140**:82-9  
13  
14 60. Roelands B, Hasegawa H, Watson P, Piacentini MF, Buyse L, De Schutter G,  
15 Meeusen RR. The effects of acute dopamine reuptake inhibition on performance. *Med Sci*  
16 *Sports Exerc* 2008;**40**:879-85  
17  
18 61. Watson P, Hasegawa H, Roelands B, Piacentini MF, Looverie R, Meeusen R. Acute  
19 dopamine/noradrenaline reuptake inhibition enhances human exercise performance in  
20 warm, but not temperate conditions. *J Physiol* 2005;**565**:873-83  
21  
22 62. Palmer SS, Mortimer JA, Webster DD, Bistevins R, Dickinson GL. Exercise therapy  
23 for Parkinson's disease. *Arch Phys Med Rehabil* 1986;**67**:741-5  
24  
25 63. Herman T, Giladi N, Gruendlinger L, Hausdorff JM. Six weeks of intensive treadmill  
26 training improves gait and quality of life in patients with Parkinson's disease: a pilot  
27 study. *Arch Phys Med Rehabil* 2007;**88**:1154-8  
28  
29 64. Sacheli MA, Neva JL, Lakhani B, Murray DK, Vafai N, Shahinfard E, English C,  
30 McCormick S, Dinelle K, Neilson N, McKenzie J, Schulzer M, McKenzie DC, Appel-  
31 Cresswell S, McKeown MJ, Boyd LA, Sossi V, Stoessl AJ. Exercise increases caudate  
32 dopamine release and ventral striatal activation in Parkinson's disease. *Mov Disord*  
33 2019;**34**:1891-900  
34  
35 65. Meeusen R, Roelands B. Fatigue: Is it all neurochemistry?. *Eur J Sport Sci*  
36 2018;**18**:37-46  
37  
38 66. Roelands B, Goekint M, Heyman E, Piacentini MF, Watson P, Hasegawa H, Buyse L,  
39 Pauwels F, De Schutter G, Meeusen R. Acute norepinephrine reuptake inhibition  
40 decreases performance in normal and high ambient temperature. *J Appl Physiol (1985)*  
41 2008;**105**:206-12  
42  
43 67. Klass M, Duchateau J, Rabec S, Meeusen R, Roelands B. Noradrenaline Reuptake  
44 Inhibition Impairs Cortical Output and Limits Endurance Time. *Med Sci Sports Exerc*  
45 2016;**48**:1014-23  
46  
47 68. Klass M, Roelands B, Levenez M, Fontenelle V, Pattyn N, Meeusen R, Duchateau J.  
48 Effects of noradrenaline and dopamine on supraspinal fatigue in well-trained men. *Med*  
49 *Sci Sports Exerc* 2012;**44**:2299-308  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 69. Struder HK, Hollmann W, Platen P, Donike M, Gotzmann A, Weber K. Influence of  
4 paroxetine, branched-chain amino acids and tyrosine on neuroendocrine system responses  
5 and fatigue in humans. *Horm Metab Res* 1998;**30**:188-94  
6  
7  
8 70. Chinevere TD, Sawyer RD, Creer AR, Conlee RK, Parcell AC. Effects of L-tyrosine  
9 and carbohydrate ingestion on endurance exercise performance. *J Appl Physiol (1985)*  
10 2002;**93**:1590-7  
11  
12 71. Sutton EE, Coill MR, Deuster PA. Ingestion of tyrosine: effects on endurance, muscle  
13 strength, and anaerobic performance. *Int J Sport Nutr Exerc Metab* 2005;**15**:173-85  
14  
15  
16 72. Tumilty L, Davison G, Beckmann M, Thatcher R. Failure of oral tyrosine  
17 supplementation to improve exercise performance in the heat. *Med Sci Sports Exerc*  
18 2014;**46**:1417-25  
19  
20  
21 73. Watson P, Enever S, Page A, Stockwell J, Maughan RJ. Tyrosine supplementation  
22 does not influence the capacity to perform prolonged exercise in a warm environment. *Int*  
23 *J Sport Nutr Exerc Metab* 2012;**22**:363-73  
24  
25  
26 74. Owasoyo JO, Neri DF, Lamberth JG. Tyrosine and its potential use as a  
27 countermeasure to performance decrement in military sustained operations. *Aviat Space*  
28 *Environ Med* 1992;**63**:364-9  
29  
30  
31 75. Deijen JB, Wientjes CJ, Vullings HF, Cloin PA, Langefeld JJ. Tyrosine improves  
32 cognitive performance and reduces blood pressure in cadets after one week of a combat  
33 training course. *Brain Res Bull* 1999;**48**:203-9  
34  
35  
36 76. Dollins AB, Krock LP, Storm WF, Wurtman RJ, Lieberman HR. L-tyrosine  
37 ameliorates some effects of lower body negative pressure stress. *Physiol Behav*  
38 1995;**57**:223-30  
39  
40  
41 77. Neri DF, Wiegmann D, Stanny RR, Shappell SA, McCardie A, McKay DL. The  
42 effects of tyrosine on cognitive performance during extended wakefulness. *Aviat Space*  
43 *Environ Med* 1995;**66**:313-9  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

Table 1. Research protocol during the first 10 weeks of the military training.

	Week									
	1	2	3	4	5	6	7	8	9	10
VO <sub>2max</sub> test	X				X			X		X
Submaximal marching test	X			X			X		X	
POMS	X	X		X			X	X	X	
Questionnaire							X	X	X	
Inquiry about symptoms of URTI	X			X		X	X	X	X	
Blood samples	X			X			X		X	

Table 2. The descriptive statistics of non-OR and OR conscripts at week 1.

	Non-overreached	Overreached	P*
<b>*Total Physical activity</b> (MET·min·wk <sup>-1</sup> )	3607 ± 3403	7808 ± 11026	<b>0.40</b>
VO <sub>2</sub> max (mL·kg <sup>-1</sup> ·min <sup>-1</sup> )	43.0 ± 6.5	45.0 ± 9.4	<b>0.57</b>
Height (cm)	177.6 ± 6.7	176.6 ± 6.1	<b>0.583</b>
<b>Body mass (kg)</b>	77.9 ± 12.9	76.9 ± 18.3	<b>0.85</b>
BMI (kg/m <sup>2</sup> )	24.7 ± 3.8	24.5 ± 4.4	<b>0.70</b>
Fatmass (kg)	15.1 ± 8.3	17.1 ± 11.6	<b>0.95</b>
Fatfreemass (kg)	62.8 ± 8.5	59.8 ± 7.7	<b>0.36</b>
Body fat percentage	18.7 ± 1.1	20.6 ± 3.1	<b>0.65</b>

**\*Total PA: Light PA + Moderate PA + Vigorous PA reported by international physical activity questionnaire (IPAQ).**

Table 3. Descriptive statistics of non-significantly changed amino acids at rest (nmol/ml) among nOR and OR conscripts during the first 9 weeks of the military service.

		Week 1	Week 4	Week 7	Week 9	P*
		Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
Asparagine	nOR	85.8 ± 25.3	82.4 ± 25.4	88.2 ± 25.6	86.5 ± 26.8	0.27
	OR	66.1 ± 19.3	60.4 ± 21.8	65.2 ± 20.9	69.4 ± 16.0	
Aspartate	nOR	4.1 ± 2.2	4.0 ± 1.7	4.0 ± 1.9	4.7 ± 1.7	0.073
	OR	6.0 ± 3.7	4.8 ± 2.4	4.7 ± 1.7	6.1 ± 3.0	
Glutamine	nOR	641.8 ± 72.7	622.8 ± 80.7	675.8 ± 80.8	646.1 ± 86.7	0.12
	OR	588.0 ± 52.6	559.0 ± 76.7	637.3 ± 94.2	600.7 ± 39.8	
Glycine	nOR	257.5 ± 92.8	250.2 ± 83.7	264.7 ± 88.8	271.8 ± 105.7	0.19
	OR	240.6 ± 19.3	220.7 ± 35.0	254.8 ± 53.4	294.5 ± 62.1	
Histidine	nOR	73.2 ± 10.3	73.1 ± 7.9	72.7 ± 8.9	71.4 ± 10.2	0.44
	OR	76.1 ± 4.8	72.8 ± 13.3	72.9 ± 13.9	74.9 ± 7.3	
Isoleucine	nOR	55.5 ± 8.9	54.2 ± 8.0	53.6 ± 8.1	54.9 ± 9.6	0.094
	OR	59.2 ± 9.8	60.4 ± 10.6	55.8 ± 9.2	52.9 ± 6.5	
Leusine	nOR	151.5 ± 21.7	150.7 ± 21.9	153.0 ± 18.0	154.5 ± 23.0	0.36
	OR	158.0 ± 13.5	152.8 ± 18.1	153.5 ± 23.9	145.1 ± 21.6	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Lysine	nOR	146.9 ± 23.9	134.3 ± 25.6	151.9 ± 24.4	148.4 ± 29.9	0.65
	OR	146.6 ± 24.8	122.4 ± 35.5	144.6 ± 44.6	150.2 ± 35.7	
Methionine	nOR	29.0 ± 3.9	25.9 ± 4.6	29.7 ± 4.3	28.4 ± 5.8	0.59
	OR	28.7 ± 3.4	21.3 ± 3.2	28.6 ± 3.9	24.6 ± 3.2	
Phenylalanine	nOR	59.6 ± 7.8	57.4 ± 7.2	60.5 ± 6.4	62.3 ± 8.7	0.61
	OR	60.3 ± 2.7	52.0 ± 5.8	60.1 ± 7.1	54.4 ± 8.6	
Serine	nOR	80.7 ± 15.9	77.6 ± 16.5	83.9 ± 16.9	81.9 ± 20.1	0.79
	OR	84.9 ± 13.8	74.9 ± 12.0	80.2 ± 12.2	86.6 ± 15.5	
Taurine	nOR	37.1 ± 7.8	36.2 ± 7.4	33.8 ± 6.7	60.7 ± 9.1	0.91
	OR	39.0 ± 11.2	34.7 ± 8.6	33.9 ± 5.8	37.8 ± 9.9	
Treonine	nOR	59.4 ± 8.0	55.6 ± 10.7	60.8 ± 11.8	58.8 ± 12.4	0.33
	OR	56.2 ± 9.4	50.2 ± 7.3	52.3 ± 12.3	56.8 ± 13.0	
Valine	nOR	217.9 ± 34.9	216.0 ± 35.5	213.2 ± 26.2	226.2 ± 40.4	0.83
	OR	218.7 ± 18.3	217.8 ± 34.8	205.7 ± 25.7	214.7 ± 14.9	

Note: The data is presented in the form of mean ± SD.