

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Khair, Raad M.; Stenroth, Lauri; Cronin, Neil J.; Reito, Aleksii; Paloneva, Juha; Finni, Taija

Title: In vivo localized gastrocnemius subtendon representation within the healthy and ruptured human Achilles tendon

Year: 2022

Version: Accepted version (Final draft)

Copyright: © 2022, Journal of Applied Physiology

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Khair, R. M., Stenroth, L., Cronin, N. J., Reito, A., Paloneva, J., & Finni, T. (2022). In vivo localized gastrocnemius subtendon representation within the healthy and ruptured human Achilles tendon. *Journal of Applied Physiology*, 133(1), 11-19.
<https://doi.org/10.1152/jappphysiol.00084.2022>

Achilles MG and LG subtendon representation

1 **In vivo localised gastrocnemius subtendon representation within the healthy and** 2 **ruptured human Achilles tendon**

3 Ra'ad M. Khair ^{1*}, Lauri Stenroth ², Neil J. Cronin ^{1,4}, Aleksi Reito ³, Juha Paloneva ³ and Taija Finni ¹.

4 ¹ Faculty of Sport and Health Sciences, Neuromuscular Research Center, University of Jyväskylä,
5 Jyväskylä, Finland; ² Department of Applied Physics, University of Eastern Finland, Kuopio, Finland;
6 ³ Central Finland Health Care District, Finland and University of Eastern Finland, Finland, ⁴ School of
7 Sport and Exercise, University of Gloucestershire, UK.

8
9
10 *Author for Correspondence

11 Ra'ad M. Khair

12 Email: raad.m.khair@jyu.fi

13 Phone : +358469221362

14 P.O. Box 35

15 40014 Jyväskylä, Finland

16
17
18 Author contributions

19 Conceptualization: T.F., N.J.C., A.R., J.P., R.M.K; Methodology: T.F., N.J.C., A.R., L.S., J.P.; Data
20 acquisition: T.F., R.M.K.; Data curation - analysis: R.M.K.; Writing original draft: R.M.K.; Writing –
21 review & editing: R.M.K., L.S., N.J.C., A.R., J.P., T.F.; Visualization: R.M.K.; Supervision: T.F.,
22 N.J.C.; Funding acquisition: T.F; Project administration: T.F.

23

24 **Abstract**

25 The Achilles tendon (AT) is composed of three distinct in-series elastic subtendons, arising
26 from different muscles in the triceps surae. Independent activation of any of these muscles is
27 thought to induce sliding between the adjacent AT subtendons. We aimed to investigate
28 displacement patterns during voluntary contraction (VOL) and selective transcutaneous
29 stimulation of medial (MG_{stim}) and lateral (LG_{stim}) gastrocnemius between ruptured and
30 healthy tendons, and to examine the representative areas of AT subtendons. Twenty-eight
31 patients with unilateral AT rupture performed bilateral VOL at 30% of the maximal
32 isometric un-injured plantarflexion torque. AT displacement was analysed from sagittal B-
33 mode ultrasonography images during VOL, MG_{stim} and LG_{stim}. Three-way ANOVA
34 revealed a significant two-way interaction of contraction type*location on the tendon
35 displacement ($F(10-815)=3.72$, $p<0.001$). The subsequent two-way analysis revealed a
36 significant contraction type*location interaction for tendon displacement ($F(10-410)=3.79$,
37 $p<0.001$) in the un-injured limb only, where LG_{stim} displacement pattern was significantly
38 different from MG_{stim} ($p=0.008$) and VOL ($p=0.005$). When comparing contraction types
39 between limbs there were no difference in the displacement patterns, but displacement
40 amplitudes differed. There was no significant difference in the location of maximum or
41 minimum displacement between limbs. The displacement pattern was not different in non-
42 surgically treated compared to un-injured tendons one-year post rupture. Our results suggest
43 that near the calcaneus, LG subtendon is located in the most anterior region adjacent to
44 medial gastrocnemius. However, free tendon stiffness seems to be lower in the injured AT,
45 leading to more displacement during electrically-induced contractions compared to the un-
46 injured.

47

Achilles MG and LG subtendon representation

48 **New & Noteworthy**

49 Using selective electrical stimulation, we report the distributions of medial and lateral
50 gastrocnemius subtendon representations within the healthy and ruptured Achilles tendon. In
51 the majority of our sample, lateral gastrocnemius subtendon was found in the most anterior
52 region adjacent to medial gastrocnemius both in the healthy and ruptured, non-surgically
53 treated tendon. The tendon internal displacement pattern does not seem to differ, but
54 displacement amplitude and non-uniformity differed between healthy and ruptured tendons
55 one-year post rupture.

56

57 **Key words:** Achilles tendon, architecture, geometry, anatomy, rupture, human.

58

59 **Introduction**

60 The Achilles tendon (AT) provides critical series elasticity to the triceps surae, amplifying
61 power for activities such as walking and running (1) and playing a significant role in
62 mechanical energy storage (2). Normal tendon function is disrupted by AT disorders that
63 also cause pain and disability. Achilles tendon rupture (ATR) is prevalent in sport-related
64 activities with an incidence of 31/100,000 individuals per year (3, 4). Understanding the
65 normal and pathological biomechanical function of the AT is crucial to the diagnosis and
66 management of AT-related maladies.

67 The AT has a complex hierarchical structure and is composed of distinct bundles of fascicles
68 running continuously along the tendon, called subtendons. AT subtendons each arises from a
69 different muscular head of the triceps surae: soleus (SOL), medial gastrocnemius (MG), and
70 lateral gastrocnemius (LG) (5, 6). The tendon twists so that at the calcaneal tuberosity
71 insertion, the MG fibres are located on the lateral surface, LG fibres more deeply, and the
72 SOL fibres on the medial surface (5, 6). The degree of twist varies among individuals and can
73 be classified into three types (5). This variation might lead to interindividual differences in
74 the location of the MG, LG and SOL tendon fascicles along the length of the tendon (5). Due
75 to this structure, AT is subjected to complex non-uniform loading that can cause
76 heterogeneity of strain within the tendon (7).

77 *In vivo* studies have exploited advances in ultrasonic imaging and speckle tracking algorithms
78 to reveal non-uniform motion within the AT (8, 9). The ability of subtendons to slide relative
79 to each other is considered to be a function of a healthy tendon (9, 10). Healthy non-
80 uniformity is characterised by smaller displacement of the superficial (posterior) tendon and
81 larger displacement of the deep (anterior) tendon. Assuming that posteriorly the tendon
82 consists of fascicles arising from both gastrocnemius muscles and that the anterior tendon
83 consists of fascicles arising from soleus, researchers have tried to identify structure-function

Achilles MG and LG subtendon representation

84 relationships (11, 12). However, the mechanism of non-uniform displacement and the
85 representation of each subtendon within free AT in different individuals remains elusive due
86 to potential differences in neural control strategies (7), the architecturally complex twisted
87 tendon structure (5, 6), and the difficulty of visualizing individual subtendons using
88 conventional imaging techniques (ultrasound or magnetic resonance imaging).

89 In recent studies, ruptured ATs have been found to display more uniform within-tendon
90 displacement 1-year post-rupture (13, 14). In addition to an increase in length of the tendon,
91 ATR leads to morphomechanical changes in the triceps surae muscles and subtendons (15,
92 16). These changes seem to occur regardless of whether they were treated surgically or
93 conservatively (13, 14), and might alter the force transmission mechanism in the muscle-
94 tendon unit.

95 Voluntary contraction typically activates all synergistic muscles to a variable degree (17, 18)
96 and leads to disproportionate tissue displacement within the tendon due to mechanical and
97 structural differences between triceps surae muscles (19). During voluntary contractions,
98 complex neuromuscular control of the triceps surae within and across healthy and injured
99 individuals may confound interpretations of tissue displacement in adjacent subtendons. By
100 removing the effects of neural control, one could potentially identify if changes in structure
101 and material properties due to ATR modify the displacement pattern within the AT. Electrical
102 transcutaneous stimulation can be used to stimulate a given muscle selectively (20, 21).
103 Using this method, it can be assumed that selective activation of one of the triceps surae
104 muscles induces serial force transmission that is observed as tendon displacement mainly in
105 the area containing tendon fascicles arising from the activated muscle belly. Therefore, the
106 stimulation method may also help to understand AT subtendon organization *in vivo*.

107 By using selective transcutaneous stimulation to medial and lateral gastrocnemius muscles
108 we aimed to find out whether AT tissue displacement pattern differs in voluntary contraction

Achilles MG and LG subtendon representation

109 and electrically evoked contractions between injured (INJ) and un-injured (UNJ) tendons.
110 Examination of the displacement patterns during selective activation was expected to yield
111 information about the representative areas of AT subtendons. We hypothesized that different
112 contraction types would lead to different displacement patterns. Furthermore, it was
113 hypothesized that INJ tendon would show less, and more uniform displacement compared to
114 the UNJ tendon.

Achilles MG and LG subtendon representation

115 **Methods**

116 **Participants**

117 Twenty-eight ATR patients (24 males, 4 females) treated at the Central Finland Health Care
118 District agreed to participate (Table 1). ATR was diagnosed according to the American
119 Academy of Orthopaedic Surgeons guidelines. Inclusion criteria were a minimum of 2 of the
120 following 4 criteria: a positive Thompson test, decreased plantarflexion strength, presence of
121 a palpable gap, and increased passive ankle dorsiflexion with gentle manipulation.
122 Participants with re-occurring rupture were treated surgically and excluded from the sample,
123 which contains only individuals with non-surgical treatment and early mobilization (22). This
124 study was approved by the Ethics committee of Central Finland health care district
125 (2U/2018). Participants signed an informed consent explaining the details of the study,
126 possible risks, and
127 gave permission to use data for research purposes. Participants were invited to the laboratory
128 1-year \pm 1.8 months after rupture.

129

130 **Experimental procedure**

131 B-mode ultrasound was used to examine tendon properties. Scans were done using a 3.6-cm
132 linear probe (UST-5411, Aloka alpha10, Japan). First, the subtendon lengths of MG, LG and
133 SOL were measured from a resting prone position with the subjects' feet over the edge of a
134 table. The limb was scanned to find the most distal point of the muscle-tendon junction of
135 each muscle head and the tendon insertion on the calcaneus, all of which were marked on the
136 skin. The distance between the points was then measured with a measuring tape (23). The
137 reliability of this method was tested, whereby four un-injured limbs was measured on two
138 separate days. The subtendon lengths of the triceps surae muscles were measured and the
139 intraclass correlation coefficient (ICC) was calculated (24). ICC was 0.99 (90% CI 0.97-

Achilles MG and LG subtendon representation

140 0.99) with a coefficient of variation (CV) of 6.6%. Ultrasound imaging was then used to
141 locate the thickest part of both gastrocnemius muscles, where the stimulating electrodes were
142 placed. Participants' skin was shaved and cleaned with alcohol to ensure good conductivity.
143 A pair of 32 mm diameter electrodes (Niva Medical Oy) was attached over each muscle with
144 ~1 cm inter-electrode distance. During measurements, participants sat in a custom-made
145 ankle dynamometer (University of Jyväskylä, Finland) with the hip at 120°, knee at 0° (fully
146 extended), and the ankle and first metatarsophalangeal joints at 90° and 0° respectively. The
147 foot was strapped to the dynamometer pedal and the thigh secured to the seat above the knee.
148 To image tendon displacement, the ultrasound probe was attached longitudinally with the
149 distal edge ~2 cm above the calcaneus.

150 A warm-up was done in the form of a series of standardized submaximal contractions.
151 Starting with UNJ, unilateral maximal voluntary isometric contractions (MVCs) were
152 performed followed by contractions corresponding to 30% of UNJ MVC. Then, with the
153 participant relaxed, single stimulation pulses were elicited with increasing intensity using a
154 constant current electrical stimulator (DS7AH; Digitimer, Hertfordshire, UK) until the motor
155 threshold was exceeded, as confirmed by a visible muscle twitch (20, 21). If a corresponding
156 displacement was not observed clearly in the US image of the AT, higher stimulation
157 intensity was used. AT displacement was imaged 1 s before and throughout a tetanic pulse of
158 1000 μ s at 100 Hz at the pre-determined stimulation intensity. MG and LG were stimulated in
159 random order. The entire protocol was then repeated for INJ, starting with voluntary
160 isometric contractions, followed by electrically induced contractions.

161 Force data were collected via a strain gauge transducer in the foot pedal of the ankle
162 dynamometer. A potentiometer placed under the heel was used to detect heel lift during
163 contractions. Data were sampled at 1 kHz via a 16-bit A/D board (Power 1401, Cambridge
164 Electronic Design, Cambridge, UK) connected to the computer, and signals were recorded

Achilles MG and LG subtendon representation

165 using Spike2 software (Cambridge Electronic Design, Cambridge, UK). To synchronize data,
166 a TTL-pulse was sent manually via Spike2 to first trigger the data acquisition with the US
167 device for 8 seconds and after 1 s to deliver the 0.7 s tetanus to either MG or LG. Ultrasound
168 videos were sampled at 50 HZ and stored for further offline analysis.

169 **Data analysis**

170 Ultrasound B-mode image analysis of tendon displacement was done using a speckle tracking
171 algorithm implemented in Matlab (R2020a, MathWorks Inc, Natick, MA, USA) according to
172 the previously validated and published configuration of Slane and Thelen (9, 25). The region
173 of interest location and size were defined for each subject manually to ensure that only
174 tendon tissue was analysed. A grid of six nodes across the width of the tendon and eleven
175 across the length of the tendon was generated (14). All tracking results were visually
176 inspected to ensure that the nodes remained inside the tendon throughout the movement.
177 Incremental displacements were fitted with a low-order polynomial (25). Displacements of
178 nodes along each of the six antero-posterior rows were averaged and peak displacement of
179 the average data were extracted for analysis. The six locations across the tendon starting from
180 the posterior part to the anterior part are referred to as locations 1-6, respectively. The
181 average peak displacement across the six locations was used to represent mean displacement.
182 Locations of the maximum and minimum displacement were extracted. Tendon non-
183 uniformity was expressed as the difference between minimal and maximal displacement in
184 the tendon. To facilitate the comparison of displacement patterns between electrically
185 induced contractions and volitional activation, the displacement data were normalized to a
186 range between 0-1 where 0 is minimum displacement location and 1 is maximum
187 displacement location. The relative displacement relation between 6 locations across the
188 tendon is hereafter referred to as the displacement pattern. Displacement was normalized
189 since voluntary contraction produced higher torque and overall AT displacement than

Achilles MG and LG subtendon representation

190 electrically induced contractions. Peak torque was calculated for both voluntary and
191 electrically induced contractions.

192

193 **Statistical analysis**

194 Statistical analysis was performed using JASP (JASP version 0.14.1, Amsterdam,
195 Netherlands). The level of significance was set at $p < 0.05$. Three-way repeated-measures
196 ANOVA was performed to investigate the effects of contraction type (VOL, MG, and LG
197 stimulations), limb condition (INJ vs UNJ) and tendon location (across 6 locations) on the
198 normalized displacement of the tendon. The main interest of the analysis is in three- and two-
199 way interaction effects, indicating how the displacements are distributed between the tendon
200 locations (i.e. are affecting the displacement pattern) in the different conditions and limbs. If
201 significant three-way interactions were detected, two-way analysis was performed, followed
202 by simple pairwise comparisons with Bonferroni-adjustment when a significant main effect
203 was found. Greenhouse-Geisser adjustment was applied when the assumption of sphericity
204 was violated. Skewness and kurtosis was checked to insure the normality of the data. If
205 outliers were detected, the test was done with (i.e. the entire sample) and without the outlier.
206 Limb differences (UNJ vs INJ) in AT non-uniformity, displacement amplitude, maximum
207 and minimum displacement locations were compared using two-sided paired t-tests.

208

Achilles MG and LG subtendon representation

209 **Results**

210 Free tendon length below the SOL muscle insertion site was significantly longer in INJ
211 compared to UNJ with a mean difference (95%CI) of 1.6 cm (0.6-2.6 cm; $p=0.003$). The INJ
212 MG subtendon was also longer by 2.1 cm (1.5–2.7 cm; $p<0.01$), and LG by 1.9 cm (1.2–2.6
213 cm; $p<0.01$) than in UNJ. There were no statistically significant differences in stimulation
214 threshold or intensity between limb muscles or between limbs (Table 2).

215

216 Absolute displacement values and torque levels are reported in (Table 3). There was no
217 statistically significant difference in stimulation evoked torque levels between limbs in
218 response to stimulation of either muscle despite the stimulation inducing a significantly
219 higher mean displacement in both INJ in muscles compared to UNJ. The mean (SD)
220 magnitude of heel lift during electrically induced contractions was 0.04 mm (0.5) and 2.5 mm
221 (4.0) during voluntary contractions.

222

223 **Voluntary and stimulation-induced displacement patterns**

224 To explore the differences in displacement patterns, the absolute values of the 6 locations
225 were normalized to enable comparison between VOL and stimulation conditions (Figure 1).
226 Three-way repeated-measures ANOVA was performed to evaluate the effects of contraction
227 type, location and limb condition on tendon displacement. There was a significant two-way
228 interaction of contraction type*location on the tendon displacement ($F(10,978) = 3.7$,
229 $p<0.001$). Initial three-way analysis was followed by a two-way repeated-measures ANOVA
230 for the effect of contraction type*location on tendon displacement at the two levels of limb
231 condition and the location*limb condition on tendon displacement at each contraction type
232 level.

Achilles MG and LG subtendon representation

233 There was no significant location*limb condition interaction effect on tendon displacement at
234 each contraction type level. There was a significant contraction type*location interaction
235 effect on tendon displacement ($F(10-492) = 3.8, p < 0.001$) at the UNJ limb, while the
236 interaction effect was not significant for the INJ limb ($F(10-486) = 1.11, p = 0.353$). Simple
237 pairwise comparisons were done between the contraction types for the UNJ with a Bonferroni
238 adjustment applied. The analysis showed that the LG_{stim} displacement pattern was
239 significantly different to MG_{stim} ($p = 0.007$), and VOL ($p = 0.003$) (Figure 1). Individual
240 displacement patterns are shown in (Figure 2).

241 In UNJ, maximum displacement during MG_{stim} occurred most frequently in the three most
242 anterior locations, while during LG_{stim} , maximum displacement occurred most often in the
243 most anterior (6th) location (Figure 4). This pattern was also found in INJ, where the most
244 frequent locations of maximum displacement during MG_{stim} were in the anterior half of the
245 tendon (frequency of maximal displacement: 4th: 21.4%, 5th: 32.2% and 6th: 32.7%), while
246 during LG_{stim} , maximum displacement occurred in the 6th location in 48.2% of participants.
247 Minimum displacement was found in the most posterior location for the stimulation of both
248 muscles in both limbs. There was no statistically significant difference in maximum or
249 minimum displacement location between limbs.

250

251 **Tendon non-uniformity and displacement amplitude during electrical stimulation**

252 Tendon non-uniformity was higher in UNJ compared with INJ with a mean difference
253 (95%CI) of 0.11 mm (0.04 – 0.18mm, $p = 0.005$) during MG_{stim} , and 0.09 mm (0.03 – 1.42
254 mm, $p < 0.001$) during LG_{stim} (Figure 3). When non-uniformity was compared between
255 stimulated muscles in the same limb, there was no statistically significant difference in either
256 limb, with a mean difference (95%CI) of 0.016 mm (-0.06 – 0.09 mm) for UNJ and 0.003

Achilles MG and LG subtendon representation

257 mm (-0.05 – 0.05 mm) for INJ. One outlier was detected in LG_{stim} mean displacement group
258 (Higher range:1.9mm, outlier:2.7mm), when the whole sample was used there was no
259 significant difference in mean tendon displacement of the INJ between the contractions
260 induced when stimulating different muscles, with a mean difference (95%CI) of 0.28 mm (-
261 0.004 – 0.56 mm, p=0.053), however when the test was done without the outlier there was a
262 significant difference (95%CI) of mean difference of 0.34 mm (0.007 – 0.61 mm, p=0.015).
263 In the UNJ, there was a significant difference in the mean displacement depending on the
264 stimulated muscle, with a greater displacement when MG was stimulated (95%CI) of mean
265 difference of 0.22 mm (0.04 – 0.40 mm, p=0.016).

266

267 **Discussion**

268 In this study, we examined internal AT displacement patterns during voluntary and selective
269 transcutaneous stimulation of medial and lateral gastrocnemius to investigate differences
270 within the AT tissue displacement between INJ and UNJ limbs of patients after AT rupture
271 and to inspect the representative areas of subtendons. The lowest stimulation intensity that
272 induced a visible contraction was used to ensure selective activation of only the targeted
273 muscle. As hypothesized, displacement patterns during voluntary and electrically induced
274 contractions were different; the displacement pattern was significantly different during LG_{stim}
275 compared to VOL and MG_{stim} in both limbs. There was no statistically significant difference
276 when the displacement patterns were compared for each stimulated contraction between
277 limbs. Thus, with the assumption that the stimulation-induced force is primarily serially
278 transmitted to tendon fascicles, the subtendon organization does not seem to be altered in the
279 non-surgically treated limb of ATR patients. In UNJ, peak tendon displacement during
280 MG_{stim} tended to occur more posteriorly compared to VOL. Overall, the anterior half of the
281 AT underwent larger displacement than the superficial posterior part in all contraction
282 conditions.

283 Despite higher mean displacement in INJ, displacement was more uniform when compared to
284 UNJ during contractions induced by stimulating MG and LG. Tendon stiffness also seemed
285 to be lower in INJ, since ankle joint torque was similar during muscle stimulations in both
286 limbs, but the displacement was larger in INJ than in UNJ. However, this was not observed
287 for voluntary contractions in which tendon mean tendon displacement did not differ between
288 the limbs. Marked inter-individual differences were observed in internal tendon motion.
289 Thus, when investigating AT anatomical organization and internal force sharing, an
290 individualized approach might help to understand AT force sharing mechanisms and tendon
291 recovery from injury.

Achilles MG and LG subtendon representation

292 **Voluntary vs. stimulated contractions**

293 Internal tendon displacement patterns were different during LG_{stim} compared to VOL and
294 MG_{stim} in UNJ. In VOL, peak displacement was typically found in the two most anterior
295 locations. Voluntary contraction leads to disparate tissue displacement within the tendon due
296 to disproportionate activation of synergistic muscles and mechanical structural differences
297 between triceps surae muscles (7, 19). On the other hand, the low, stimulation-induced force
298 can be assumed to be mainly transmitted serially to the targeted muscle's subtendon (26).
299 Although lateral force transmission may occur (21), the main pathway of force is the stiffest
300 structure. Hence, the location of peak displacement in response to stimulation can be
301 considered to reveal the location of tendon fascicles within the cross-section of AT.

302 Displacement during LG_{stim} peaked in the anterior tendon, implying that the most anterior area
303 could be occupied by tendon fascicles arising from LG subtendon. In anatomical studies,
304 Pękala et al. (2017) and Edama et al. (2015) found that SOL occupied the anterior portion and
305 LG the lateral portion of the tendon at the level of the SOL muscle-tendon junction.
306 However, due to high torsion within AT, LG tendon fascicles are likely located anteriorly in
307 the more distal tendon (5, 6). Furthermore, in a recent study, three tendons were dissected,
308 and 3D computer aided models were constructed based on these tendons. In the model that
309 twisted the most, LG subtendon was found to completely occupy the anterior portion of the
310 distal AT (27). Therefore, anatomical studies are consistent with the present observations
311 regarding the location of the LG subtendon.

312 There was no difference between VOL and MG_{stim} displacement patterns (Figure 1).
313 However, during MG_{stim} displacement peaked around the 4th and 5th locations in UNJ,
314 indicating that fascicles originating from MG could be present in the mid-to-anterior part of
315 the tendon. Unlike in LG_{stim}, there was more individual variation in the location of peak
316 displacement in MG_{stim}. Due to individual differences in free tendon length, the superior-

Achilles MG and LG subtendon representation

317 inferior field of view may not have been consistent across subjects relative to tendon length.
318 When comparing these observations to the anatomical maps provided by previous cadavers
319 studies, natural anatomical variation may explain the observed heterogeneity in peak
320 displacement in response to MG_{stim} (5, 6, 27).

321 In addition to the anatomical origin, the observed peak displacement locations may have been
322 affected by lateral force transmission between different subtendons within the AT. Each
323 subtendon transmits the force from a single muscle belly but not fully independently, and
324 force could be laterally transmitted between triceps surae muscle bellies or even subtendons
325 (28, 29). AT force and subsequent displacement might be distributed unevenly with a bias
326 toward the SOL subtendon since SOL subtendon fascicles have been found previously to be
327 compliant in rats (29) and in human cadavers (30) although contradictory results have also
328 been reported (27). This raises questions about the forces transmitted through connective
329 tissue or inter-fascicular matrix, which could be crucial for force transmission mechanisms
330 and inter-fascicular gliding within the tendon (10, 31).

331 In summary, during VOL and electrically induced contractions of MG and LG, minimum
332 displacement always occurred in the posterior tendon and maximum displacement in the mid-
333 to-anterior tendon. The same observation was made in a recent study where SOL and MG
334 were electrically stimulated (32), and the tendon was split into two halves for analysis
335 purposes; the anterior half always displaced the most in response to MG and SOL
336 stimulations in different ankle positions. This is consistent with the observations made in this
337 study as we found that the mid-to-deep part of the tendon displaced most when MG was
338 stimulated and the deep part when LG was stimulated. The difference between the two
339 studies is in the interpretation of the data in regard to which subtendon are presented in the
340 deep part of the tendon. In Lehr et al. the tendon was split in consideration to the function-
341 structure relationship (11, 12), and the authors interpreted a larger non-uniformity and

Achilles MG and LG subtendon representation

342 displacement in the representative part of the stimulated muscle tendon when SOL was
343 stimulated compared to MG as an evidence of consistency with the anatomical function-
344 structure consideration (32). We relied on the principle that the main pathway of force is the
345 stiffest structure. Thus, when a muscle is selectively stimulated the arising regional tendon
346 displacement can inform us about regions of the tendon corresponding to fascicles arising
347 from different triceps surae muscles. Based on the beforementioned we found that the
348 gastrocnemius subtendons are located in the mid-to-anterior part of the tendon, and that LG is
349 probably located most anteriorly.

350

351 Gastrocnemius and soleus have different functional roles, despite having a common distal
352 tendon and working synergistically as ankle plantar flexors (33, 34). It has been suggested
353 that in order to perform their differing functional roles, these muscles rely on the ability of
354 the subtendons to displace relative to each other (35, 36). It is of interest to investigate if
355 normal subtendon organization can be restored after ATR as this most likely is a prerequisite
356 for restoring normal Achilles tendon and triceps surae function including the functional
357 independence of the muscles. If tendon fascicles that were originally part of different
358 subtendons would merge during the healing process this could result in reduced capacity for
359 relative movement between subtendons and disruption of the normal function of the Achilles
360 tendon. In fact, ATR followed by surgical reconstruction has been shown to reduce non-
361 uniform tendon motion observed using speckle tracking (8, 37). Our tendon displacement
362 data (Figure 2) and our previous report (14) suggest that there are considerable individual
363 variations in the subtendon organization in both ruptured and un-injured tendons. This
364 signifies the importance of an individualized assessment and interpretation of the subtendon
365 organization and function after ATR.

366

Achilles MG and LG subtendon representation

367 **Tendon non-uniformity and displacement amplitude during electrical stimulation**

368 Consistent with previous studies, we found a more uniform displacement pattern in INJ
369 compared to the contralateral tendon 1-year post rupture (13, 14), suggesting impaired sliding
370 within the injured tendon. Limited inter-fascicular sliding might be a result of interfascicular
371 matrix adhesions caused by the rupture (10). Mean displacement in INJ was higher than in
372 UNJ. As the same amount of torque was produced during stimulation, this result suggests
373 lower stiffness in INJ. However, this differs from our previous results, where we reported that
374 stiffness of the entire MG tendon during isometric voluntary contraction was similar between
375 injured and un-injured tendons 1-year post rupture (38). This would suggest that in the free
376 distal AT, mechanical properties (stiffness) may be altered locally and manifest themselves at
377 low force levels, while globally stiffness seems to be similar between limbs. This discrepancy
378 between our observations could indicate an extension of the toe region, or slackness of the
379 tendon in the INJ limb while the linear region of the force-displacement curve would be
380 similar between the limbs. A similar phenomenon of an extended range of tendon strain at
381 low stresses has been reported previously after 4 weeks of limb unloading by suspension
382 (39).

383

384 We found no differences between limbs (UNJ vs INJ) in the locations of maximum or
385 minimum displacement when stimulating either MG or LG, consistent with our previous
386 findings during voluntary contractions (14). Furthermore, in the three contraction types the
387 displacement patterns were similar when compared between limbs. Thus, the anatomical
388 subtendon organization does not seem to be altered after a rupture in non-surgically treated
389 tendons.

Achilles MG and LG subtendon representation

390 **Limitations**

391 There are several limitations of this study. First, the nature of two-dimensional imaging may
392 not fully capture the complex three-dimensional behaviour of the triceps surae subtendons,
393 which could lead to errors when estimating AT tissue displacement. The speckle tracking
394 algorithm uses a low order polynomial fit to regularize displacement (25). This may reduce
395 variation in displacement between the six locations across the tendon. However, filtering has
396 been deemed necessary to reduce noise and erroneous estimates (40), and was applied here in
397 the same manner as in previous studies (32). Furthermore, it should be noted that LG muscle
398 has different compartments that are innervated by two main nerves and numerous sub-
399 branches (41), so stimulation might activate different branches of the muscle causing more
400 variability to the displacement pattern. Furthermore, selective activation of LG might stiffen
401 the connective tissue between SOL and LG, facilitating force transmission (29). Thus, the
402 representation of LG or MG subtendon that we observed within the AT may have been
403 influenced by lateral force transmission at the level of the muscle or tendon. However, this
404 effect was likely minimal since it has been shown that lateral force sharing within the human
405 Achilles tendon is small at low forces (42).

406 **Conclusion**

407 To conclude, Achilles tendon displacement patterns were different in response to selective
408 stimulation of LG compared to MG stimulation or voluntary contraction. Our results suggest
409 that when imaged from a mid-sagittal view, the gastrocnemius subtendons are located in the
410 mid-to-anterior part of the tendon, and that LG is probably located most anteriorly. Previous
411 anatomical studies support these results, but more investigations are needed since results in
412 the literature are inconsistent. The stimulation method could allow for a more individualized
413 approach for investigation of tendon organization, that might help to better understand the
414 complex mechanics and triceps surae subtendon representations within the Achilles tendon.

Achilles MG and LG subtendon representation

415 We found no evidence that non-surgical treatment of ATR alters the displacement pattern
416 within the tendon suggesting that non-surgical treatment may preserve the normal subtendon
417 organization. However, differences in displacement amplitude and non-uniformity of the
418 tendon displacement were present between the limbs in electrically stimulated conditions.
419 These findings suggest an extended toe region of the tendon force-displacement curve after
420 ATR and potential adhesions preventing non-uniform displacements.

Achilles MG and LG subtendon representation

421 **Grants**

422 This study was funded by Academy of Finland grant #323168, UNderstanding REStoration
423 of Achilles Tendon function after rupture (UNRESAT), and in part by Academy of Finland
424 grant #332915.

425

426 **Disclosure**

427 The authors declare that they have no competing interests.

428 **References**

- 429 1. **Fukashiro S, Hay DC, Nagano A.** Biomechanical Behavior of Muscle-Tendon Complex during
430 Dynamic Human Movements. *J Appl Biomech* 22: 131–147, 2006. doi: 10.1123/jab.22.2.131.
- 431 2. **Roberts TJ, Azizi E.** Flexible mechanisms: the diverse roles of biological springs in vertebrate
432 movement. *J Exp Biol* 214: 353–361, 2011.
- 433 3. **Ganestam A, Kallemose T, Troelsen A, Barfod KW.** Increasing incidence of acute Achilles
434 tendon rupture and a noticeable decline in surgical treatment from 1994 to 2013. A nationwide
435 registry study of 33,160 patients. *Knee Surg Sports Traumatol Arthrosc* 24: 3730–3737, 2016.
- 436 4. **Lantto I, Heikkinen J, Flinkkilä T, Ohtonen P, Leppilähti J.** Epidemiology of Achilles tendon
437 ruptures: increasing incidence over a 33-year period. *Scand J Med Sci Sports* 25: e133-138,
438 2015. doi: 10.1111/sms.12253.
- 439 5. **Edama M, Kubo M, Onishi H, Takabayashi T, Inai T, Yokoyama E, Hiroshi W, Satoshi N,
440 Kageyama I.** The twisted structure of the human Achilles tendon. *Scand J Med Sci Sports* 25:
441 e497–e503, 2015.
- 442 6. **Pełkala PA, Henry BM, Ochoła A, Kopacz P, Tatoń G, Młyniec A, Walocha JA, Tomaszewski KA.**
443 The twisted structure of the Achilles tendon unraveled: A detailed quantitative and qualitative
444 anatomical investigation. *Scand J Med Sci Sports* 27: 1705–1715, 2017. doi:
445 10.1111/sms.12835.
- 446 7. **Bojsen-Møller J, Magnusson SP.** Heterogeneous Loading of the Human Achilles Tendon In
447 Vivo. *Exerc Sport Sci Rev* 43: 190–197, 2015. doi: 10.1249/JES.0000000000000062.
- 448 8. **Beyer R, Agergaard A-S, Magnusson SP, Svensson RB.** Speckle tracking in healthy and
449 surgically repaired human Achilles tendons at different knee angles—A validation using
450 implanted tantalum beads. *Transl Sports Med* 1: 79–88, 2018. doi: 10.1002/tsm2.19.
- 451 9. **Slane LC, Thelen DG.** Non-uniform displacements within the Achilles tendon observed during
452 passive and eccentric loading. *J Biomech* 47: 2831–2835, 2014.
- 453 10. **Thorpe CT, Udeze CP, Birch HL, Clegg PD, Screen HR.** Capacity for sliding between tendon
454 fascicles decreases with ageing in injury prone equine tendons: a possible mechanism for age-
455 related tendinopathy. *Eur Cell Mater* 25, 2013.
- 456 11. **Clark WH, Franz JR.** Do triceps surae muscle dynamics govern non-uniform Achilles tendon
457 deformations? *PeerJ* 6: e5182, 2018. doi: 10.7717/peerj.5182.
- 458 12. **Stenroth L, Thelen D, Franz J.** Biplanar ultrasound investigation of in vivo Achilles tendon
459 displacement non-uniformity. *Transl Sports Med* 2: 73–81, 2019.
- 460 13. **Fröberg Å, Cissé A-S, Larsson M, Mårtensson M, Peolsson M, Movin T, Arndt A.** Altered
461 patterns of displacement within the Achilles tendon following surgical repair. *Knee Surg Sports
462 Traumatol Arthrosc* 25: 1857–1865, 2017.
- 463 14. **Khair RM, Stenroth L, Péter A, Cronin NJ, Reito A, Paloneva J, Finni T.** Non-uniform
464 displacement within ruptured Achilles tendon during isometric contraction. *Scand J Med Sci
465 Sports* 31: 1069–1077, 2021. doi: <https://doi.org/10.1111/sms.13925>.

Achilles MG and LG subtendon representation

- 466 15. **Peng WC, Chao YH, Fu ASN, Fong SSM, Rolf C, Chiang H, Chen S, Wang HK.** Muscular
467 Morphomechanical Characteristics After an Achilles Repair. *Foot Ankle Int* 40: 568–577, 2019.
468 doi: 10.1177/1071100718822537.
- 469 16. **Svensson RB, Couppé C, Agergaard A-S, Ohrhammar Josefsen C, Jensen MH, Barfod KW,**
470 **Nybing JD, Hansen P, Krogsgaard M, Magnusson SP.** Persistent functional loss following
471 ruptured Achilles tendon is associated with reduced gastrocnemius muscle fascicle length,
472 elongated gastrocnemius and soleus tendon, and reduced muscle cross-sectional area. *Transl*
473 *SPORTS Med* 2: 316–324, 2019. doi: 10.1002/tsm2.103.
- 474 17. **Hug F, Del Vecchio A, Avrillon S, Farina D, Tucker K.** Muscles from the same muscle group do
475 not necessarily share common drive: evidence from the human triceps surae. *J Appl Physiol*
476 130: 342–354, 2021. doi: 10.1152/jappphysiol.00635.2020.
- 477 18. **Masood T, Bojsen-Møller J, Kalliokoski KK, Kirjavainen A, Äärimaa V, Peter Magnusson S,**
478 **Finni T.** Differential contributions of ankle plantarflexors during submaximal isometric muscle
479 action: A PET and EMG study. *J Electromyogr Kinesiol* 24: 367–374, 2014. doi:
480 10.1016/j.jelekin.2014.03.002.
- 481 19. **Albracht K, Arampatzis A, Baltzopoulos V.** Assessment of muscle volume and physiological
482 cross-sectional area of the human triceps surae muscle in vivo. *J Biomech* 41: 2211–2218, 2008.
483 doi: 10.1016/j.jbiomech.2008.04.020.
- 484 20. **Bojsen-Møller J, Schwartz S, Kalliokoski KK, Finni T, Magnusson SP.** Intermuscular force
485 transmission between human plantarflexor muscles in vivo. *J Appl Physiol* 109: 1608–1618,
486 2010. doi: 10.1152/jappphysiol.01381.2009.
- 487 21. **Finni T, Cronin NJ, Mayfield D, Lichtwark GA, Cresswell AG.** Effects of muscle activation on
488 shear between human soleus and gastrocnemius muscles. *Scand J Med Sci Sports* 27: 26–34,
489 2017.
- 490 22. **Reito A, Logren H-L, Ahonen K, Nurmi H, Paloneva J.** Risk factors for failed nonoperative
491 treatment and rerupture in acute Achilles tendon rupture. *Foot Ankle Int* 39: 694–703, 2018.
- 492 23. **Barfod KW, Riecke AF, Boesen A, Hansen P, Maier JF, Døssing S, Troelsen A.** Validation of a
493 novel ultrasound measurement of Achilles tendon length. *Knee Surg Sports Traumatol Arthrosc*
494 23: 3398–3406, 2015.
- 495 24. **Hopkins WG.** Measures of Reliability in Sports Medicine and Science. *Sports Med* 30: 1–15,
496 2000. doi: 10.2165/00007256-200030010-00001.
- 497 25. **Slane LC, Thelen DG.** The use of 2D ultrasound elastography for measuring tendon motion and
498 strain. *J Biomech* 47: 750–754, 2014.
- 499 26. **Tian M, Herbert RD, Hoang P, Gandevia SC, Bilston LE.** Myofascial force transmission between
500 the human soleus and gastrocnemius muscles during passive knee motion. *J Appl Physiol* 113:
501 517–523, 2012. doi: 10.1152/jappphysiol.00111.2012.
- 502 27. **Yin N-H, Fromme P, McCarthy I, Birch HL.** Individual variation in Achilles tendon morphology
503 and geometry changes susceptibility to injury. *eLife* 10: e63204, 2021. doi:
504 10.7554/eLife.63204.

Achilles MG and LG subtendon representation

- 505 28. **Bernabei M, van Dieën JH, Baan GC, Maas H.** Significant mechanical interactions at
506 physiological lengths and relative positions of rat plantar flexors. *J Appl Physiol* 118: 427–436,
507 2015. doi: 10.1152/jappphysiol.00703.2014.
- 508 29. **Finni T, Bernabei M, Baan GC, Noort W, Tijs C, Maas H.** Non-uniform displacement and strain
509 between the soleus and gastrocnemius subtendons of rat Achilles tendon. *Scand J Med Sci*
510 *Sports* 28: 1009–1017, 2018.
- 511 30. **Ekiert M, Tomaszewski KA, Mlyniec A.** The differences in viscoelastic properties of subtendons
512 result from the anatomical tripartite structure of human Achilles tendon - ex vivo experimental
513 study and modeling. *Acta Biomater* 125: 138–153, 2021. doi: 10.1016/j.actbio.2021.02.041.
- 514 31. **Thorpe CT, Godinho MS, Riley GP, Birch HL, Clegg PD, Screen HR.** The interfascicular matrix
515 enables fascicle sliding and recovery in tendon, and behaves more elastically in energy storing
516 tendons. *J Mech Behav Biomed Mater* 52: 85–94, 2015.
- 517 32. **Lehr NL, Clark WH, Lewek MD, Franz JR.** The effects of triceps surae muscle stimulation on
518 localized Achilles subtendon tissue displacements. *J Exp Biol* 224, 2021. doi:
519 10.1242/jeb.242135.
- 520 33. **Francis CA, Lenz AL, Lenhart RL, Thelen DG.** The modulation of forward propulsion, vertical
521 support, and center of pressure by the plantarflexors during human walking. *Gait Posture* 38:
522 993–997, 2013. doi: 10.1016/j.gaitpost.2013.05.009.
- 523 34. **Lenhart RL, Francis CA, Lenz AL, Thelen DG.** Empirical evaluation of gastrocnemius and soleus
524 function during walking. *J Biomech* 47: 2969–2974, 2014. doi: 10.1016/j.jbiomech.2014.07.007.
- 525 35. **Clark WH, Franz JR.** Age-related changes to triceps surae muscle-subtendon interaction
526 dynamics during walking. *Sci Rep* 11: 21264, 2021. doi: 10.1038/s41598-021-00451-y.
- 527 36. **Franz JR, Thelen DG.** Imaging and simulation of Achilles tendon dynamics: implications for
528 walking performance in the elderly. *J Biomech* 49: 1403–1410, 2016. doi:
529 10.1016/j.jbiomech.2016.04.032.
- 530 37. **Fröberg Å, Cissé A-S, Larsson M, Mårtensson M, Peolsson M, Movin T, Arndt A.** Altered
531 patterns of displacement within the Achilles tendon following surgical repair. *Knee Surg Sports*
532 *Traumatol Arthrosc* 25: 1857–1865, 2017. doi: 10.1007/s00167-016-4394-5.
- 533 38. **Khair RM, Stenroth L, Cronin NJ, Reito A, Paloneva J, Finni T.** Muscle-tendon
534 morphomechanical properties of non-surgically treated Achilles tendon 1-year post-rupture.
535 *Clin Biomech* 92: 105568, 2022. doi: 10.1016/j.clinbiomech.2021.105568.
- 536 39. **Shin D, Finni T, Ahn S, Hodgson JA, Lee H-D, Edgerton VR, Sinha S.** Effect of chronic unloading
537 and rehabilitation on human Achilles tendon properties: a velocity-encoded phase-contrast
538 MRI study. *J Appl Physiol Bethesda Md* 1985 105: 1179–1186, 2008. doi:
539 10.1152/jappphysiol.90699.2008.
- 540 40. **Svensson RB, Slane LC, Magnusson SP, Bogaerts S.** Ultrasound-based speckle-tracking in
541 tendons: a critical analysis for the technician and the clinician. *J Appl Physiol* 130: 445–456,
542 2021. doi: 10.1152/jappphysiol.00654.2020.

Achilles MG and LG subtendon representation

543 41. **Segal RL, Wolf SL, DeCamp MJ, Chopp MT, English AW.** Anatomical Partitioning of Three
544 Multiarticular Human Muscles. *Cells Tissues Organs* 142: 261–266, 1991. doi:
545 10.1159/000147199.

546 42. **Haraldsson BT, Aagaard P, Qvortrup K, Bojsen-Moller J, Krogsgaard M, Koskinen S, Kjaer M,**
547 **Magnusson SP.** Lateral force transmission between human tendon fascicles. *Matrix Biol* 27:
548 86–95, 2008. doi: 10.1016/j.matbio.2007.09.001.

549

550

Achilles MG and LG subtendon representation

551 **Figure captions:**

552 Figure 1. Mean normalized displacement patterns \pm SD during voluntary and selective electrically
553 induced contractions of the medial (MG) and lateral (LG) gastrocnemius muscles in the un-injured
554 (left) and injured limb (right). Graphs represent group means Individual patterns in the un-injured
555 limb are shown in Figure 2. * Difference between the contraction types ($p < 0.05$).

556

557 Figure 2. Normalized displacement patterns in the un-injured limb. Left: Raw data points for each
558 participant during voluntary and selective electrical stimulation across the 6 locations of the Achilles
559 tendon. Right: Box plots of means and SD for each location (1-6 respectively).
560 * Difference between the contraction types ($p < 0.05$).

561

562 Figure 3. Tendon displacement (mm) of the whole sample during gastrocnemius muscle stimulation at
563 each of the six locations across the tendon width. The values are expressed as mean \pm SD.

564

565 Figure 4. Distribution of peak displacement locations across the tendon in the sagittal view when
566 medial gastrocnemius (MG, upper) or lateral gastrocnemius (LG, lower) was selectively stimulated in
567 the un-injured limb.

568

Table 1. Patient characteristics, free Achilles tendon length, and medial and lateral gastrocnemius subtendon lengths (mean \pm SD).

Participants (N=28)		
Age (years)	42.4 \pm 9.3	
Height (m)	1.76 \pm 0.08	
Body mass (Kg)	82.5 \pm 12.2	
Limb condition	Un-injured	Injured
Free tendon length (cm)	8.79 \pm 3.47	10.36 \pm 3.71
MG subtendon length (cm)	18.90 \pm 1.92	20.99 \pm 2.20
LG subtendon length (cm)	21.59 \pm 1.60	23.51 \pm 1.99

Table 2. Descriptive data of motor thresholds, selective electrical stimulation-induced contractions intensities of medial (MG) and lateral (LG) gastrocnemius muscles in the un-injured and injured limbs, and comparisons between limbs and muscles.



	Injured		Un-injured		<i>P</i>-values comparing stimulations			
	MG	LG	MG	LG	between limbs		between muscles	
					MG	LG	INJ	UNJ
Stimulation intensity mA (SD)	20.36 (9.26)	18.48 (5.63)	17.75 (9.26)	16.07 (7.37)	0.164	0.155	0.413	0.097
Threshold mA (SD)	15.50 (9.75)	15.25 (4.76)	14.07 (7.98)	19.75 (7.71)	0.513	0.408	0.634	0.210

P-values using un-adjusted pairwise t-test.

Table 3. Descriptive data of mean displacement, non-uniformity, and absolute torque of electrically induced and voluntary contractions.

	Injured			Un-injured		
	MG	LG	VOL	MG	LG	VOL
Mean displacement mm (SD)	0.93 (0.65)	0.65 (0.57)	3.52 (1.71)	0.61 (0.48)	0.39 (0.27)	3.63 (1.18)
Tendon non-uniformity mm (SD)	0.14 (0.11)	0.15 (0.12)	0.85 (0.79)	0.25 (0.23)	0.24 (0.17)	1.48 (1.04)
Torque Nm (SD)	5.18 (2.98)	3.24 (2.72)	57.98 (16.30)	5.56 (4.05)	3.67 (2.53)	57.78 (16.23)


Normalized displacement





Contraction type


- LG_{stim}
- MG_{stim}
- VOL

Normalized displacement






MG




Post./Super



ANT./Deep

LG



Post./Super




ANT./Deep

In vivo localised gastrocnemius subtendon representation within the healthy and ruptured human Achilles tendon

N=28


METHODS

Selective electrical stimulation & Voluntary plantarflexion



RESULTS

Frequency of maximal displacement locations during selective electrical stimulations



MG subtendon → anterior mid region
LG subtendon → most anterior region

CONCLUSION

When comparing healthy and ruptured tendons one-year post rupture, tendon internal displacement patterns were not different, but absolute displacement amplitude and non-uniformity did differ. **LG** subtendon was found in the most anterior region adjacent to **MG** in both healthy and non-surgically treated tendons.

