

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Contardo Ayala, Ana María; Parker, Kate; Mazzoli, Emiliano; Lander, Natalie; Ridgers, Nicola D.; Timperio, Anna; Lubans, David R.; Abbott, Gavin; Koorts, Harriet; Salmon, Jo

Title: Effectiveness of Intervention Strategies to Increase Adolescents' Physical Activity and Reduce Sedentary Time in Secondary School Settings, Including Factors Related to Implementation : A Systematic Review and Meta-Analysis

Year: 2024

Version: Published version

Copyright: © The Author(s) 2024

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:


Contardo Ayala, A. M., Parker, K., Mazzoli, E., Lander, N., Ridgers, N. D., Timperio, A., Lubans, D. R., Abbott, G., Koorts, H., & Salmon, J. (2024). Effectiveness of Intervention Strategies to Increase Adolescents' Physical Activity and Reduce Sedentary Time in Secondary School Settings, Including Factors Related to Implementation : A Systematic Review and Meta-Analysis. *Sports Medicine - Open*, 10, Article 25. <https://doi.org/10.1186/s40798-024-00688-7>

SYSTEMATIC REVIEW

Open Access



Effectiveness of Intervention Strategies to Increase Adolescents' Physical Activity and Reduce Sedentary Time in Secondary School Settings, Including Factors Related to Implementation: A Systematic Review and Meta-Analysis

Ana María Contardo Ayala^{1,4*} , Kate Parker^{1,4}, Emiliano Mazzoli^{1,5}, Natalie Lander^{1,4}, Nicola D. Ridgers^{1,2}, Anna Timperio^{1,4}, David R. Lubans^{3,6,7}, Gavin Abbott^{1,4}, Harriet Koorts^{1,4} and Jo Salmon^{1,4}

Abstract

Background Globally, just one in five adolescents meet physical activity guidelines and three-quarters of the school day is spent sitting. It is unclear which types of school-based interventions strategies increase physical activity and reduce sedentary time among adolescents, or how these interventions are implemented influences their effectiveness.

Objective The three aims of our systematic review were to (a) identify intervention strategies used within secondary school settings to improve students' movement behaviours throughout school-based initiatives, delivered at or by the school; (b) determine the overall effect of the interventions (meta-analysis) on physical activity (all intensities), sedentary time, cognitive/academic, physical health and/or psychological outcomes; and (c) describe factors related to intervention implementation.

Methods Searches were conducted in MEDLINE complete, EMBASE, CINAHL, SPORTDiscus, APA PsycINFO, and ERIC in January 2023 for studies that (a) included high school-aged adolescents; (b) involved a school-based intervention to increase physical activity and/or decrease sedentary time; and (c) were published in English. Reported effects were pooled in meta-analyses where sufficient data were obtained.

Results Eighty-five articles, representing 61 interventions, met the inclusion criteria, with 23 unique intervention strategies used. Interventions that involved whole-school approaches (i.e., physical activity sessions, environmental modifications, teacher training, peer support and/or educational resources) were favourably associated with most of the outcomes. The meta-analyses showed: (a) non-significant effects for sedentary time (Standardized mean

*Correspondence:

Ana María Contardo Ayala
a.contardoayala@deakin.edu.au

Full list of author information is available at the end of the article

difference [SMD] = -0.02; 95%CI, -0.14, 0.11), physical activity at all intensities (light: SMD= -0.01; 95%CI, -0.08, 0.05; moderate: SMD= 0.06; 95%CI, -0.09, 0.22; vigorous: SMD= 0.08; 95%CI, -0.02, 0.18; moderate-to-vigorous: SMD= 0.05; 95%CI, -0.01, 0.12) and waist circumference (SMD= 0.09; 95%CI, -0.03, 0.21), and (b) a small statistically significant decrease in body mass index (SMD= -0.09, 95%CI -0.16, -0.0). Factors related to intervention implementation were reported in 51% of the articles.

Conclusion While some intervention approaches demonstrated promise, small or null effects were found in meta-analyses. Future school-based interventions should utilize a whole-school approach designed to increase adolescents' activity across the day. Consistent reporting of implementation will increase understanding of how interventions are adopted, implemented and sustained.

Registration PROSPERO (CRD42020169988).

Key Points

- School-based interventions with adolescents have had null effects on increasing physical activity and reducing sedentary time.
- The impact on physical and psychological outcomes has been poorly reported among secondary school students.
- This review included articles with all experimental study designs, any intervention length, and that reported a variety of outcomes among adolescents in secondary schools (11–18 years old).
- Although meta-analyses showed that these interventions did not increase physical activity or reduce sedentary time, studies that quantify adiposity markers showed a small decrease in body mass index.
- The review identified that additional physical activity sessions, environmental modifications, teacher training, peer support, educational resources and/or active lesson strategies were associated with greater benefits.
- Fidelity and participant responses were the most commonly reported implementation factors, adaptation was the least commonly reported; however, factors associated with effective implementation were not a focus of the included studies.

Keywords Adolescents, Physical activity, Sedentary behaviour, School-based interventions, Implementation

Background

Adequate physical activity has a multitude of benefits for adolescents including a reduced risk of developing adverse psychological [1] and physical health conditions [2]. Moreover, physical activity is positively associated with academic outcomes, including cognitive skills (e.g., executive functioning, memory) [3], attitude (e.g., motivation, self-concept) [4], academic behaviour (e.g., on-task time, organization) [5], and academic achievement (e.g., standardized test scores) [6]. Conversely, excess sedentary time particularly recreational screen-time, during adolescence has negative implications for psychological [1] and physical health [7]. It is currently recommended that adolescents engage in at least an average of 60 min per day of moderate- to vigorous-intensity physical activity (MVPA) and limit the amount of time they spend sedentary [8].

Approximately 81% of adolescents globally do not meet physical activity guidelines [9]. Additionally, adolescents spend an average of nine hours per day sitting, which includes three hours per day engaged in sedentary screen-time [10]. Secondary (middle and high) school students can spend up to 75% of their class time sedentary; often accumulated in long, unbroken bouts of sitting [11]. Although secondary schools are typically required to provide regular physical education (PE) classes, just

36% of this time is spent in MVPA [12]. Moreover, scheduled PE time declines across secondary school years and is not always compulsory in the upper secondary or high school years [13]. There are many other opportunities for students to engage in movement behaviours (i.e., increase physical activity and/or reduce sedentary time) throughout the day, including recess and lunch breaks, and during, between and after lessons [14].

Several reviews have summarised school-based movement behaviour initiatives in the secondary school context, concluding that these have been largely ineffective (null to small positive effects) [15–23]. However, the majority have reported intervention effects on MVPA [15, 16, 18, 21], or sedentary time [23], included just PE interventions (already contributing to school-hours physical activity) [16, 19, 21], focused on older adolescents [17] or girls only [22], restricted inclusion criteria to specific intervention designs (only RCTs) [17, 18] and study lengths [15, 18], or focused on low-middle income countries [20]. Meta-analyses were performed in only five reviews [15, 16, 18, 21, 23] that described the impact of school-based interventions on physical activity (small or null effects) [16, 18, 21], and just two analyzed the intervention components used [15, 16]. None have reported the factors crucial for intervention implementation effectiveness.

To our knowledge, there is currently no synthesis of non-PE interventions delivered at or by schools to improve movement behaviours across the school day, and existing reviews have generally not reported outcomes beyond movement behaviours, such as cognitive and academic outcomes, physical health, and/or psychological outcomes. The three aims of our systematic review were to (a) identify intervention strategies used within secondary school settings to improve students' movement behaviours throughout school-based initiatives, delivered at or by the school; (b) determine the overall effect of the interventions (meta-analysis) on physical activity (all intensities), sedentary time, cognitive/academic, physical health and/or psychological outcomes; and (c) describe factors related to intervention implementation.

Methods

Protocol and Registration

This review was registered with PROSPERO (CRD42020169988) and followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Statement guidelines (PRISMA). Following the PROSPERO registration, the eligibility criteria were expanded to include strategies targeting school-related work (i.e. homework) and a third aim was added: to describe factors related to intervention implementation.

Eligibility Criteria and Search Strategies

Eligibility criteria and search strategy were modelled on the Participants, Interventions, Comparisons, Outcomes, and Study design (PICOS) framework. Studies were included if they included: (a) Secondary/middle/high school-age adolescents (>11 years and <18 years); (b) Interventions delivered in the school setting (i.e., during class, recess and lunch time) and/or related to schoolwork (e.g., homework); (c) Strategy/ies to increase physical activity of any intensity (i.e., light, moderate, vigorous or moderate to vigorous) and/or decrease sedentary behaviour; (d) Any outcomes (for movement behaviours measures, have to report an effect on daily and/or school-based physical activity and sedentary behaviour); (e) Any study design (e.g., randomized controlled trials [individual and cluster], controlled trials, pre-post studies design, quasi-experimental studies), and with any comparison (e.g., pre-post intervention comparison) or control groups (e.g., non-exposed control/comparison groups); (f) written in English. Studies were excluded from this review if they included: (a) Target population was adolescents with special needs; primary (elementary), middle and secondary school studies were combined with results reported together; participants were in year 6 only (considered a primary school year in some countries); (b) Before- or after-school hours programs; (c) Intervention targeted PE lessons, before- or

after-school hours, active travel, or educational programs (e.g., non-active lifestyle/health lessons) in isolation; (d) Non-experimental studies (e.g., cross-sectional and case studies); and (e) Findings were only reported in abstracts (including poster abstracts), conference proceedings, dissertations, commentaries, editorials, review articles, and letters. The search strategy is reported in Supplementary Table 1 (Search strategy used [i.e., EBSCO and EMBASE]).

Information Sources

A systematic search was conducted using six online databases: MEDLINE complete, Embase, CINAHL, SPORT-Discus, APA PsycINFO, and ERIC. Further articles were identified via forwards and backward citation tracking of included articles and relevant systematic reviews. Peer-reviewed articles between January 2000 and January 2023 were considered for inclusion, as most school-based movement behaviour interventions have been conducted in the last ~20 years. Reference lists of the included articles were also screened to identify additional eligible interventions.

Study Selection

All search results were exported into a reference manager (Endnote x9, Clarivate analytics) and duplicates were removed. Titles and abstracts were exported to Covidence (Melbourne, Australia). Two authors (KP, AMC) screened all titles and abstracts; discussed discrepancies and came to a consensus for inclusion. Both authors reviewed the full text and discussed full-text discrepancies. Any disagreements were solved in a meeting involving four authors (AMC, KP, AT, JS).

Data Collection Process

After identifying published articles, the authors identified whether they constituted individual studies or multiple articles from the same study. Hereafter, "articles" refers to the count of papers identified, while "studies" pertains to the number of distinct research studies/interventions represented by these articles. Data extraction was performed by two authors (AMC and KP), and all articles were cross-checked for accuracy by an additional author (EM). Data extracted and quantified included study and participant characteristics, intervention strategies and intervention effects. Intervention outcomes were classified into five categories: movement behaviours (e.g., physical activity, sedentary time, energy expenditure), cognitive and academic outcomes (e.g., working memory, on task behaviour), physical health (e.g., fitness, obesity, and musculoskeletal health), and psychological outcomes. Data extraction of factors relevant to the implementation of the programs (performed by AMC and KP;




Fig. 1 Study selection process (PRISMA – 2020 version). Legend: IV, intervention; PA, physical activity; SB, sedentary behaviour

checked for accuracy by NL) was guided by the United Kingdom Medical Research Council Process Evaluation of Complex Interventions [24], which includes the following three components: *Implementation* (implementation process, fidelity, dose, adaptations, reach), *Mechanism of impact* (participants' responses, mediators, unanticipated pathways/consequences, and *Context* (contextual factors which affect/shape intervention). Descriptive analysis was used to describe studies, participants' characteristics and intervention components and effects on physical activity (all intensities), sedentary time, cognitive/academic, physical health, and/or psychological outcomes.

Methodological Quality

Quality assessment screening was completed by three authors (AMC, KP, NL), in duplicate, using an adapted version of the Effective Public Health Practice Project tool for quantitative studies [25]. The three authors discussed discrepancies and came to a consensus for the scores. The components assessed were selection bias (sample representation), study design (e.g., RCT), confounders (e.g., control for baseline differences), blinding (researcher and participant), data collection methods (e.g., validity and reliability of the assessment tool used), withdrawals and dropouts (e.g., report of number of

withdrawals and reason), intervention integrity (e.g., percentage of participants that received the intervention), and statistical analysis (e.g., use of intention to treat approach) [26]. Individual criteria for methodological quality scores and an overall quality score were created and rated as strong (no weak rating), moderate (one weak rating) or weak (>1 weak rating) [25]. Inter-rater reliability for assessment of study quality between authors was moderate (Cohen's Kappa=0.6, $p < 0.001$).

Statistical Analysis

Descriptive analysis (number and/or percentage) was used to summarise studies and participant characteristics, intervention strategies and the effect of these strategies on physical activity (all intensities), sedentary time, cognitive/academic, physical health and/or psychological outcomes, and factors related to intervention implementation. In addition, meta-analyses were conducted using Stata v18 when at least three articles reported intervention effects on outcomes with a comparable device (e.g., actigraphy), same outcome (e.g. daily MPA, and/or measurement scales/unit (kg/m^2 , cm), provided complete data for pre- and post- intervention measurements, and were randomised controlled trials. Self-reported movement behaviour outcomes were excluded from the meta-analyses. Consequently, meta-analyses included articles

reporting on comparable: (a) device-measured movement behaviours (i.e. sedentary time, LPA, MPA, VPA, MVPA assessed during the whole day); and (b) body mass index and waist circumference (measured with standardized protocols). The sample size, mean difference, and standard error/confidence interval between intervention and control groups were entered into Microsoft® Excel and effect sizes were calculated (due to the varying units for a given outcome, standardized mean differences were calculated and used in meta-analysis). Reported immediate intervention effects were used for these meta-analyses and, in case of incomplete reporting, the corresponding authors were contacted to request additional information. For cluster randomized controlled trials, the point estimate and the 95% confidence interval intervention-reported data from their intervention effect that accounted for clustering were extracted. For cluster-randomised controlled trials, we ensured that all studies included in the meta-analyses accounted for clustered data in their statistical analyses. In cases where a study included multiple groups receiving the same or different intervention, a group combination to create a single pair-wise comparison was used [28]. A random-effects restricted maximum-likelihood model was utilized for the meta-analyses.

The Q statistic and I^2 [27], and visual inspection of the forest plots, were used to examine statistical heterogeneity. I^2 indicated high heterogeneity when $>75\%$ and moderate when $>50\%$. Random effect meta-regression models were used to explore heterogeneity induced by the relationship between moderators (i.e., participants' age [years], duration of the intervention [weeks], type of intervention [single-or multi-component]) and study effect sizes, when there were more than ten studies in the meta-analysis [28]. Where the meta-regression suggested the presence of a potentially important covariate, subgroup analyses were used to further investigate the data. For all meta-analysis and meta-regression models, statistical significance was set at $p < 0.05$ and effect sizes were interpreted as 0.2 small, 0.5 medium and 0.6 as large [29]. To assess potential small-study effects and publication bias for meta-analysis with at least 10 studies (i.e. MVPA, BMI, and WC) [30], funnel plots were produced and the Egger regression asymmetry test [31] was conducted. The trim-and-fill [32] computation was also used to assess the effect of publication bias on the interpretation of results.

Results

Characteristics of the Included Studies

In total, 85 articles (from 63 studies) fulfilled the inclusion criteria (Table 1). Of the included studies, 17 (27%) were conducted in Australia, followed by the United Kingdom ($n=9$, 14.3%), the USA ($n=6$, 9.5%) and other countries ($n=31$, 49%). Study sample sizes ranged from

33 [33] to 6,476 [34] (Table 1). In total, the studies included 45,733 participants, with 50 articles (58.9%) reporting on physical activity, 31 (36.4%) on sedentary time, 35 (41.2%) on physical health, and 12 (14.1%) on psychological outcomes. Participant ages ranged from 11 to 17 years and the average age was 13.7 years (reported in 44 articles, 51.8%). The percentage of girls varied between 36% [35] and 100% [36–41]. The participants' sex was not reported in five studies [34, 42–45]. Most of the studies ($n=46$, 73%) used a randomised controlled trial design [33–39, 42, 44, 46–78], 15 studies (23.8%) used a quasi-experimental design [40, 79–91], one study (1.6%) used a cross-over trial design [43] and one (1.6%) a hybrid effectiveness-implementation design [92]. Intervention length ranged from a single 20-min session [52, 62] to three years [81].

Strategies Used within Secondary-School Interventions and their Effect on Movement Behaviours, Cognitive/Academic, Physical Health and/or Psychological Outcomes

Various unique intervention strategies were reported ($n=23$) across 11 categories (i.e., active lessons, community involvement, educational resources, environmental, incentives/rewards, peer support, physical activity session/s, research support, school policy, teacher training, and technology strategies). A definition, detailed description, and examples of these categories can be found in Table 2 and Supplementary Table 2, respectively. Single component interventions using only one strategy ($n=19$ studies, 28.6%) involved physical activity sessions ($n=10$), peer-led support ($n=6$), or an environmental modification strategy ($n=3$). Most interventions ($n=45$ studies, 71.4%) involved the use of a combination of two or more strategies ('*multicomponent interventions*'). The strategies most frequently used were: physical activity sessions ($n=31$); environmental modifications ($n=29$); educational resources ($n=25$ interventions); peer support ($n=20$); teacher training ($n=18$); supporting technology ($n=11$); active lessons ($n=8$); community involvement ($n=7$); research support ($n=6$); incentives/rewards ($n=4$); and school policies ($n=3$) (Table 3). Although some interventions targeted several elements of the school day (e.g., class, recess/lunch, homework), 29 studies (46%) included a class-time component [37, 43, 44, 46, 50–55, 57–59, 63, 64, 66–72, 74, 79, 80, 84, 87, 89, 90], 22 studies (34.9%) a recess/lunch component [34, 35, 37, 38, 46, 49, 56, 57, 63, 65, 68–70, 72, 76, 77, 83, 86, 88, 91–94], 15 (23.8%) studies a whole-school day approach [36, 39, 40, 50, 56, 57, 61, 63, 68, 69, 72, 73, 75, 81, 83, 95], and one intervention (1.6%) [96] included an active homework component. Table 4 summarizes studies that report a positive impact on movement behaviours, cognitive/academic performance, physical wellbeing, and psychological outcomes, sorted by their methodological

Table 1 Characteristics of the included studies and study populations description

Authors	Design*	n	Sex (%)	Age (y) (Mean, range)	Length (weeks)	Outcomes [‡]	Overall methodological quality rating
Aceves-Martins et al., 2017 [55]	RCT - cluster	393	I=50.6% B C=47.5% B	14.6±0.7, 13–16	52	PA, SED	Moderate
Ahmed et al., 2022 [95]	RCT - cluster	320	I=65.6% B C=51.9% B	I=14.4±1.2 C=14.2±0.9, 13–17	12	PA, SED	Weak
Altunkurek et al., 2019 [54]	RCT	132	I ₁ : 42.4% B I ₂ : 50% B C: 51.5% B	12–15	12	PA, WB	Weak
Amoah et al., 2021 [94]	RCT - cluster	848	51.3% G	16.9±1.4 14–19	26	PA, H	Weak
Andrade et al., 2014 [70]	RCT - cluster	1234	37.6% B 62.4% G	12.8±0.8	121	PA, SED, H, F	Moderate
Andrade et al., 2015 [96]						SED	Moderate
Ardic et al., 2016 [89]	Quasi - Ex	87	50.6% G	12.8±0.8, 12–15	52	PA, H	Weak
Barbosa Filho et al., 2016 [63]	RCT - cluster	1085	I=51.8% B C=51.2% B	11–17	17	PA	Moderate
Barbosa Filho et al., 2017 [125]						PA, H	Moderate
Barbosa Filho et al., 2019 [97]						PA	Moderate
Bandeira et al., 2020 [109]						SED	Moderate
Bell et al., 2017 [42]	RCT	928	No reported	12–13	17	PA, SED	Weak
Bogart et al., 2016 [49]	RCT	1368	49.1% B	12.2±0.68	5	H	Weak
Bonhauser et al., 2005 [87]	Quasi - Ex	198	I=45.9% G C=57% G	I=15.5±0.8 C=15.5±0.9	43	PA, F, WB	Weak
Budde et al., 2010 [52]	RCT	59	I ₁ =50% B I ₂ =55% B C=62% B	14.4±0.5	12 min	H, AC	Weak
Bush et al., 2010 [126]	Quasi - Ex	191	I=48% B C=32% B	I=13.9±0.7 B I=13.8±0.9 G C=12.5±0.6 B C=12.5±0.6 G	16	PA AC	Weak Weak
Carlin et al., 2018 [76]	RCT - cluster	197	100% G	11–13, 12.4±0.6	12	PA, SED, H, F, WB	Moderate
Chen et al., 2020 [85]	Quasi - Ex	377	I ₁ =53.6% G I ₂ =55.4% G	No reported	12	PA	Weak
Contardo Ayala et al., 2018 [80]	Quasi - Ex	105	43.2% G	14.8±1.7, 12–17	17	H, F PA, SED, AC	Weak Weak
Sudholz et al., 2020 [100]							
Corder et al., 2016 [51]	RCT	788	I=47.7% B C=43.5% B	I=13.2±0.4 C=13.1±0.3	8	PA, WB	Moderate
Corder et al., 2020 [73]	RCT - cluster	2862	I=46.6% G C=48.9% G	I=13.2±0.4 C=13.2±0.4	12	PA, SED, H, WB	Weak
Corepal et al., 2019 [56]	RCT - cluster	224	46.9% B	12–14	22	PA, SED	Weak
Costigan et al., 2018 [46]	RCT	65	69% B	15.8±0.6	8	PA	Weak
Cui et al., 2012 [53]	RCT	682	I=50.9% B C=52.7% B	I=12.7±0.5 B I=12.6±0.5 G C=12.8±0.5 B C=12.6±0.4 G	4	PA, SED	Weak
Gammon et al., 2019 [71]	RCT - cluster	222	51% B	No reported	12	PA, SED	Weak
Ghammam et al., 2017 [81]	Quasi - Ex	4003	I=50.2% B C=46.5% B	11–16	156	PA, SED	Weak
Haapala et al., 2017 [86]	Quasi - Ex	319	66.7% G	14.0±0.6	87	PA, SED	Weak
Haerens et al., 2006 [35]	RCT - cluster	2840	36.6% G	13.1±0.8, 11–15	43	PA, H PA	Moderate Moderate
Haerens et al., 2007 [98]							
Harrington, 2018 [36]	RCT - cluster	1752	100% G	11–14	61	PA, SED	Strong
Gorely et al., 2019 [115]						FRI	Strong

Table 1 (continued)

Authors	Design*	n	Sex (%)	Age (y) (Mean, range)	Length (weeks)	Outcomes [‡]	Overall methodological quality rating
Hollis et al., 2016 [60]	RCT - cluster	1150	48% B	12	104	H	Strong
Sutherland et al., 2016 [102]						PA, FRI	Strong
Sutherland et al., 2016 [127]						PA, FRI	Moderate
Sutherland et al., 2020 [34]	RCT - cluster	6476	No reported	No reported	52	FRI	Weak
Sutherland et al., 2021 [45]					104	FRI	Weak
James et al., 2020 [48]	RCT	909	51.2% B 48.8% G	13–14	52	H, F	Weak
Kariippanon et al., 2019 [43]	Cross-over trial	171	No reported	13.2 ± 1	2	PA, SED	Weak
Kennedy et al., 2018 [77]	RCT - cluster	607	50.1% G	14.1 ± 0.5	10	F, PA, FRI	Weak
Kennedy et al., 2019 [117]						FRI	Weak
Kennedy et al., 2021 [92]	Hybrid	750	47.6% G	14.4 ± 1	10	F, FRI	Weak
Knebel et al., 2020 [68]	RCT - cluster	597	I = 51% G C = 54.1% G	13.0 ± 1.0	39	SED	Weak
Knox et al., 2012 [84]	Quasi - Ex	192	No reported	I = 12.4 ± 0.5 C = 12.1 ± 1.1	18	PA, H	Weak
Kolle et al., 2020 [65]	RCT - cluster	2084	I ₁ = 50% G I ₂ = 49% G C = 49% G	14 ± 0.3	39	PA, SED, F	Strong
Solberg et al., 2021 [78]						AC	Weak
Lazorick et al., 2015 [90]	Quasi - Ex	362	I = 53% B C = 51% B	I = 13.3 ± 0.8 C = 13.1 ± 0.5	14	PA, H	Weak
Leme et al., 2016 [107]	RCT - cluster	253	100% G	15.61 ± 0.05	26	PA, SED, H	Weak
Leme et al., 2018 [37]						PA, SED, H	Weak
Lubans et al., 2010 [93]	RCT	108	I ₁ = 59% B I ₂ = 49% B C = 47% B	15.0 ± 0.7	8	H, F	Weak
Lubans et al., 2011 [104]	RCT	100	100% B	14.3 ± 0.6	26	PA, H	Moderate
Morgan et al., 2012 [114]						WB	Weak
Lubans et al., 2012 [50]						PA, H	Moderate
Lubans et al., 2012 [118]	RCT - cluster	357	100% G	13.2 ± 0.5	52	PA, SED, H, F	Strong
Dewar et al., 2013 [59]						PA, SED, H	Strong
Dewar et al., 2014 [103]						PA, SED, AC	Strong
Lubans et al., 2021 [99]	RCT - cluster	670	44.6% G	16.0 ± 0.43	52	PA, H, AC, F, WB	Moderate
Mavilidi et al., 2020 [66]	RCT - cluster	221	49.8% G	16.0 ± 0.5 16–18	52	AC	Weak
Valkenborghs et al., 2022 [113]	RCT - cluster	56	61% G	16.1 ± 0.4	26	H	Strong
Ludyga et al., 2018 [64]	RCT - cluster	36	I = 12% G C = 24.2% G	12–15	8	AC	Weak
Ludyga et al., 2019 [62]	RCT - cluster	94	100% B	13.9 ± 0.8, 12–15	20 min session	AC	Weak
Melnyk et al., 2013 [67]	RCT - cluster	779	51.5% G	14–16	26	PA, H, AC	Weak
Melnyk et al., 2015 [111]						H, WB	Weak
Murphy et al., 2022 [41]	Quasi - Ex	85	100% G	13 ± 0.7	10	PA, WB, F	Weak
Okely et al., 2017 [38]	RCT - cluster	1769	100% G	13.6 ± 0.0	78	PA, SED	Strong
Parrish et al., 2018 [57]	RCT - cluster	88	50% B	14.7 ± 0.7, 13–16	22	PA, SED, AC	Moderate
Peralta et al., 2009 [33]	RCT	33	100% B	12.5 ± 0.4	26	PA, H, F	Weak
Schofield et al., 2005 [40]	Quasi - Ex	90	100% G	15.8 ± 0.8, 15–18	12	PA, H	Weak
Sebire et al., 2018 [39]	RCT - cluster	427	100% G	12–13	22	PA	Moderate
Sebire et al., 2019 [116]						FRI	Moderate

Table 1 (continued)

Authors	Design*	n	Sex (%)	Age (y) (Mean, range)	Length (weeks)	Outcomes [‡]	Overall methodological quality rating
Smith et al., 2014 [106]	RCT - cluster	361	100% B	12.7 ± 0.5	20	PA, SED, H	Weak
Lubans et al., 2016 [108]						PA, SED	Weak
Lubans et al., 2016 [69]						SED, H, F, WB	Weak
Subramanian et al., 2015 [47]	RCT	439	43.1% G	12–17	26	AC	Weak
Suchert et al., 2015 [61]	RCT - cluster	1162	48% G	13.7 ± 0.7, 12–17	12	PA, SED, F	Moderate
Sudholz et al., 2016 [79]	Quasi - Ex	43	49% G	13.7 ± 1.4, 12–16	7	PA, SED	Weak
Tarp et al., 2016 [72]	RCT - cluster	632	48.9% G	12.9 ± 0.6	20	PA, H, AC, F	Moderate
Torbeyns et al., 2017 [44]	RCT	56	51.8% B	14.3 ± 0.6	22	H, F	Weak
Tymms et al., 2016 [58]	RCT - cluster	1494	I=51.0% G C=53.3% G	I=11.72 C=11.79	6	PA	Weak
Van Woudenberg et al., 2018 [75]	RCT - cluster	190	53.68% G	12.17, 11–14	1	PA	Weak
Verloigne et al., 2018 [74]	RCT - cluster	156	54.5% G	15.5 ± 0.5	26	PA, SED, WB	Moderate
Yang et al., 2017 [83]	Quasi - Ex	820	I=73.2% B C=79.1% B	I=10.9 ± 1.6 C=11.0 ± 1.5	52	PA, H, F	Weak
Yli-Piipari et al., 2016 [82]	Quasi - Ex-	94	51.1% G	11–15	4	PA, F	Weak
Yu et al., 2021 [91]	Quasi - Ex	514	No reported	No reported	21	PA, FRI	Weak

Abbreviations: n=number; y=years. * (Study design) RCT=randomised controlled trial; Quasi - ex=quasi experimental; Hybrid=hybrid effectiveness-implementation trial design

[‡] (Sex) B=boys; G=girls; I=Intervention group; C=comparison/control group [‡] (Outcomes) PA=physical activity; SED=sedentary behaviour; WB=wellbeing; H=health; F=fitness, AC=academic outcomes; FRI=factors relating to the implementation

Table 2 Definitions of the intervention strategies reported

Name of the strategy	Definition
Active lessons	Teachers' normal planned class lessons, where the delivery method rather than the content is changed (PA into curriculum subjects)
Community involvement:	Participation/support of community members/facilities outside schools
Educational	Theory based sessions only (health-related)
Environmental modifications:	A supportive school environment encourages physical activity throughout the school day (e.g. active indoor and outdoor environments, active equipment)
Incentives/rewards:	Incentives to promote or reward physical activities or certain goals
Physical activity sessions	Opportunities for PA (supervised or unsupervised) during schools' hours (e.g. during recess and lunchbreak)
Peer support:	Adolescent leaders to encourage PA among their peers
Research support:	Direct support from the research team involved
School policy	Changes, adaptation of the school policies to encourage physical activity and reduce sitting time
Teacher training	Teacher development sessions, pedagogical strategies

Abbreviations: PA=physical activity

quality. These interventions encompassed a range of strategies, including physical activity sessions, environmental enhancements, teacher training, peer support, and educational resources.

Intervention Effects on Movement Behaviours

Sixty-three articles (74%) described intervention effects on *movement behaviours* (self-reported and/or device measured) (Table 1, outcomes). Of these, 40 articles (47%) reported on device-measured physical activity (i.e., light- [LPA], moderate- [MPA], vigorous- [VPA],

moderate- to vigorous-intensity [MVPA] physical activity). Sedentary time was reported in 31 articles (36.4%); of these, 20 (23.5%) used device-measured sedentary/sitting time. A descriptive summary of the number of intervention strategies and their impact on different outcomes is presented in Supplementary Table 3.

Sufficient data for meta-analysis were available for sedentary time, LPA, MPA, VPA, MVPA assessed during the whole day (Summary statistics used for calculation of standardized mean difference across the studies are presented in Supplementary Table 7). All meta-analyses

Table 3 Frequency of intervention strategies used across studies

Strategies used	References
PA session	[33, 34, 38, 41, 46, 47, 54, 60, 62, 64–67, 71, 72, 77, 82–84, 86, 90, 92, 94, 95, 99, 104, 106, 107, 112, 113, 118]
Environment	[34, 35, 40, 43, 44, 49, 55–57, 66, 68, 70, 74, 77, 79–81, 83, 85, 89, 91, 95, 99, 104, 106, 113, 118, 125]
Educational	[33, 38, 43, 44, 47, 54, 57, 66, 74, 76, 78, 79, 81, 86, 89, 90, 92, 94–96, 101, 112, 116, 117, 127]
Peer support	[36, 39, 42, 48, 49, 51, 53, 55, 56, 58, 60, 67, 70, 73, 75, 76, 81, 104, 106, 126]
Teacher training	[36, 43, 57, 58, 60, 63, 66–68, 74, 77, 80, 85, 90, 99, 107, 113, 117, 118]
Technology	[55, 56, 61, 72, 77, 92, 99, 106, 113, 118]
Community involvement	[41, 76, 86, 94, 101, 115, 116, 127]
Active lessons	[52, 63, 65, 80, 87, 89, 126]
Research support	[41, 43, 45, 52, 115, 117]
Reward/ Incentives	[33, 45, 51, 55]
School policy	[41, 115, 127]

showed non-significant effects: Sedentary time, SMD = -0.02 (95% CI: -0.14, 0.11), Fig. 2.A; LPA, SMD = -0.01 (95% CI: -0.08, 0.05), Fig. 2.B; MPA, SMD=0.06 (95% CI: -0.09,0.22), Fig. 2.C; VPA, SMD=0.08 (95% CI: -0.02, 0.18), Fig. 2.D; MVPA, SMD=0.05 (95% CI: -0.01, 0.12), Fig. 2.E).

Intervention Effects on Cognitive and Academic Outcomes

Cognitive and academic outcomes were measured in 12 articles (14.1%) which did not fulfill the inclusion criteria for meta-analyses. The most frequently explored outcomes were executive functions. Evidence of a positive impact was reported on working memory [52, 57, 64], inhibitory control [62], cognitive flexibility [47], verbal fluency [47], attention and concentration [47, 110], and on-task behaviour [66]. No effects were found on inhibitory control [72, 99], and working memory [99]. A positive intervention effect was found on health course grades [67], numeracy and reading performance [78] and academically-relevant social skills ratings (cooperation, assertion, academic competence) [67]. No effect was found on mathematical skills [72].

Intervention Effects on Physical Health Outcomes

Physical health related outcomes were assessed in 29 articles (34.1%). Meta analysis of 16 articles (18.8%) showed a statistically significant small effect on BMI (SMD= -0.09 [95% CI: -0.16, -0.02]) with a moderate level of heterogeneity and 56% of the articles were assessed with high-risk of bias; Fig. 3.A). For waist circumference, 5 articles (5.9%) provided necessary data for meta-analysis. A non-significant intervention effect was found (SMD=0.09 [95% CI: -0.03, 0.21]; Fig. 3, B).

Evidence of a positive effect was reported for blood pressure [48, 83, 84, 94], high-density lipoprotein (HDL) to total cholesterol ratio [84], and glucose [84]. No effects on body fat [33, 106], blood pressure [76] or musculoskeletal health [80] were found. Fitness-related outcomes were assessed in 19 articles (22.4%). Positive effects on cardiorespiratory fitness (e.g. increase distance run) [44,

48, 65, 72, 87, 92, 99], and muscular fitness (e.g. muscular endurance strength) were found [69, 70, 83, 87, 99]. No effects on cardiorespiratory fitness [33, 41, 61, 76, 82] and muscular fitness [41, 50, 65, 82, 112] were reported. Hippocampal metabolism was assessed [113], showing a significant intervention effect on N-acetylaspartate and glutamate+ glutamine in the left hippocampus.

Intervention Effects on Psychological Outcomes

Psychological outcomes were assessed in 12 articles (14.1%), which did not fulfill the inclusion criteria for meta-analyses. Positive effects were found in: anxiety and self-esteem [87], social support [51, 76], wellbeing [41, 51, 54, 69], self-perception [114], and lower depressive symptoms in participants with elevated depressive symptoms at baseline [111]. No effects on wellbeing [73, 99] or on stress and internalization/externalization problems [99] were reported, and one study indicated that the relationship with classmates deteriorated in the intervention group [74]. Also, there were no effects on social cognitive variables for physical activity (i.e., self-efficacy, perceived environment, social support, behavioural strategies, outcome expectations and outcome expectancies related to physical activity) [103].

Methodological Quality

Of the included articles, 65% received an overall weak quality rating, 25% were rated as moderate, and 11% were considered strong quality (Supplementary Table 4). A breakdown of the methodological quality per article is shown in Table 5. Study design, data collection methods, confounders, withdrawals and dropout were assessed as strong in >50% of the articles. Selection bias was considered weak in 45% of the articles. For articles that had a positive effect on an outcome reported in this review ($n=58$) [68%], 16 were classified as strong, 17 as moderate, and 25 as weak (Table 4).

Table 4 (continued)

	Methodological quality	Movement behaviour	Academic outcomes	Physical Health	Psychological outcomes
Knox et al., 2012 [84]	Weak			✓	✓
Laberge et al., 2012 [110]	Weak		✓		
Lazorick et al., 2015 [90]	Weak			✓	
Leme et al., 2016 [107]	Weak	✓			
Lubans et al., 2010 [93]	Weak			✓	
Lubans et al., 2016 [108]	Weak	✓			
Lubans et al., 2016 [69]	Weak	✓			✓
Ludyga et al., 2018 [64]	Weak		✓		
Ludyga et al., 2019 [62]	Weak		✓		
Mavilidi et al., 2020 [66]	Weak		✓		
Melnyk et al., 2013 [67]	Weak	✓	✓		
Melnyk et al., 2015 [111]	Weak			✓	✓
Morgan et al., 2012 [114]	Weak			✓	✓
Murphy et al., 2022 [41]	Weak				✓
Schofield et al., 2005 [40]	Weak	✓			
Smith et al., 2014 [106]	Weak				
Solberg et al., 2021 [78]	Weak	✓	✓		
Subramanian et al., 2015 [47]	Weak		✓		
Sudholz et al., 2016 [79]	Weak	✓			
Sudholz et al., 2020 [100]	Weak	✓	✓		
Torbeyns et al., 2017 [44]	Weak				
Yang et al., 2017 [83]	Weak				
Yu et al., 2021 [91]	Weak	✓			✓

Legend: ✓ indicates a study that reported a positive significant effect on movement behaviours, academic outcomes, physical health and/or psychological outcomes




Fig. 2 Forest plots for the effect of school-based initiatives on device measured sedentary time (A), light intensity physical activity (B), moderate physical activity (C), vigorous physical activity (D), moderate-to-vigorous physical activity (E). Legend: Risk of bias: (SB) selection bias; (SD) study design; (C) confounder; (B) blinding; (DC) data collection; (W) withdrawal and (OR) overall risk




Fig. 3 Forest plots for the effect of school-based initiatives on body mass index (A) and waist circumference (B). Legend: Risk of bias: (SB) selection bias; (SD) study design; (C) confounder; (B) blinding; (DC) data collection; (W) withdrawal and (OR) overall risk

Factors Relevant to the Implementation

At least one *factor relevant to the implementation* of the programs (implementation, mechanism of impact and/or context) was reported in 43 articles (50.5%). Specific components reported included *fidelity* ($n=26$, 30.5%) [33, 34, 38, 45, 53, 57, 58, 60, 63, 66, 70, 73, 77, 82, 92, 99, 101, 102, 105, 107, 111, 115–117]. Fidelity was assessed based on adherence to the intervention protocol and consistency in delivery, of which 14 articles (16.4%) indicated that interventions were delivered as planned (>60% of consistency with the original plan); *participant responses* ($n=25$, 29.4%) [33, 38, 46, 51, 53, 55–57, 67, 70, 73, 74,

77, 79, 88, 92, 101, 106, 107, 111, 115–117], where 20 articles (23.5%), indicated the majority of the participants enjoyed or participated in the intervention; *dose* ($n=18$, 21.2%) [34, 45, 48, 57, 60, 63, 66, 70, 73, 77, 85, 87, 99, 101, 102, 105, 115, 117], of which 10 indicated that the intervention was delivered and received as planned (>50% of the frequency and duration of the intervention was delivered); *reach* ($n=18$, 21.2%) [33, 34, 45, 55, 60, 63, 64, 70, 88, 92, 93, 106, 107, 115–117], where 13 indicated that the intervention targeted the expected audience; *context* ($n=15$, 17.6%) [33, 38, 42, 47, 48, 51, 74, 87, 99, 107, 115, 118], of which 4 indicated no adverse effects of the

Table 5 Methodological quality assessment for each article

Study	Selection bias	Study design	Confounder	Blinding	Data collection	Withdrawal	Overall Rating
Aceves-Martins et al., 2017 [55]	2	1	1	3	1	2	2
Ahmed et al., 2022 [95]	3	1	1	3	1	1	3
Altunkurek et al., 2019 [54]	3	1	1	3	3	3	3
Amoah et al., 2021 [94]	2	1	1	3	3	1	3
Andrade et al., 2014 [70]	1	1	1	3	1	2	2
Andrade et al., 2015 [96]	1	1	1	3	1	2	2
Ardic et al., 2016 [89]	3	2	1	3	1	1	3
Bandeira et al., 2020 [109]	1	1	1	3	1	1	2
Barbosa Filho et al., 2016 [63]	1	1	1	3	1	1	2
Barbosa Filho et al., 2017 [125]	1	1	1	3	1	1	2
Barbosa Filho et al., 2019 [97]	1	1	1	3	1	1	2
Bell et al., 2017 [42]	3	1	3	3	1	1	3
Bogart et al., 2016 [49]	2	1	1	3	1	3	3
Bonhauser et al., 2005 [87]	3	3	1	3	3	1	3
Budde et al., 2010 [52]	3	1	3	3	3	1	3
Bush et al., 2010 [126]	2	3	1	3	1	2	3
Carlin et al., 2018 [76]	2	1	1	3	1	1	2
Chen et al., 2020 [85]	3	3	3	3	2	2	3
Contardo Ayala et al., 2018 [80]	3	3	1	3	1	1	3
Corder et al., 2016 [51]	2	1	3	3	1	1	3
Corder et al., 2020 [73]	1	1	3	2	1	2	2
Corepal et al., 2019 [56]	2	1	3	3	1	1	3
Costigan et al., 2018 [46]	3	1	3	2	1	3	3
Cui et al., 2012 [53]	3	1	1	3	2	1	3
Dewar et al., 2013 [59]	2	1	1	2	1	1	1
Dewar et al., 2014 [103]	2	1	1	2	1	1	1
Gammon et al., 2019 [71]	3	1	3	3	1	1	3
Ghammam et al., 2017 [81]	1	3	1	3	3	1	3
Gorely et al., 2019 [115]	2	1	3	2	1	3	3
Haapala et al., 2017 [86]	3	3	1	3	1	3	3
Haerens et al., 2006 [35]	1	1	1	3	1	1	2
Haerens et al., 2007 [98]	1	1	1	3	1	1	2
Harrington, 2018 [36]	2	1	1	2	1	2	1
Hollis et al., 2016 [60]	1	1	1	2	1	1	1
James et al., 2020 [48]	2	1	3	3	1	1	3
Kariippanon et al., 2019 [43]	2	3	1	3	1	1	3
Kennedy et al., 2018 [77]	2	1	3	3	1	2	3
Kennedy et al., 2019 [117]	2	1	3	3	1	2	3
Kennedy et al., 2021 [92]	2	3	3	3	1	3	3
Knebel et al., 2020 [68]	3	1	1	3	3	2	3
Knox et al., 2012 [84]	3	3	1	3	1	3	3
Kolle et al., 2020 [65]	2	1	2	2	1	1	1
Laberge et al., 2012 [110]	2	3	3	3	1	2	3
Lazorick et al., 2015 [90]	2	3	1	3	1	2	3
Leme et al., 2016 [107]	3	1	3	2	1	2	3
Leme et al., 2018 [37]	3	1	3	2	1	2	3
Lubans et al., 2010 [93]	3	1	3	3	1	2	3
Lubans et al., 2011 [104]	2	1	1	3	1	1	2
Lubans et al., 2012 [118]	2	1	1	2	1	1	1
Lubans et al., 2012 [50]	2	1	1	3	1	1	2
Lubans et al., 2016 [108]	3	1	3	2	1	1	3
Lubans et al., 2016 [69]	3	1	3	2	1	1	3
Lubans et al., 2021 [99]	2	1	1	3	1	1	2

Table 5 (continued)

Study	Selection bias	Study design	Confounder	Blinding	Data collection	Withdrawal	Overall Rating
Ludyga et al., 2018 [64]	3	1	1	3	1	1	3
Ludyga et al., 2019 [62]	3	1	3	3	1	1	3
Mavilidi et al., 2020 [66]	2	1	3	3	1	1	3
Melnyk et al., 2013 [67]	3	1	1	3	1	2	3
Melnyk et al., 2015 [111]	3	1	1	3	1	2	3
Morgan et al., 2012 [114]	2	1	3	3	1	1	3
Murphy et al., 2022 [41]	3	3	3	3	3	3	3
Okely et al., 2017 [38]	1	1	1	2	1	2	1
Parrish et al., 2018 [57]	3	1	1	2	1	1	2
Peralta et al., 2009 [33]	3	1	3	2	1	1	3
Schofield et al., 2005 [40]	2	3	1	3	1	2	3
Sebire et al., 2018 [39]	1	1	1	3	1	1	2
Sebire et al., 2019 [116]	1	1	1	3	1	1	2
Smith et al., 2014 [106]	3	1	3	3	1	2	3
Solberg et al., 2021 [78]	2	1	3	3	1	1	3
Subramanian et al., 2015 [47]	3	1	1	3	3	2	3
Suchert et al., 2015 [61]	1	1	1	3	1	1	2
Sudholz et al., 2016 [79]	3	3	3	3	1	3	3
Sudholz et al., 2020 [100]	3	3	1	3	1	1	3
Sutherland et al., 2020 [34]	3	1	1	3	1	1	3
Sutherland et al., 2021 [45]	3	1	3	3	1	1	3
Sutherland et al., 2016 [102]	1	1	1	2	1	2	1
Sutherland et al., 2016 [127]	1	1	3	2	1	2	2
Tarp et al., 2016 [72]	2	1	1	3	1	2	2
Torbeyns et al., 2017 [44]	3	1	3	3	1	1	3
Tymms et al., 2016 [58]	3	1	1	3	1	1	3
Valkenborghs et al., 2022 [113]	2	1	1	2	1	1	1
Van Woudenberg et al., 2018 [75]	3	1	1	3	1	1	3
Verloigne et al., 2018 [74]	2	1	1	3	1	1	2
Yang et al., 2017 [83]	3	3	3	3	3	1	3
Yli-Piipari et al., 2016 [82]	3	1	3	3	1	3	3
Yu et al., 2021 [91]	3	3	3	3	2	3	3

Legend: 1=strong, 2=moderate, 3=weak methodological quality

intervention, and 5 articles showed that the intervention needed some modifications (e.g. different devices, time for planning, resources); and *adaptation* was required in 6 studies (i.e. content and frequency adaptations) [34, 45, 63, 74, 77, 117].

Publication Bias

Funnel plots for MVPA, BMI and WC are presented in Supplementary material 6, and show little sign of asymmetry. In addition, the Egger's tests for all three outcomes were not statistically significant, indicating absence of small-sample effects. There was little evidence of publication bias with pooled effect size estimates using the trim-and-fill method similar to the main findings for MVPA (SMD=0.06 vs. 0.05), with overall interpretations of effects unchanged for BMI (SMD=-0.08 vs. -0.08) and WC (SMD=1.46 vs.1.46).

Discussion

Our systematic review provided a narrative synthesis of the intervention strategies used within the secondary school settings to improve students' movement behaviours throughout school-based initiatives and the factors related to intervention implementation, and a meta-analysis (where possible) of the effectiveness of school-based initiatives to increase physical activity and reduce sedentary time on adolescents' movement behaviours, energy expenditure, cognitive/academic, behavioural and physical and psychological health outcomes. Intervention strategies that reported favourable effects on movement behaviours, cognitive/academic, physical and psychological outcomes tended to include physical activity sessions, environmental modifications, teacher training, peer support and/or educational resources. Despite some promising findings for BMI, the meta-analysis showed no significant effects of interventions on the total

accumulated daily movement behaviours, assessed using accelerometers, and waist circumference.

Interventions incorporated a wide variety of strategies, with the majority being multicomponent. Previous reviews have concluded multicomponent interventions elicit greater effects on youth health and wellbeing compared to single component interventions [119], particularly those that use a whole-school approach [9]. However, it can be difficult to establish which strategies embedded within multicomponent interventions are most effective and may be context specific. Homework and recess time were seldom targeted in interventions. This is consistent with a recent review and meta-analysis of school-based recess interventions, which reported that just one of 43 interventions targeted secondary school students [120]. Physical activity sessions, educational resources, environmental modifications, peer-support, and teacher training were the most commonly used strategies, consistent with a previous review of school-based interventions to increase adolescent physical activity [16]. Although active lessons are commonly used and can have positive effects on lesson-time physical activity in primary/elementary school interventions [121], only eight of the 63 interventions in this review incorporated active lessons. There is clearly a need for evidence of the effectiveness of interventions involving active lessons, recess and lunch breaks and active homework.

Our meta-analyses revealed small positive effects of overall interventions on BMI; however, most articles were assessed with a high risk of bias. Results related to BMI are consistent with findings from a previous review indicating that school-based physical activity interventions result in a very small decrease in BMI z-score in children and adolescents [23]. We did not find significant effects for accelerometer-determined physical activity of any intensity. This is consistent with previous meta-analyses, which found that school-based physical activity interventions had small or no benefits on whole-day moderate-intensity physical activity levels of adolescents [15–17]. This review found few studies reported the impact on sedentary time and LPA, especially during school hours. It found a statistically significant effect for sitting and stepping time (data not shown, as two out of the four studies reporting these data did not meet the inclusion criteria for analysis [i.e. randomised controlled trials]). The effect of school-based initiatives on movement behaviours may have been under-estimated as these initiatives could be more effective at reducing sedentary time and increasing LPA rather than higher intensity activity during school hours. Future research should consider assessing multiple movement behaviour intensities (i.e., sedentary time, LPA) with the appropriate devices [122].

The quality assessment of the studies showed different distributions of effectiveness between weak, moderate

and strong quality. Further analyses showed no apparent differences in quality between studies that reported significant effects and those that did not for each outcome; however, only few studies were classified with low risk of bias. Only 43 articles (50.5%) reported factors relevant to the implementation of the programs. Therefore, with the lack of and inconsistencies in reporting implementation-relevant information, it is difficult to determine whether the interventions were sufficiently or appropriately implemented as intended. Insufficient equipment or supportive environments (e.g., standing desks), high dropout rates due to academic commitments, cancelled sessions due to unplanned activities, lack of time to plan and deliver, and programs not viewed as a priority were reported as factors that could influence intervention effectiveness. Evaluating factors related to implementation before and during the intervention, that allows researchers and schools to adapt and improve the implementation strategies to the school's and students' needs, could achieve better results [123].

Limitations of included studies were the high level of heterogeneity between outcome measures, and that only a small number of studies reported movement behaviours during school hours. It could be possible that interventions were effective during school hours, but effects were compensated for outside the school [124]. Future school-based interventions should consider (a) incorporating whole of school approaches (b) including additional outcomes such as academic performance and reporting movement behaviour outcomes for school hours and the whole day and multiple intensities, (c) reporting factors relevant to the implementation to assist in interpreting effectiveness.

Schools should be encouraged to develop and implement policies that support whole-of-school physical activity strategies. These policies could include providing teaching relief so teachers can receive professional development (e.g., to deliver active lessons in the classroom), and allocating space in the curriculum for additional physical activity sessions and delivery of educational resources.

Conclusions

While some intervention approaches for increasing adolescents' physical activity and reducing sedentary time in secondary schools demonstrated promise (e.g. physical activity sessions, environmental modifications, teacher training, peer support, educational resources and/or active lesson strategies), small or non-significant effects were found in the meta-analyses. Future movement behaviour interventions in secondary schools should utilize a whole-school approach to beneficially change adolescents' activity levels. Consistent reporting

of implementation will increase understanding of how interventions effect outcomes.

Abbreviations

SMD	Standardized mean difference
CI	Confidence interval
MVPA	Moderate- to vigorous-intensity physical activity
PE	Physical education
PICOS	Participants, Interventions, Comparisons, Outcomes, and Study design framework
RCTs	Randomized controlled trials
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
LPA	Light physical activity
MPA	Moderate physical activity
VPA	Vigorous physical activity
BMI	Body mass index
HDL	High-density lipoprotein
N	Number
SED	Sedentary behaviour
EE	Energy expenditure
WC	Waist circumference
CRF	Cardiorespiratory fitness
SB	Selection bias
SD	Study design
C	Confounder
B	Blinding
DC	Data collection
W	Withdrawal
OR	Overall risk

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40798-024-00688-7>.

Supplementary Material 1

Acknowledgements

Not applicable.

Author Contributions

JS, AMC, KP, EM, NL, DL, HK, NDR, and AT conceptualized and designed the study. AMC and KP searched the databases, screened the titles and abstracts of all identified studies, obtained the full-text papers of the relevant studies, applied the selection criteria, collected the data, entered the results, and assessed for study quality. EM cross-checked for accuracy of full text selection, data entry and study quality. AMC and GA performed and interpreted the statistical analysis. AMC and KP wrote the initial draft manuscript. AMC, JS, KP, EM, AT, NL, DL, HK, NDR, and GA revised and approved the final manuscript.

Funding

This study is supported by an NHMRC (National Health and Medical Research Council) Investigator Grant (APP 1176885).

Data Availability

The datasets used and/or analyzed in the current study will be supplied by the corresponding author upon reasonable request.

Declarations

Ethics Approval and Consent to Participate

Not applicable.

Consent for Publication

Not applicable.

Competing interests

AMCA and EM are supported by a NHMRC Investigator Grant (APP 1176885). NDR is supported by a National Heart Foundation of Australia Future Leader Fellowship (ID 101895). DRL is supported by a National Health and Medical Research Council (NHMRC) senior Research Fellowship (APP1154507). JS is supported by a Leadership Level 2 Fellowship, NHMRC (APP 1176885). Other authors report no competing interests.

Author details

¹Institute for Physical Activity and Nutrition (IPAN), Deakin University, Geelong, Victoria, Australia

²Alliance for Research in Exercise, Nutrition and Activity (ARENA), Allied Health & Human Performance, University of South Australia, Adelaide, South Australia, Australia

³Centre for Active Living and Learning, College of Human and Social Futures, University of Newcastle, Newcastle, NSW, Australia

⁴School of Exercise and Nutrition Sciences, Deakin University, Geelong, Victoria, Australia

⁵School of Health and Social Development, Deakin University, Geelong, Victoria, Australia

⁶Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

⁷Hunter Medical Research institute, New Lambton Heights, NSW, Australia

Received: 20 September 2022 / Accepted: 21 February 2024

Published online: 13 March 2024

References

- Rodriguez-Ayllon M, Estévez-López F, Cadenas-Sanchez C, Gracia-Marco L, Lubans DR, Ortega FB, et al. Physical activity, sedentary behaviour and Mental Health in Young people: a review of reviews. In: Pingitore A, Matorci F, Vassalle C, editors. Adolescent health and wellbeing. Cham: Springer International Publishing; 2019. pp. 35–73.
- Poitras VJ, Gray CE, Borghese MM, Carson V, Chaput JP, Janssen I, et al. Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Appl Physiol Nutr Metab*. 2016;41(6 Suppl 3):197–239.
- Gunnell KE, Poitras VJ, LeBlanc A, Schibli K, Barbeau K, Hedayati N, et al. Physical activity and brain structure, brain function, and cognition in children and youth: a systematic review of randomized controlled trials. *Ment Health Phys Act*. 2019;16:105–27.
- Liu M, Wu L, Ming Q. How does physical activity intervention improve Self-Esteem and Self-Concept in Children and adolescents? Evidence from a Meta-analysis. *PLoS ONE*. 2015;10(8):e0134804.
- Mahar MT. Classroom-based physical activity and On-Task Behavior. *Translational J Am Coll Sports Med*. 2019;4(17).
- Watson A, Timperio A, Brown H, Best K, Hesketh KD. Effect of classroom-based physical activity interventions on academic and physical activity outcomes: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act*. 2017;14(1):114.
- Carson V, Hunter S, Kuzik N, Gray CE, Poitras VJ, Chaput JP, et al. Systematic review of sedentary behaviour and health indicators in school-aged children and youth: an update. *Appl Physiol Nutr Metab*. 2016;41(6 Suppl 3):240–65.
- Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451–62.
- van Sluijs EMF, Ekelund U, Crochemore-Silva I, Guthold R, Ha A, Lubans D, et al. Physical activity behaviours in adolescence: current evidence and opportunities for intervention. *Lancet*. 2021;398(10298):429–42.
- Bauman AE, Petersen CB, Blond K, Rangul V, Hardy LL. The descriptive epidemiology of sedentary Behaviour. In: Leitzmann MF, Jochem C, Schmid D, editors. *Sedentary Behaviour Epidemiology*. Cham: Springer International Publishing; 2018. pp. 73–106.
- Arundell L, Salmon J, Koorts H, Contardo Ayala AM, Timperio A. Exploring when and how adolescents sit: cross-sectional analysis of activPAL-measured patterns of daily sitting time, bouts and breaks. *BMC Public Health*. 2019;19(1):653.
- Hollis JL, Sutherland R, Williams AJ, Campbell E, Nathan N, Wolfenden L, et al. A systematic review and meta-analysis of moderate-to-vigorous physical

- activity levels in secondary school physical education lessons. *Int J Behav Nutr Phys Act.* 2017;14(1):52.
13. Hardman K. The world-wide survey of physical education in schools: findings, issues and strategies for a sustainable future. The fellows lecture (part 3). *Br J Teach Phys Educ.* 2001 //;32(2).
 14. Ridgers ND, Parrish A-M, Salmon J, Timperio A. School Recess Physical Activity interventions. *The Routledge Handbook of Youth Physical Activity.* Routledge; 2020. pp. 504–22.
 15. van de Kop JH, van Kernebeek WG, Otten RHJ, Toussaint HM, Verhoeff AP. School-based physical activity interventions in Prevocational adolescents: a systematic review and Meta-analyses. *J Adolesc Health.* 2019;65(2):185–94.
 16. Borde R, Smith JJ, Sutherland R, Nathan N, Lubans DR. Methodological considerations and impact of school-based interventions on objectively measured physical activity in adolescents: a systematic review and meta-analysis. *Obes Rev.* 2017;18(4):476–90.
 17. Hynynen ST, van Stralen MM, Sniehotta FF, Araujo-Soares V, Hardeman W, Chinapaw MJ, et al. A systematic review of school-based interventions targeting physical activity and sedentary behaviour among older adolescents. *Int Rev Sport Exerc Psychol.* 2016;9(1):22–44.
 18. Love R, Adams J, van Sluijs EMF. Are school-based physical activity interventions effective and equitable? A meta-analysis of cluster randomized controlled trials with accelerometer-assessed activity. *Obes Rev.* 2019;20(6):859–70.
 19. Morton KL, Atkin AJ, Corder K, Suhrcke M, van Sluijs EM. The school environment and adolescent physical activity and sedentary behaviour: a mixed-studies systematic review. *Obes Rev.* 2016;17(2):142–58.
 20. Verstraeten R, Roberfroid D, Lachat C, Leroy JL, Holdsworth M, Maes L, et al. Effectiveness of preventive school-based obesity interventions in low- and middle-income countries: a systematic review. *Am J Clin Nutr.* 2012;96(2):415–38.
 21. Hollis JL, Sutherland R, Williams AJ, Campbell E, Nathan N, Wolfenden L et al. A systematic review and meta-analysis of moderate-to-vigorous physical activity levels in secondary school physical education lessons. *Int J Behav Nutr Phys Activity.* 2017;14(1).
 22. Owen MB, Curry WB, Kerner C, Newson L, Fairclough SJ. The effectiveness of school-based physical activity interventions for adolescent girls: a systematic review and meta-analysis. *Prev Med.* 2017;105:237–49.
 23. Neil-Sztramko SE, Caldwell H, Dobbins M. School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. *Cochrane Database Syst Rev.* 2021;9(9):CD007651.
 24. Moore Graham F, Audrey S, Barker M, Bond L, Bonell C, Hardeman W et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ: Br Med J.* 2015 03/19;350.
 25. Thomas BH, Ciliska D, Dobbins M, Micucci S. A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs.* 2004;1(3):176–84.
 26. Day SJ, Altman DG. Statistics notes: blinding in clinical trials and other studies. *BMJ* 2000 Aug 19–26;321(7259):504.
 27. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21(11):1539–58. 2002/06/15.
 28. Higgins JPTJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. *Cochrane Handbook for systematic reviews of interventions version 6.2* (updated February 2021). In: *Cochrane*, editor; 2021.
 29. Cohen J. *Statistical Power Analysis for the behavioral sciences.* Elsevier Science; 2013.
 30. Page MJ, Shamseer L, Altman DG, Tetzlaff J, Sampson M, Tricco AC, et al. *Epidemiology and Reporting Characteristics of Systematic Reviews of Biomedical Research: a cross-sectional study.* *PLoS Med.* 2016;13(5):e1002028.
 31. Sterne JAC, Egger M, Smith GD. Investigating and dealing with publication and other biases in meta-analysis. *BMJ.* 2001;323(7304):101.
 32. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics.* 2000;56(2):455–63.
 33. Peralta LR, Jones RA, Okely AD. Promoting healthy lifestyles among adolescent boys: the Fitness Improvement and Lifestyle Awareness Program RCT. *Prev Med.* 2009;48(6):537–42.
 34. Sutherland R, Campbell E, McLaughlin M, Nathan N, Wolfenden L, Lubans DR, et al. Scale-up of the physical activity 4 everyone (PA4E1) intervention in secondary schools: 12-month implementation outcomes from a cluster randomized controlled trial. *Int J Behav Nutr Phys Act.* 2020;17(1):100.
 35. Haerens L, Deforche B, Maes L, Cardon G, Stevens V, De Bourdeaudhuij I. Evaluation of a 2-year physical activity and healthy eating intervention in middle school children. *Health Educ Res.* 2006;21(6):911–21.
 36. Harrington DM, Davies MJ, Bodicoat DH, Charles JM, Chudasama YV, Gorely T, et al. Effectiveness of the 'Girls active' school-based physical activity programme: a cluster randomized controlled trial. *Int J Behav Nutr Phys Act.* 2018;15(1):40.
 37. Leme ACB, Baranowski T, Thompson D, Nicklas T, Philippi ST. Sustained impact of the Healthy Habits, Healthy Girls - Brazil school-based randomized controlled trial for adolescents living in low-income communities. *Prev Med Rep.* 2018;10((Thompson D.) Department of Pediatrics, USDA/Children's Nutrition Research Center, Baylor College of Medicine, Houston, TX, United States):346–52.
 38. Okely AD, Lubans DR, Morgan PJ, Cotton W, Peralta L, Miller J, et al. Promoting physical activity among adolescent girls: the girls in Sport group randomized trial. *Int J Behav Nutr Phys Act.* 2017;14(1):81.
 39. Sebire SJ, Jago R, Banfield K, Edwards MJ, Campbell R, Kipping R, et al. Results of a feasibility cluster randomized controlled trial of a peer-led school-based intervention to increase the physical activity of adolescent girls (PLAN-A). *Int J Behav Nutr Phys Act.* 2018;15(1):50.
 40. Schofield L, Mummery WK, Schofield G. Effects of a controlled pedometer-intervention trial for low-active adolescent girls. *Med Sci Sports Exerc.* 2005;37(8):1414–20.
 41. Murphy J, Sweeney MR, McGrane B. The effect of a games-based intervention on wellbeing in adolescent girls. *Health Educ J.* 2022 2022/06/01;81(4):463–78.
 42. Bell SL, Audrey S, Cooper AR, Noble S, Campbell R. Lessons from a peer-led obesity prevention programme in English schools. *Health Promot Int.* 2017;32(2):250–9.
 43. Kariippanon KE, Cliff DP, Okely AD, Parrish AM. Flexible learning spaces reduce sedentary time in adolescents. *J Sci Med Sport.* 2019;22(8):918–23.
 44. Torbeyns T, de Geus B, Bailey S, Decroix L, Van Cutsem J, De Pauw K, et al. Bike desks in the Classroom: Energy Expenditure, Physical Health, Cognitive Performance, Brain Functioning, and academic performance. *J Phys Act Health.* 2017;14(6):429–39.
 45. Sutherland R, Campbell E, McLaughlin M, Nathan N, Wolfenden L, Lubans DR, et al. Scale-up of the physical activity 4 everyone (PA4E1) intervention in secondary schools: 24-month implementation and cost outcomes from a cluster randomized controlled trial. *Int J Behav Nutr Phys Act.* 2021;18(1):137.
 46. Costigan SA, Ridgers ND, Eather N, Plotnikoff RC, Harris N, Lubans DR. Exploring the impact of high intensity interval training on adolescents' objectively measured physical activity: findings from a randomized controlled trial. *J Sports Sci.* 2018;36(10):1087–94.
 47. Subramanian SK, Sharma VK, Arunachalam V, Radhakrishnan K, Ramamurthy S. Effect of structured and unstructured physical activity training on cognitive functions in adolescents - a Randomized Control Trial. *J Clin Diagn Res.* 2015;9(11):CC04–9.
 48. James ML, Christian D, Scott SC, Todd CE, Stratton G, Demmler J, et al. Active children through individual vouchers evaluation: a mixed-method RCT. *Am J Prev Med.* 2020;58(2):232–43.
 49. Bogart LM, Elliott MN, Cowgill BO, Klein DJ, Hawes-Dawson J, Uyeda K et al. Two-year BMI outcomes from a School-based intervention for Nutrition and Exercise: a Randomized Trial. *Pediatrics.* 2016;137(5).
 50. Lubans DR, Morgan PJ, Callister R. Potential moderators and mediators of intervention effects in an obesity prevention program for adolescent boys from disadvantaged schools. *J Sci Med Sport.* 2012;15(6):519–25.
 51. Corder K, Brown HE, Schiff A, van Sluijs EM. Feasibility study and pilot cluster-randomized controlled trial of the GoActive intervention aiming to promote physical activity among adolescents: outcomes and lessons learnt. *BMJ Open.* 2016;6(11):e012335.
 52. Budde H, Voelcker-Rehage C, Pietrassyk-Kendziorra S, Machado S, Ribeiro P, Arafat AM. Steroid hormones in the saliva of adolescents after different exercise intensities and their influence on working memory in a school setting. *Psychoneuroendocrinology.* 2010;35(3):382–91.
 53. Cui Z, Shah S, Yan L, Pan Y, Gao A, Shi X et al. Effect of a school-based peer education intervention on physical activity and sedentary behaviour in Chinese adolescents: a pilot study. *BMJ Open.* 2012;2(3).
 54. Altunkurek SZ, Bebis H. The effects of wellness coaching on the wellness and health behaviors of early adolescents. *Public Health Nurs.* 2019;36(4):488–97.
 55. Aceves-Martins M, Llaurado E, Tarro L, Morina D, Papell-Garcia I, Prades-Tena J, et al. A School-Based, Peer-Led, Social Marketing intervention to engage

- Spanish adolescents in a healthy lifestyle (we are cool-Som La Pera Study): a parallel-cluster Randomized Controlled Study. *Child Obes.* 2017;13(4):300–13.
56. Corepal R, Best P, O'Neill R, Kee F, Badham J, Dunne L, et al. A feasibility study of 'The StepSmart Challenge' to promote physical activity in adolescents. *Pilot Feasibility Stud.* 2019;5(1):132.
57. Parrish AM, Trost SG, Howard SJ, Batterham M, Cliff D, Salmon J, et al. Evaluation of an intervention to reduce adolescent sitting time during the school day: the 'Stand up for Health' randomized controlled trial. *J Sci Med Sport.* 2018;21(12):1244–9.
58. Tymms PB, Curtis SE, Routen AC, Thomson KH, Bolden DS, Bock S, et al. Clustered randomized controlled trial of two education interventions designed to increase physical activity and wellbeing of secondary school students: the MOVE project. *BMJ Open.* 2016;6(1):e009318.
59. Dewar DL, Morgan PJ, Plotnikoff RC, Okely AD, Collins CE, Batterham M, et al. The nutrition and enjoyable activity for teen girls study: a cluster randomized controlled trial. *Am J Prev Med.* 2013;45(3):313–7.
60. Hollis JL, Sutherland R, Campbell L, Morgan PJ, Lubans DR, Nathan N, et al. Effects of a 'school-based' physical activity intervention on adiposity in adolescents from economically disadvantaged communities: secondary outcomes of the 'Physical activity 4 everyone' RCT. *Int J Obes (Lond).* 2016;40(10):1486–93.
61. Suchert V, Isensee B, Sargent J, Weisser B, Hanewinkel R, Lauf G, et al. Prospective effects of pedometer use and class competitions on physical activity in youth: a cluster-randomized controlled trial. *Prev Med.* 2015;81:399–404.
62. Ludyga S, Puhse U, Lucchi S, Marti J, Gerber M. Immediate and sustained effects of intermittent exercise on inhibitory control and task-related heart rate variability in adolescents. *J Sci Med Sport.* 2019;22(1):96–100.
63. Barbosa Filho VC, da Silva KS, Mota J, Beck C, da Silva Lopes A. A physical activity intervention for Brazilian students from low Human Development Index areas: a cluster-randomized controlled trial. *J Phys Act Health.* 2016;13(11):1174–82.
64. Ludyga S, Gerber M, Kamijo K, Brand S, Puhse U. The effects of a school-based exercise program on neurophysiological indices of working memory operations in adolescents. *J Sci Med Sport.* 2018;21(8):833–8.
65. Kolle E, Solberg RB, Safvenbom R, Dyrstad SM, Berntsen S, Resaland GK, et al. The effect of a school-based intervention on physical activity, cardiorespiratory fitness and muscle strength: the School in Motion Cluster randomized trial. *Int J Behav Nutr Phys Act.* 2020;17(1):154.
66. Mavilidi MF, Mason C, Leahy AA, Kennedy SG, Eather N, Hillman CH, et al. Effect of a time-efficient physical activity intervention on Senior School Students' On-Task Behaviour and subjective vitality: the 'Burn 2 learn' Cluster Randomised Controlled Trial. *Educ Psychol Rev.* 2021;33(1):299–323.
67. Melnyk BM, Jacobson D, Kelly S, Belyea M, Shaibi G, Small L, et al. Promoting healthy lifestyles in high school adolescents: a randomized controlled trial. *Am J Prev Med.* 2013;45(4):407–15.
68. Knebel MTG, Borgatto AF, Lopes MVV, Dos Santos PC, Matias TS, Narciso FV, et al. Mediating role of screen media use on adolescents' total sleep time: a cluster-randomized controlled trial for physical activity and sedentary behaviour. *Child Care Health Dev.* 2020;46(3):381–9.
69. Lubans DR, Smith JJ, Morgan PJ, Beauchamp MR, Miller A, Lonsdale C, et al. Mediators of psychological wellbeing in adolescent boys. *J Adolesc Health.* 2016;58(2):230–6.
70. Andrade S, Lachat C, Ochoa-Aviles A, Verstraeten R, Huybregts L, Roberfroid D, et al. A school-based intervention improves physical fitness in Ecuadorian adolescents: a cluster-randomized controlled trial. *Int J Behav Nutr Phys Act.* 2014;11:153.
71. Gammon C, Morton K, Atkin A, Corder K, Daly-Smith A, Quarmby T, et al. Introducing physically active lessons in UK secondary schools: feasibility study and pilot cluster-randomized controlled trial. *BMJ Open.* 2019;9(5):e025080.
72. Tarp J, Domazet SL, Froberg K, Hillman CH, Andersen LB, Bugge A. Effectiveness of a School-based physical activity intervention on cognitive performance in Danish adolescents: LCoMotion-Learning, Cognition and Motion - A Cluster Randomized Controlled Trial. *PLoS ONE.* 2016;11(6):e0158087.
73. Corder K, Sharp SJ, Jong ST, Foubister C, Brown HE, Wells EK, et al. Effectiveness and cost-effectiveness of the GoActive intervention to increase physical activity among UK adolescents: a cluster randomized controlled trial. *PLoS Med.* 2020;17(7):e1003210.
74. Verloigne M, Ridgers ND, De Bourdeaudhuij I, Cardon G. Effect and process evaluation of implementing standing desks in primary and secondary schools in Belgium: a cluster-randomized controlled trial. *Int J Behav Nutr Phys Act.* 2018;15(1):94.
75. van Woudenberg TJ, Bevelander KE, Burk WJ, Smit CR, Buijs L, Buijzen M. A randomized controlled trial testing a social network intervention to promote physical activity among adolescents. *BMC Public Health.* 2018;18(1):542.
76. Carlin A, Murphy MH, Nevill A, Gallagher AM. Effects of a peer-led walking in Schools intervention (the WISH study) on physical activity levels of adolescent girls: a cluster randomized pilot study. *Trials.* 2018;19(1):31.
77. Kennedy SG, Smith JJ, Morgan PJ, Peralta LR, Hilland TA, Eather N, et al. Implementing resistance training in secondary schools: a Cluster Randomized Controlled Trial. *Med Sci Sports Exerc.* 2018;50(1):62–72.
78. Solberg RB, Steene-Johannessen J, Anderssen SA, Ekelund U, Safvenbom R, Haugen T, et al. Effects of a school-based physical activity intervention on academic performance in 14-year old adolescents: a cluster randomized controlled trial - the School in Motion study. *BMC Public Health.* 2021;21(1):871.
79. Sudholz B, Timperio A, Ridgers ND, Dunstan DW, Baldock R, Holland B, et al. The impact and feasibility of introducing height-adjustable desks on adolescents' sitting in a secondary School Classroom. *AIMS Public Health.* 2016;3(2):274–87.
80. Contardo Ayala AM, Sudholz B, Salmon J, Dunstan DW, Ridgers ND, Arundell L, et al. The impact of height-adjustable desks and prompts to break-up classroom sitting on adolescents' energy expenditure, adiposity markers and perceived musculoskeletal discomfort. *PLoS ONE.* 2018;13(9):e0203938.
81. Ghammam R, Maatoug J, Zammit N, Kebaili R, Boughammoura L, Al'Absi M, et al. Long term effect of a school based intervention to prevent chronic diseases in Tunisia, 2009–2015. *Afr Health Sci.* 2017;17(4):1137–48.
82. Yli-Piipari S, Layne T, McCollins T, Knox T. The Impact of Classroom Physical Activity Breaks on Middle School Students' Health-Related Fitness: An Xbox One Kinetic Delivered 4-Week Randomized Controlled Trial. *JTRM in Kinesiology*; 2016.
83. Yang Y, Kang B, Lee EY, Yang HK, Kim HS, Lim SY, et al. Effect of an obesity prevention program focused on motivating environments in childhood: a school-based prospective study. *Int J Obes (Lond).* 2017;41(7):1027–34.
84. Knox GJ, Baker JS, Davies B, Rees A, Morgan K, Cooper SM, et al. Effects of a novel school-based cross-curricular physical activity intervention on cardiovascular disease risk factors in 11- to 14-year-olds: the activity knowledge circuit. *Am J Health Promot.* 2012 Nov-Dec;27(2):75–83.
85. Chen S, Rosenkranz RR, McLoughlin GM, Vazou S, Lanningham-Foster L, Gentile DA et al. Evaluating the implementation and effectiveness of the SWITCH-MS: an Ecological, multi-component adolescent obesity Prevention intervention. *Int J Environ Res Public Health.* 2020;17(15).
86. Haapala HL, Hirvensalo MH, Kulmala J, Hakonen H, Kankaanpaa A, Laine K, et al. Changes in physical activity and sedentary time in the Finnish schools on the move program: a quasi-experimental study. *Scand J Med Sci Sports.* 2017;27(11):1442–53.
87. Bonhauser M, Fernandez G, Puschel K, Yanez F, Montero J, Thompson B, et al. Improving physical fitness and emotional wellbeing in adolescents of low socioeconomic status in Chile: results of a school-based controlled trial. *Health Promot Int.* 2005;20(2):113–22.
88. Louise Bush P, Laberge S, Laforest S. Physical activity promotion among underserved adolescents: make it fun, easy, and popular. *Health Promot Pract.* 2010;11(3 Suppl):795–875.
89. Ardic A, Erdogan S. The effectiveness of the COPE healthy lifestyles TEEN program: a school-based intervention in middle school adolescents with 12-month follow-up. *J Adv Nurs.* 2017;73(6):1377–89.
90. Lazorick S, Fang X, Hardison GT, Crawford Y. Improved body Mass Index measures following a Middle School-based obesity intervention-the MATCH Program. *J Sch Health.* 2015;85(10):680–7.
91. Yu H, Kulinna PH, Mulhearn SC. The effectiveness of equipment provisions on rural Middle School Students' physical activity during lunch recess. *J Phys Act Health.* 2021;18(3):287–95.
92. Kennedy SG, Smith JJ, Estabrooks PA, Nathan N, Noetel M, Morgan PJ, et al. Evaluating the reach, effectiveness, adoption, implementation and maintenance of the resistance training for teens program. *Int J Behav Nutr Phys Act.* 2021;18(1):122.
93. Lubans DR, Sheaman C, Callister R. Exercise adherence and intervention effects of two school-based resistance training programs for adolescents. *Prev Med* 2010 Jan-Feb;50(1–2):56–62.
94. Amoah J, Said S, Rampal L, Manaf R, Ibrahim N, Owusu-Agyei S, et al. Effects of a school-based intervention to reduce cardiovascular disease risk factors among secondary school students: a cluster-randomized, controlled trial. *PLoS ONE.* 2021;16(11):e0259581.

95. Ahmed KR, Kolbe-Alexander T, Khan A. Effectiveness of a school-based intervention on physical activity and screen time among adolescents. *J Sci Med Sport*. 2022;25(3):242–8.
96. Andrade S, Verloigne M, Cardon G, Kolsteren P, Ochoa-Aviles A, Verstraeten R, et al. School-based intervention on healthy behaviour among Ecuadorian adolescents: effect of a cluster-randomized controlled trial on screen-time. *BMC Public Health*. 2015;15(1):942.
97. Barbosa Filho VC, Bandeira ADS, Minatto G, Linard JG, Silva JAD, Costa RMD, et al. Effect of a Multicomponent intervention on lifestyle factors among Brazilian adolescents from low Human Development Index areas: a cluster-randomized controlled trial. *Int J Environ Res Public Health*. 2019;16(2):267.
98. Haerens L, De Bourdeaudhuij I, Maes L, Cardon G, Deforche B. School-based randomized controlled trial of a physical activity intervention among adolescents. *J Adolesc Health*. 2007;40(3):258–65.
99. Lubans DR, Smith JJ, Eather N, Leahy AA, Morgan PJ, Lonsdale C et al. Time-efficient intervention to improve older adolescents' cardiorespiratory fitness: findings from the 'Burn 2 Learn' cluster randomized controlled trial. *Br J Sports Med*. 2020 Dec 21.
100. Sudholz B, Contardo Ayala AM, Timperio A, Dunstan DW, Conroy DE, Abbott G et al. The impact of height-adjustable desks and classroom prompts on classroom sitting time, social, and motivational factors among adolescents. *J Sport Health Sci*. 2020 May 20.
101. Sutherland R, Reeves P, Campbell E, Lubans DR, Morgan PJ, Nathan N, et al. Cost effectiveness of a multi-component school-based physical activity intervention targeting adolescents: the 'Physical activity 4 everyone' cluster randomized trial. *Int J Behav Nutr Phys Act*. 2016;13:94.
102. Sutherland RL, Campbell EM, Lubans DR, Morgan PJ, Nathan NK, Wolfenden L, et al. The physical activity 4 everyone cluster Randomized Trial: 2-Year outcomes of a School Physical activity intervention among adolescents. *Am J Prev Med*. 2016;51(2):195–205.
103. Dewar DL, Morgan PJ, Plotnikoff RC, Okely AD, Batterham M, Lubans DR. Exploring changes in physical activity, sedentary behaviors and hypothesized mediators in the NEAT girls group randomized controlled trial. *J Sci Med Sport*. 2014;17(1):39–46.
104. Lubans DR, Morgan PJ, Aguiar EJ, Callister R. Randomized controlled trial of the physical activity leaders (PALs) program for adolescent boys from disadvantaged secondary schools. *Prev Med*. 2011;52(3/4):239–46.
105. Lubans DR, Morgan PJ, Okely AD, Dewar D, Collins CE, Batterham M, et al. Preventing obesity among adolescent girls: one-year outcomes of the Nutrition and enjoyable activity for Teen girls (NEAT girls) Cluster Randomized Controlled Trial. *Arch Pediatr Adolesc Med*. 2012;166(9):821–7.
106. Smith JJ, Morgan PJ, Plotnikoff RC, Dally KA, Salmon J, Okely AD, et al. Smartphone obesity prevention trial for adolescent boys in low-income communities: the ATLAS RCT. *Pediatrics*. 2014;134(3):e723–31.
107. Leme AC, Lubans DR, Guerra PH, Dewar D, Toassa EC, Philippi ST. Preventing obesity among Brazilian adolescent girls: six-month outcomes of the Healthy habits, Healthy Girls-Brazil school-based randomized controlled trial. *Prev Med*. 2016;86:77–83.
108. Lubans DR, Smith JJ, Plotnikoff RC, Dally KA, Okely AD, Salmon J, et al. Assessing the sustained impact of a school-based obesity prevention program for adolescent boys: the ATLAS Cluster randomized controlled trial. *Int J Behav Nutr Phys Act*. 2016;13:92.
109. Bandeira ADS, Silva KS, Bastos JLD, Silva DAS, Lopes ADS, Barbosa Filho VC. Psychosocial mediators of screen time reduction after an intervention for students from schools in vulnerable areas: a cluster-randomized controlled trial. *J Sci Med Sport*. 2020;23(3):264–9.
110. Laberge S, Bush PL, Chagnon M. Effects of a culturally tailored physical activity promotion program on selected self-regulation skills and attitudes in adolescents of an underserved, multiethnic milieu. *Am J Health Promot*. 2012;26(4):e105–15.
111. Melnyk BM, Jacobson D, Kelly SA, Belyea MJ, Shaibi GQ, Small L, et al. Twelve-Month effects of the COPE Healthy lifestyles TEEN Program on overweight and depressive symptoms in High School adolescents. *J Sch Health*. 2015;85(12):861–70.
112. Lubans DR, Morgan PJ, Dewar D, Collins CE, Plotnikoff RC, Okely AD, et al. The Nutrition and enjoyable activity for Teen girls (NEAT girls) randomized controlled trial for adolescent girls from disadvantaged secondary schools: rationale, study protocol, and baseline results. *BMC Public Health*. 2010;10(1):652.
113. Valkenborghs SR, Hillman CH, Al-ledani O, Nilsson M, Smith JJ, Leahy AA, et al. Effect of high-intensity interval training on hippocampal metabolism in older adolescents. *Psychophysiology*. 2022;59(11):e14090.
114. Morgan PJ, Saunders KL, Lubans DR. Improving physical self-perception in adolescent boys from disadvantaged schools: psychological outcomes from the physical activity leaders randomized controlled trial. *Pediatr Obes*. 2012;7(3):e27–32.
115. Gorely T, Harrington DM, Bodicoat DH, Davies MJ, Khunti K, Sherar LB, et al. Process evaluation of the school-based girls active programme. *BMC Public Health*. 2019;19(1):1187.
116. Sebire SJ, Banfield K, Jago R, Edwards MJ, Campbell R, Kipping R, et al. A process evaluation of the PLAN-A intervention (peer-Led physical activity iNtervention for adolescent girls). *BMC Public Health*. 2019;19(1):1203.
117. Kennedy SG, Peralta LR, Lubans DR, Foweather L, Smith JJ. Implementing a school-based physical activity program: process evaluation and impact on teachers' confidence, perceived barriers and self-perceptions. *Phys Educ Sport Ped*. 2019;24(3):233–48.
118. Lubans D, Morgan P, Okely A, Dewar D, Collins C, Batterham M, et al. Preventing obesity among adolescent girls: outcomes of the nutrition and enjoyable activity for teen girls cluster randomized controlled trial. *J Sci Med Sport*. 2012;15:332–5.
119. Messing S, Rutten A, Abu-Omar K, Ungerer-Rohrich U, Goodwin L, Burlacu I et al. How can physical activity be promoted among children and adolescents? A systematic review of reviews across settings. *Front Public Health*. 2019 2019;7(55):55.
120. Parrish AM, Chong KH, Moriarty AL, Batterham M, Ridgers ND. Interventions to Change School recess activity levels in children and adolescents: a systematic review and Meta-analysis. *Sports Med*. 2020;50(12):2145–73.
121. Norris E, van Steen T, Direito A, Stamatakis E. Physically active lessons in schools and their impact on physical activity, educational, health and cognition outcomes: a systematic review and meta-analysis. *Br J Sports Med*. 2020;54(14):826–38.
122. Ridgers ND, Salmon J, Ridley K, O'Connell E, Arundell L, Timperio A. Agreement between activPAL and ActiGraph for assessing children's sedentary time. *Int J Behav Nutr Phys Act*. 2012;9(1):15.
123. Cassar S, Salmon J, Timperio A, Naylor PJ, van Nassau F, Contardo Ayala AM, et al. Adoption, implementation and sustainability of school-based physical activity and sedentary behaviour interventions in real-world settings: a systematic review. *Int J Behav Nutr Phys Act*. 2019;16(1):120.
124. Beets MW, Okely A, Weaver RG, Webster C, Lubans D, Brusseau T, et al. The theory of expanded, extended, and enhanced opportunities for youth physical activity promotion. *Int J Behav Nutr Phys Act*. 2016;13(1):120.
125. Barbosa Filho VC, da Silva KS, Mota J, Vieira NFC, Gubert FA, Lopes AdS. For whom was it effective? Moderators of the effect of a school-based intervention on potential physical activity determinants among Brazilian students. *Prev Med*. 2017;97:80–5.
126. Bush PL, Laberge S, Laforest S. Physical activity promotion among underserved adolescents: 'make it fun, easy, and popular'. *Health Promot Pract*. 2010;11(3S):79S–87S.
127. Sutherland R, Campbell E, Lubans DR, Morgan PJ, Okely AD, Nathan N, et al. Physical activity 4 everyone' school-based intervention to prevent decline in adolescent physical activity levels: 12 month (mid-intervention) report on a cluster randomized trial. *Br J Sports Med*. 2016;50(8):488–95.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.